

Home

2. Ruxolitinib

Ruxolitinib

Generic name: ruxolitinib [RUX-oh-LI-ti-nib]

Brand name: Jakafi

Dosage form: oral tablet (10 mg; 15 mg; 20 mg; 25 mg; 5 mg)

Drug class: Multikinase inhibitors



Medically reviewed by Drugs.com on Feb 20, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is ruxolitinib?

Ruxolitinib is used in adults to treat myelofibrosis or polycythemia vera, which are bone marrow disorders that affect your body's ability to produce blood cells.

Ruxolitinib is also used to treat graft versus host disease in adults and children at least 12 years old.

Ruxolitinib is usually given after other treatments have failed.

Ruxolitinib may also be used for purposes not listed in this medication guide.

Ruxolitinib side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Some side effects may be similar to the symptoms of myelofibrosis. Call your doctor at once if you have:

- changes in the size, shape, or color of a mole or skin lesion;
- problems with speech, thought, vision, or muscle movement (these symptoms may start gradually and get worse quickly);
- nausea, vomiting, weakness, general ill feeling;
- cold sores around your mouth, skin sores or blisters, itching, tingling, skin rash, burning pain in your thigh or lower back;
- pain in your arms, back, neck, jaw, or stomach;
- signs of a stroke--sudden numbness or weakness, severe headache, slurred speech, problems with vision or balance;
- signs of a blood clot in the lung--chest pain, sudden cough or shortness of breath, dizziness, coughing up blood;

- signs of a blood clot deep in the body--pain, swelling, or warmth in one leg;
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- signs of infection--fever, chills, sore throat, body aches, unusual tiredness, loss of appetite, bruising or bleeding;
- low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath; or
- signs of tuberculosis: fever, cough, night sweats, loss of appetite, weight loss, and feeling very tired.

Common side effects of ruxolitinib may include:

- low blood cell counts;
- · swelling;
- infection;
- · bruising;
- · diarrhea;
- dizziness; or
- headache.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Ruxolitinib side effects (more detail)

Related/similar drugs

Imbruvica

Imbruvica (ibrutinib) is used to treat chronic lymphocytic leukemia (CLL), small lymphocytic ...

Reviews & ratings

6.4 / 10

66 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Jakafi

Jakafi (ruxolitinib) is used to treat myelofibrosis, polycythemia vera, and graft versus host ...

Reviews & ratings

7.5 / 10

33 Reviews

View more

Gleevec

Gleevec is targeted cancer therapy for specific types of leukemia (blood cancer), bone marrow ...

Reviews & ratings

8.6 / 10

37 Reviews

View more

Imatinib

Imatinib is used to treat certain types of leukemia, bone marrow disorders, and skin cancer ...

Reviews & ratings

8.8 / 10

47 Reviews

View more

Ibrutinib

Ibrutinib (Imbruvica) is a Bruton's tyrosine kinase inhibitor used for certain blood cancers and ...

Reviews & ratings

6.8 / 10

83 Reviews

View more

Rayos

Rayos (prednisone, delayed-release) is used to treat rheumatoid arthritis, polymyalgia rheumatica ...

Reviews & ratings

3 Reviews

View more

Warnings

Follow all directions on your prescription label and read all medication guides or instruction sheets. Tell your doctor if you use other medicines or have other medical conditions or allergies.

Before taking this medicine

You should not use ruxolitinib if you are allergic to it.

Tell your doctor if you have ever had tuberculosis or if anyone in your household has tuberculosis. Also, tell your doctor if you've had or been exposed to tuberculosis, or if you recently traveled. Some infections are more common in certain parts of the world, and you may have been exposed during travel.

Tell your doctor if you have or have ever had:

- · any type of infection;
- shingles (herpes zoster);
- a blood clot, stroke, heart attack, or other heart problems;
- · low white or red blood cell counts;
- any type of cancer;
- are a current or past smoker;
- kidney disease (or if you are on dialysis);
- liver disease (especially hepatitis B); or
- high cholesterol or triglycerides (types of fat in the blood).

It is not known if ruxolitinib will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

Do not breastfeed while using ruxolitinib, and for at least 2 weeks after your last dose.

Ruxolitinib is not approved for the treatment of myelofibrosis or polycythemia vera by anyone younger than 18 years old.

How should I take ruxolitinib?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine

exactly as directed.

Your doctor may occasionally change your dose. You will need frequent medical tests to help your doctor determine the best dose.

You may take ruxolitinib with or without food. The ruxolitinib tablet can also be given through a **nasogastric (NG) feeding tube.**

Read and carefully follow any Instructions for Use provided with the medicine. Ask your doctor or pharmacist if you do not understand these instructions.

If you are on dialysis, ruxolitinib should be taken after your dialysis.

Using ruxolitinib may increase your risk of developing other cancers. Ask your doctor about this risk.

Store at room temperature away from moisture and heat.

You should not stop using ruxolitinib suddenly. Follow your doctor's instructions about tapering your dose.

You may be given other medications to help prevent infection. Keep taking these medicines for as long as your doctor has prescribed.

Detailed Ruxolitinib dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking ruxolitinib?

Grapefruit may interact with ruxolitinib and cause side effects. Avoid consuming grapefruit products.

What other drugs will affect ruxolitinib?

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which may increase side effects or make the medicines less effective.

Tell your doctor about all your other medicines, especially fluconazole.

This list is not complete and many other drugs may affect ruxolitinib. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Ruxolitinib drug interactions (more detail)

Does ruxolitinib interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

ruxolitinib	
+	
Enter a drug name	Add

Frequently asked questions

• What are the most common skin conditions? (with photos)

More about ruxolitinib

- · Check interactions
- Compare alternatives
- Reviews (42)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: multikinase inhibitors
- Breastfeeding
- En español

Patient resources

• Ruxolitinib (Oral) advanced reading

Other brands

Jakafi

Professional resources

• Ruxolitinib Phosphate monograph

Other brands

Jakafi

Related treatment guides

- Myelofibrosis
- Polycythemia Vera

- · Graft Versus Host Disease
- Myeloproliferative Disorders

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 7.02.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ড Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

7.4 / 10

42 Reviews

Images

Jakafi (ruxolitinib) 10 mg (INCY 10)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×