

Home

2. Sominex

Sominex

Generic name: diphenhydramine [DYE-fen-HYE-dra-meen]

Brand names: Allergy (Diphenhydramine HCI), Allergy Relief (Diphenhydramine HCI), Banophen, Benadryl, Benadryl Allergy, ... show all 41

brands

Drug classes: Anticholinergic antiemetics, Anticholinergic antiparkinson agents, Antihistamines, Miscellaneous anxiolytics, sedatives and hypnotics



Medically reviewed by Drugs.com on Mar 31, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Sominex?

Sominex is an antihistamine that is used to treat sneezing, runny nose, watery eyes, hives, skin rash, itching, and other cold or allergy symptoms.

Sominex is also used to treat motion sickness, to induce sleep, and to treat certain symptoms of Parkinson's disease. This medicine sleep aid medicine is not for use in children younger than 12 years old.

There are many brands and forms of diphenhydramine available. Not all brands are listed on this leaflet.

Sominex may also be used for purposes not listed in this medication guide.

Sominex side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Sominex may cause serious side effects. Stop using Sominex and call your doctor at once if you have:

- severe drowsiness; or
- painful or difficult urination.

Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

Common side effects of Sominex may include:

- · drowsiness:
- dry eyes, blurred vision;
- · dry mouth, nose, or throat;

- · decreased urination;
- · constipation;
- feeling restless or excited (especially in children); or
- day-time drowsiness or "hangover" feeling after night-time use.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



Sominex side effects (more detail)

Related/similar drugs

Ativan

Ativan is a benzodiazepine used to treat anxiety disorders or anxiety associated with depression ...

Reviews & ratings

8.0 / 10

479 Reviews

View more

Benadryl

Benadryl is an antihistamine and is used to treat sneezing, runny nose, itching and other ...

Reviews & ratings

7.0 / 10

79 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Ambien

Ambien (zolpidem) is a prescription sedative used to treat insomnia, helping you fall asleep and ...

Reviews & ratings

7.0 / 10

558 Reviews

View more

Lunesta

Lunesta is a sedative and is used to treat insomnia. Lunesta's effects typically begin within 30 ...

Reviews & ratings

5.9 / 10

341 Reviews

View more

Eszopiclone

Eszopiclone (brand name Lunesta) is a sedative and is used to treat insomnia. Eszopiclone's effects ...

Reviews & ratings

5.7 / 10

442 Reviews

View more

Temazepam

Temazepam is used to treat insomnia (trouble sleeping). Learn about side effects, interactions and ...

Reviews & ratings

6.5 / 10

448 Reviews

View more

Melatonin

Melatonin is used to combat jet lag and ease sleep problems like insomnia.

Learn about uses ...

Reviews & ratings

5.7 / 10

239 Reviews

View more

Zolpidem

Zolpidem is a sedative-hypnotic medicine used for the short-term treatment of insomnia (trouble ...

Reviews & ratings

6.9 / 10

1,055 Reviews

View more

Warnings

Use Sominex exactly as directed. Taking too much Sominex can lead to serious heart problems, seizures, coma, or death.

Do not use this medicine to make a child sleepy. Sominex sleep aid medicine is not for use in children younger than 12 years old.

Before taking this medicine

You should not use Sominex if you are allergic to it.

Ask a doctor or pharmacist if this medicine is safe to use if you have ever had:

- an enlarged prostate or urination problems;
- asthma, chronic obstructive pulmonary disease (COPD), or other breathing disorder;
- glaucoma; or
- · a thyroid disorder.

Ask a doctor before using this medicine if you are pregnant or breastfeeding. Sominex may slow breast milk production.

How should I take Sominex?

Use exactly as directed on the label, or as prescribed by your doctor. Sominex is only for short-term use until your symptoms clear up.

Taking too much Sominex can lead to serious heart problems, seizures, coma, or death.

Always follow directions on the medicine label about giving Sominex to a child. **Do not use the medicine only to make a child sleepy.** Death can occur from the misuse of antihistamines in very young children.

For motion sickness, take Sominex 30 minutes before you will be in a situation that causes you motion sickness (such as a long car ride, airplane or boat travel, amusement park rides, etc). Continue taking this medicine with meals and at bedtime for the rest of the time you will be in a motion-sickness situation.

As a sleep aid, take Sominex within 30 minutes before bedtime.

You must chew the chewable tablet before you swallow it.

Measure **liquid medicine** carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

Remove an **orally disintegrating tablet** from the package only when you are ready to take the medicine. Place the tablet in your mouth and allow it to dissolve, without chewing. Swallow several times as the tablet dissolves.

Call your doctor if the condition you are treating with Sominex does not improve, or if you have a fever with a headache, cough, or skin rash.

Do not use Sominex for longer than 2 weeks to treat sleep problems, or longer than 7 days to treat cold or allergy symptoms.

Sominex can affect the results of allergy skin tests. Tell any doctor who treats you that you are using Sominex.

Store at room temperature away from moisture and heat.

Detailed Sominex dosage information

What happens if I miss a dose?

Since Sominex is used when needed, you may not be on a dosing schedule. Skip any missed dose if it's almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of diphenhydramine** can be fatal.

Overdose symptoms may include vomiting, confusion, severe drowsiness, ringing in your ears, no urination, very dry eyes and mouth, dilated pupils, fast heartbeats, tremor, agitation, hallucinations, or seizure.

What should I avoid while taking Sominex?

Avoid driving or hazardous activity until you know how Sominex will affect you. Your reactions could be impaired.

Drinking alcohol can increase certain side effects of diphenhydramine.

Ask a doctor or pharmacist before using any other medicine that may contain Sominex. This includes medicines for

sleep, cold/allergy symptoms, or anti-itch medicine used on the skin. Using too much this medicine may lead to a fatal overdose.

What other drugs will affect Sominex?

Ask a doctor or pharmacist before using Sominex with any other medicines, especially drugs that can cause drowsiness (such as opioid medication, sleep medicine, a muscle relaxer, or medicine for anxiety or seizures). Tell your doctor about all your current medicines and any medicine you start or stop using. This includes prescription and over-the-counter medicines, vitamins, and herbal products. **Not all possible interactions are listed here.**

Sominex drug interactions (more detail)

Does Sominex interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

What is the max dose of diphenhydramine for adults?	~
What are the strongest sleeping pills?	~
Does Mucinex help with Covid?	~
Does diphenhydramine raise blood pressure?	~
Does Benadryl help with or cause anxiety?	~
Can diphenhydramine be used as a sleep aid?	~
Diphenhydramine Hydrochloride vs Citrate: What's the difference?	~

More about Sominex (diphenhydramine)

- Check interactions
- · Compare alternatives
- Reviews (64)
- Drug images

- Latest FDA alerts (2)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: anticholinergic antiemetics
- Breastfeeding
- En español

Patient resources

Other brands

Benadryl, Banophen, Benadryl Allergy, ZzzQuil, ... +23 more

Professional resources

- DiphenhydrAMINE monograph
- Diphenhydramine Oral Solution (FDA)

Other brands

Benadryl, Benadryl Allergy, Diphen, Dicopanol

Related treatment guides

Insomnia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 9.02.

DRUG STATUS

Availability

Rx and/or OTC

Pregnancy & Lactation

ം Risk data available

CSA Schedule*

User Reviews & Ratings

6.3 / 10

64 Reviews

Images

Sominex diphenhydramine 25 mg (S)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







