

#### Home

Trilafon

## Trilafon ⊲

Generic name: perphenazine [ per-FEN-a-zeen ]

Drug classes: Phenothiazine antiemetics, Phenothiazine antipsychotics



Medically reviewed by Drugs.com on Aug 3, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ



The Trilafon brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be generic equivalents available.

## What is Trilafon?

Trilafon is a phenothiazine (FEEN-oh-THYE-a-zeen) anti-psychotic medicine.

Trilafon is used to treat psychotic disorders such as schizophrenia. It is also used to control severe nausea and vomiting.

Trilafon may also be used for purposes not listed in this medication guide.

## Trilafon side effects

Get emergency medical help if you have any of these **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

High doses or long-term use of perphenazine can cause a serious movement disorder that may not be reversible. The longer you use Trilafon, the more likely you are to develop this disorder, especially if you are a woman or an older adult.

Trilafon may cause serious side effects. Call your doctor at once if you have:

- uncontrolled muscle movements in your arms or legs, or your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement);
- · worsening symptoms of schizophrenia;
- · confusion, paranoia, feeling restless or excited;
- seizure (convulsions);
- a light-headed feeling, like you might pass out;
- jaundice (yellowing of your skin or eyes);

- · little or no urinating;
- slow heart rate, weak pulse, weak or shallow breathing;
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat, cough, trouble breathing, feeling light-headed; or
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, fast or uneven heartbeats.

Side effects such as dry mouth, constipation, tremors, and drowsiness may be more likely in older adults.

### Common side effects of Trilafon may include:

- mild dizziness or drowsiness;
- blurred vision, headache;
- sleep problems (insomnia), strange dreams;
- loss of appetite, vomiting, diarrhea, constipation;
- · increased sweating or urination;
- dry mouth or stuffy nose;
- breast swelling or discharge; or
- mild itching or skin rash.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Trilafon side effects (more detail)

## Related/similar drugs

## **Diphenhydramine**

Diphenhydramine is an antihistamine used to treat sneezing, runny nose, itching, hives and other ...

#### **Reviews & ratings**

5.5 / 10

541 Reviews

View more

## **Promethazine**

Promethazine is a phenothiazine and antihistamine used to treat allergies, motion sickness, nausea ...

### **Reviews & ratings**

#### View more

#### FEATURED

## **Ozempic**

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

### **Reviews & ratings**

6.6 / 10

1,504 Reviews

### View more

### **Zofran**

Zofran (ondansetron) is used to prevent nausea and vomiting caused by surgery or cancer medicines ...

## **Reviews & ratings**

8.0 / 10

270 Reviews

### View more

## **Olanzapine**

Olanzapine (brand name Zyprexa) is an atypical antipsychotic that may be used to treat adults and ...

### **Reviews & ratings**

5.9 / 10

1,091 Reviews

### View more

### **Dexamethasone**

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

### **Reviews & ratings**

5.9 / 10

162 Reviews

#### View more

## Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

### **Reviews & ratings**

7.9 / 10

1,054 Reviews

View more

### **Meclizine**

Meclizine is used to treat or prevent nausea, vomiting, and dizziness caused by motion sickness ...

### **Reviews & ratings**

7.4 / 10

234 Reviews

View more

## Metoclopramide

Metoclopramide is used short-term to treat heartburn caused by gastroesophageal reflux. Learn about

•••

## Reviews & ratings

3.9 / 10

517 Reviews

View more

## **Ativan**

Ativan is a benzodiazepine used to treat anxiety disorders or anxiety associated with depression ...

## Reviews & ratings

8.0 / 10

479 Reviews

View more

## **Warnings**

You should not use Trilafon if you have liver disease, brain damage, bone marrow depression, a blood cell disorder, or if you are also using large amounts of alcohol or medicines that make you sleepy.

Trilafon is not approved for use in older adults with dementia-related psychosis.

Call your doctor at once if you have twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs. These could be early signs of dangerous side effects.

## Before taking this medicine

You should not use Trilafon if you are allergic to any phenothiazine (such as Trilafon, chlorpromazine, prochlorperazine, promethazine, o thioridazine), or if you have:

- · liver disease:
- · brain damage;
- · bone marrow depression;
- a blood cell disorder (such as low platelets or low red or white blood cell counts); or
- if you are also using large amounts of alcohol or medicines that make you sleepy.

Trilafon may increase the risk of death in older adults with dementia-related psychosis and is not approved for this use.

Tell your doctor if you have ever had:

- · severe or untreated depression;
- · heart disease or high blood pressure;
- · kidney disease;
- · severe asthma, emphysema, or other breathing problem;
- · seizures;
- · Parkinson's disease;
- breast cancer;
- adrenal gland tumor (pheochromocytoma);
- enlarged prostate or urination problems;
- low levels of calcium in your blood (hypocalcemia);
- · glaucoma; or
- a serious side effect while using Trilafon or another phenothiazine.

Tell your doctor if you will be exposed to extreme heat or cold, or to insecticide poisons while you are taking Trilafon.

Tell your doctor if you are pregnant or you become pregnant. **Taking antipsychotic medicine in the last 3 months of pregnancy may cause breathing problems, feeding problems, or withdrawal symptoms in the newborn**.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk.

## How should I take Trilafon?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You will need frequent medical tests.

If you need surgery, tell your surgeon you currently use this medicine.

**Do not stop using Trilafon suddenly**, or you could have unpleasant symptoms such as nausea, vomiting, dizziness, or tremors. Ask your doctor how to safely stop using this medicine.

Store at room temperature away from moisture, heat, and light.

Detailed Trilafon dosage information

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of Trilafon can be fatal.** 

## What should I avoid while taking perphenazine?

Avoid driving or hazardous activity until you know how Trilafon will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

Avoid drinking alcohol. It can increase some of the side effects of perphenazine.

Avoid exposure to sunlight or tanning beds. Trilafon can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

## What other drugs will affect Trilafon?

Taking Trilafon with other drugs that make you sleepy or slow your breathing can cause dangerous or lifethreatening side effects. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Many drugs can affect Trilafon. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all your current medicines and any medicine you start or stop using.

Trilafon drug interactions (more detail)

## Does Trilafon interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



## Frequently asked questions

• What drugs cause tardive dyskinesia?

## More about Trilafon (perphenazine)

- Check interactions
- Compare alternatives
- Reviews (9)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Drug class: phenothiazine antiemetics
- Breastfeeding

### **Professional resources**

- Perphenazine monograph
- Perphenazine (FDA)

## Related treatment guides

- Nausea/Vomiting
- Psychosis

## **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.01.

#### **DRUG STATUS**

### **Availability**

O Discontinued

## **Pregnancy & Lactation**

ഴെ Risk data available

### **CSA Schedule\***

N/A Not a controlled drug

## **Approval History**

10+ years FDA approved 1957

## **User Reviews & Ratings**

6.0 / 10

9 Reviews

### **Images**

Trilafon 16 MG (LOGO 077)



# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com Advertising policy Attribution & citations

### **Terms & privacy**

Terms of use Editorial policy Privacy policy

**Support** 

Help center Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

Х