

Home

Migranal

Migranal (nasal)

Generic name: dihydroergotamine (nasal) [dye-HYE-droe-er-GOT-a-meen]

Brand names: Migranal, Trudhesa Drug class: Antimigraine agents



Medically reviewed by Drugs.com on Jul 20, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Migranal nasal?

Migranal is used to treat migraine headaches with or without aura in adults. This medicine nasal **will only treat** a headache that has already begun. It **will not prevent** headaches or reduce the number of attacks.

Migranal should not be used to treat any headache that seems to be different from your usual migraine headaches.

Migranal may also be used for purposes not listed in this medication guide.

Migranal nasal side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Migranal may cause serious side effects. Call your doctor at once if you have:

- numbness, cold feeling, or pale appearance of your fingers or toes;
- sudden or severe stomach pain, weight loss, nausea, vomiting, constipation, bloody diarrhea, fever;
- severe headache, blurred vision, pounding in your neck or ears;
- burning sensation in the nose, mouth, throat, and altered sense of taste;
- fast or slow heart rate;
- · swelling or itching;
- signs of a blood clot--sudden numbness or weakness on one side of the body, chest pain, problems with vision or speech, pain or swelling in one leg; or
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating.

Common side effects of Migranal may include:

• pain, bruising, swelling, or irritation where the medicine was applied;

- · drowsiness;
- · dizziness;
- nausea, vomiting, diarrhea;
- · runny nose;
- altered sense of taste; or
- · sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Migranal side effects (more detail)

Related/similar drugs

Metoclopramide

Metoclopramide is used short-term to treat heartburn caused by gastroesophageal reflux. Learn about

Reviews & ratings

3.9 / 10

517 Reviews

View more

FEATURED

Nurtec ODT

Nurtec ODT (rimegepant) is used to treat acute migraines and prevent episodic migraines, by ...

Reviews & ratings

6.3 / 10

320 Reviews

View more

Celecoxib

Celecoxib is a nonsteroidal antiinflammatory drug (NSAID) used to treat pain or inflammation ...

Reviews & ratings

7.0 / 10

280 Reviews

View more

Diclofenac

Diclofenac is a nonsteroidal antiinflammatory drug used to treat pain and inflammation associated ...

Reviews & ratings

7.3 / 10

695 Reviews

View more

Propranolol

Propranolol is a beta-blocker that is used to treat tremors, chest pain, high blood pressure, heart ...

Reviews & ratings

8.2 / 10

1,376 Reviews

View more

Amitriptyline

Amitriptyline is a tricyclic antidepressant used to treat depression and, off-label, conditions ...

Reviews & ratings

7.5 / 10

1,803 Reviews

View more

Ibuprofen

Ibuprofen (Advil, Motrin) treats minor aches and pains caused by the common cold, headaches ...

Reviews & ratings

7.3 / 10

240 Reviews

View more

Prednisone

Prednisone is used to treat allergic disorders, ulcerative colitis, psoriasis and arthritis. Learn ...

Reviews & ratings

7.5 / 10

867 Reviews

View more

Warnings

Tell your doctor about all your other medicines. Some drugs should not be used with Migranal.

Before taking this medicine

You should not use Migranal if you are allergic to it or similar medicines (ergotamine, ergonovine, methylergonovine, and others), or if you have or have ever had:

- heart disease, angina (chest pain), blood circulation problems, coronary artery disease (clogged arteries), or a heart attack or stroke;
- uncontrolled high blood pressure;
- · a serious infection called sepsis;
- · vascular surgery;
- narrowing of blood vessels in the legs, arms, stomach, or kidneys (peripheral vascular disease); or
- · severe liver or kidney disease.

Some drugs should not be used with Migranal. Your treatment plan may change if you also use:

- medicines that raise your blood pressure or constrict your blood vessels;
- an antibiotic such as troleandomycin, clarithromycin, or erythromycin;
- antifungal medicine such as itraconazole or ketoconazole; or
- antiviral medicine for HIV or hepatitis C.

Tell your doctor if you have ever had:

- blood circulation problems in your arms, hands, legs, or feet;
- chest pain, shortness of breath;
- risk factors for coronary artery disease (high blood pressure, high cholesterol, diabetes, menopause, smoking, family history of coronary artery disease, being overweight, being over 40 and a man); or
- liver or kidney disease.

Migranal may cause preterm labor. Tell your doctor if you are pregnant or plan to become pregnant. **Do not use**Migranal if you are pregnant.

Do not breastfeed while using this medicine, and for at least 3 days after your last dose.

How should I use Migranal nasal?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Migranal spray is absorbed quickly through your nasal passages and is for use only in the nose. The nasal spray liquid should not be injected into the body.

Your first dose may be given in a medical setting where any serious side effects can be quickly treated.

You should use Migranal as soon as your headache begins, but it may be given at any time during the migraine episode.

Before your first use, prime the nasal spray device with 4 sprays into the air, away from your face.

Trudhesa: apply one spray in each nostril then another spray in each nostril 1 hour later only if your headache comes back or if you only get some relief from your headache. Use a new nasal spray device for the second dose.

Migranal: apply one spray in each nostril followed by another spray in each nostril 15 minutes later, for a total of 4 sprays.

You may need medical tests to check your heart function.

The nasal spray vial should be thrown away 8 hours after it has been opened.

Store at room temperature away from moisture, heat, and light. Do not keep this medicine in a refrigerator and do not freeze.

Detailed Migranal dosage information

What happens if I miss a dose?

Migranal nasal is used when needed and does not have a daily dosing schedule.

Do not use more than 2 doses of **Trudhesa** within 24 hours or 3 doses within 7 days.

Do not use more than 6 sprays of Migranal within 24 hours or 8 sprays within 7 days.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include slow breathing, blood pressure changes, confusion, hallucinations, agitation, convulsions, coma, nausea, vomiting, stomach pain, and pain, numbness, cold feeling, or pale appearance of your fingers or toes.

What should I avoid while using Migranal nasal?

Do not use Migranal within 24 hours before or after using another migraine headache medicine, including:

- another ergot medicine such as ergotamine, ergonovine, methylergonovine; or
- almotriptan, eletriptan, frovatriptan, naratriptan, sumatriptan, or rizatriptan.

Grapefruit may interact with Migranal and cause side effects. Avoid consuming grapefruit products.

What other drugs will affect Migranal nasal?

Tell your doctor about all your current medicines. Many drugs can affect Migranal, especially:

- heart or blood pressure medication;
- nicotine gum or patches;
- an antidepressant; or
- medicine to treat asthma.

This list is not complete and many other drugs may affect Migranal. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

i Migranal drug interactions (more detail)

Does Migranal interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Frequently asked questions

• What are the new drugs for the treatment of migraine?

More about Migranal (dihydroergotamine)

- · Check interactions
- Compare alternatives
- Reviews (23)
- Side effects
- Dosage information

- During pregnancy
- · Generic availability
- Drug class: antimigraine agents
- Breastfeeding
- En español

Patient resources

- Migranal advanced reading
- Migranal

Other brands

Trudhesa, D.H.E. 45

Professional resources

- Migranal prescribing information
- Dihydroergotamine (AHFS Monograph)

Other brands

Trudhesa, D.H.E. 45

Related treatment guides

- Cluster Headaches
- Migraine

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

7.7 / 10

23 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





