

#### Home

Amlodipine/olmesartan

# **Amlodipine and olmesartan**

Generic name: amlodipine and olmesartan [ am-LOE-de-peen-ol-me-SAR-tan ]

Brand name: Azor

**Dosage form:** oral tablet (10 mg-20 mg; 10 mg-40 mg; 5 mg-20 mg; 5 mg-40 mg)

Drug class: Angiotensin II inhibitors with calcium channel blockers



Medically reviewed by Drugs.com on Mar 24, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

### What is amlodipine and olmesartan?

Amlodipine is a calcium channel blocker. Olmesartan is an angiotensin II receptor blocker (sometimes called an ARB).

Amlodipine and olmesartan is a combination medicine used to treat high blood pressure (hypertension). Lowering blood pressure may lower your risk of a stroke or heart attack.

Amlodipine and olmesartan may also be used for purposes not listed in this medication guide.

## Amlodipine and olmesartan side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

When you first start taking amlodipine and olmesartan, you may have new or worsening chest pain (angina), or you could have a heart attack. **Seek emergency medical attention or call your doctor right away if you have symptoms such as:** chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating.

Amlodipine and olmesartan may cause serious side effects. Call your doctor at once if you have:

- · new or worsening chest pain;
- swelling in your hands or feet, rapid weight gain;
- severe or ongoing diarrhea with weight loss;
- pounding heartbeats or fluttering in your chest;
- · a light-headed feeling, like you might pass out; or
- high potassium level--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement.

Common side effects include swelling.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

<u>Amlodipine / olmesartan side effects</u> (more detail)

## Related/similar drugs

### **Vasotec**

Vasotec (enalapril) is used to treat high blood pressure, heart failure and kidney problems caused ...

### **Reviews & ratings**

4 Reviews

View more

#### FEATURED

#### **Omvoh**

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

### **Reviews & ratings**

1 Reviews

View more

### Lisinopril

Lisinopril is an ACE inhibitor. It is used to treat high blood pressure, congestive heart failure ...

### **Reviews & ratings**

5.0 / 10

626 Reviews

View more

### **Atenolol**

Atenolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

### **Reviews & ratings**

7.2 / 10

276 Reviews

#### View more

### **Valsartan**

Valsartan is an angiotensin II receptor blocker (ARB) that may be used to treat high blood pressure ...

### Reviews & ratings

5.7 / 10

152 Reviews

View more

### Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

### Reviews & ratings

5.2 / 10

185 Reviews

View more

### Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

### **Reviews & ratings**

6.4 / 10

56 Reviews

View more

### **Furosemide**

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

### **Reviews & ratings**

6.8 / 10

121 Reviews

View more

#### Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

### **Reviews & ratings**

4.5 / 10

578 Reviews

View more

## **Warnings**

Do not use if you are pregnant. If you become pregnant, stop taking this medicine and tell your doctor right away.

If you have diabetes, do not use amlodipine and olmesartan together with any medication that contains aliskiren (a blood pressure medicine).

### Before taking this medicine

You should not use amlodipine and olmesartan if you are allergic to amlodipine or olmesartan.

If you have diabetes, do not use amlodipine and olmesartan together with any medication that contains aliskiren (a blood pressure medicine).

You may also need to avoid taking amlodipine and olmesartan with aliskiren if you have kidney disease.

Tell your doctor if you have ever had:

- · kidney disease;
- liver disease;
- · coronary artery disease;
- angina (chest pain), congestive heart failure;
- high levels of potassium in your blood;
- if you are on a low-salt diet;
- if you are 75 years or older; or
- if you recently had a heart attack.

Do not use if you are pregnant. Stop using the medicine and tell your doctor right away if you become pregnant. Olmesartan can cause injury or death to the unborn baby if you take the medicine during your second or third trimester.

You should not breastfeed while using this medicine.

Amlodipine and olmesartan is not approved for use by anyone younger than 18 years old.

### How should I take amlodipine and olmesartan?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You may take this medication with or without food.

Your blood pressure will need to be checked often. Your kidney function may also need to be checked.

Some things can cause your blood pressure to get too low. This includes vomiting, diarrhea, or heavy sweating. Call your doctor if you are sick with vomiting or diarrhea.

If you need surgery, tell the surgeon ahead of time that you are using amlodipine and olmesartan.

Keep using this medicine as directed, even if you feel well. High blood pressure often has no symptoms. You may need to use blood pressure medication for the rest of your life.

Store at room temperature away from moisture and heat.

### Amlodipine and olmesartan dosing information

### **Usual Adult Dose for Hypertension:**

Initial dose: Amlodipine 5 mg-Olmesartan 20 mg orally once a day; may increase dose after 1 to 2 weeks of therapy.

Maximum dose: Amlodipine 10 mg-Olmesartan 40 mg orally once a day

### Uses:

- -Substitution therapy for patients maintained on the individual components separately or for patients not adequately controlled with one of the components or a component within the same antihypertensive class.
- -Initial therapy in patients who are likely to need more than 1 antihypertensive agent to control blood pressure.
- Detailed Amlodipine / olmesartan dosage information

### What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

### What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

## What should I avoid while taking amlodipine and olmesartan?

Avoid driving or hazardous activity until you know how amlodipine and olmesartan will affect you. Your reactions could be impaired. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

Do not use potassium supplements or salt substitutes, unless your doctor has told you to.

### What other drugs will affect amlodipine and olmesartan?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

If you also take colesevelam, take it 4 hours after you take amlodipine and olmesartan.

Tell your doctor about all your other medicines, especially:

- cyclosporine;
- lithium;
- · simvastatin;
- · tacrolimus; or
- NSAIDs (nonsteroidal anti-inflammatory drugs)--aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others.

This list is not complete. Other drugs may affect amlodipine and olmesartan, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Amlodipine / olmesartan drug interactions (more detail)

### Does amlodipine/olmesartan interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

# Enter a drug name Add

## More about amlodipine / olmesartan

- Check interactions
- Compare alternatives
- · Pricing & coupons
- Reviews (60)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: angiotensin II inhibitors with calcium channel blockers
- En español

### Patient resources

#### Other brands

Azor

### **Professional resources**

• Amlodipine and Olmesartan Medoxomil prescribing information

### Other brands

Azor

### Related treatment guides

• High Blood Pressure

### **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

#### **DRUG STATUS**

**Availability** 

Rx Prescription only

**Pregnancy & Lactation** 

ন্দু Risk data available

**CSA Schedule\*** 

N/A Not a controlled drug

**Approval History** 

□ Drug history at FDA

### **User Reviews & Ratings**

8.4 / 10

60 Reviews

### **Images**



# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





### **About**

About Drugs.com
Advertising policy
Attribution & citations

### Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

#### Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





