

Home

- Alavert D-12 Hour Allerg...
- 3. FDA PI

Alavert D-12 Hour Allergy and Sinus: Package Insert / Prescribing Info

Package insert / product label

Generic name: loratadine, pseudoephedrine sulfate

Dosage form: tablet, film coated, extended release

Drug class: Upper respiratory combinations

Medically reviewed by Drugs.com. Last updated on Feb 27, 2025.

On This Page

Indications and Usage

Warnings

Dosage and Administration

Storage and Handling

ACTIVE INGREDIENTS (IN EACH TABLET)

Loratadine 5 mg

Pseudoephedrine sulfate 120 mg

PURPOSES

Antihistamine

Nasal decongestant

Indications and Usage for Alavert D-12 Hour Allergy and Sinus

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - · itchy, watery eyes
 - runny nose
 - · itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other respiratory allergies
- · reduces swelling of nasal passages

- temporarily relieves sinus congestion and pressure
- · temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Alavert D-12 Hour Allergy and Sinus Dosage and Administration

• do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over 1 tablet every 12 hours; not more than 2 tablets in 24 hours

children under 12 years of age ask a doctor

consumers with liver or kidney disease ask a doctor

Storage and Handling

- each tablet contains: calcium 25 mg
- store between 20° and 25°C (68° and 77°F)
- · keep in a dry place

INACTIVE INGREDIENTS

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

QUESTIONS OR COMMENTS?

Call weekdays from 9 AM to 5 PM EST at 1-800-ALAVERT (1-800-252-8378)

Distributed by: Pfizer, Madison, NJ 07940 USA

PRINCIPAL DISPLAY PANEL

NDC 0573-2660-24

Compare ours to the Active Ingredients in Claritin-D[®] 12 Hour Tablets[†]

allergy & congestion

ALAVERT®

D-12

HOUR

Pseudoephedrine Sulfate 120 mg/Nasal Decongestant Loratadine 5 mg/Antihistamine — Extended Release Tablets

Non-Drowsy*

Relief of:

- Nasal and Sinus Congestion Due to Colds or Allergies
- Sneezing Runny Nose Itchy, Watery Eyes
- Itchy Throat or Nose Due to Allergies

Indoor & Outdoor Allergies

[†]This product is not manufactured by Bayer HealthCare LLC, the distributor of Claritin-D[®] 12 Hour Tablets.

24 TABLETS

Actual Size

*When taken as directed.

See Drug Facts Panel.



sno

for most recent product information, visit www.slavert.com

Distributed by: Pitzer, Madison, NJ 07940 USA © 2018 Pitzer Inc.

Questions or comments? Cal westcays from 9 AM to 5 PM Est alt-800-Alament (1-800-252-8378)

phosphale, hyprometose, actos monohydrate, megmedum stearale, phosphale, hyprometose, atanium dioxide Inactive ingredients coscamelos sodum, disasc calcum Ong Facts (continued)

-	m yeeb in a qui, biace
	■ alone between 20° to 25°C (68° to 77°F)
	errod no resolond at linu redaild it seu fon ob m
	■ each tablet contains: calcium 35 mg
	пойватот і тэль
	 OL MODULA GENERAL BENERAL DESCRIPTION OF THE PROPERTY OF THE P

consumers with liver or lidiney disease
drifteren under 12 years of age
norbilito bras atluba navo bras assay St

Directions ando not divide, crust, chew or dissolve the tablet t

If pragment or brease1-feeding, sak a healin professional before use. Keep out of meath of children. In case of overdoss, gast medical helpor combact as Polson Control Center right away (1-600-2222-1222).

Stop uses and sak a docitor III

an admitted stop deliberation occurs. Seek medical help flyth away

an alympome to brothingrove width a Cales or are accompanied by a Pewr

from the confidences or skeeples ness occurs

en using this product do not take more than directed. Taking more than acted may cause drowsiness.

inver or leaning disease. Your doctor should determine if you need a offin Drug Facts (continued)

over) wey it eeu arohed robede s AAA esseab blovyrit um easea b haeri um eafedelo um erusee or brook right um braeg eteleory begelne ne of eub gritearhu eiduurit um
a thigh blood pressure and abbetes
esses p proving man esses p presid man
event beet on the order of the control of the contr

White the state of the state of

If you have ever had an altergic reaction to this product or any of its

sgninnsW eau fon od

esou aut rigues d'heer breathraidh fir ough frie eson

sasn

Active ingredients (in each tablet)
on to table and the most of tablets) sasodind Drug Facts

EACH TABLET IS BLISTER SEALED. DO NOT USE IF SEAL IS BROKEN.

D-12 HOUR

Pseudoephedrine Sulfate 120mg/Nasal Decongestant Loratadine 5 mg/Anthistamine — Extended Release Tablets

Non-Drowsy*

24 Tablets

*When taken as directed. See Drug Facts Panel.

Compare ours to the Active Ingredients in Claritin-D® 12 Hour Tablets†

allergy & congestion

Actual Size

Pseudoephedrine Sulfate 120 mg/Nasal Decongestant Loratadine 5 mg/Antihistamine — Extended Release Tablets

Non-Drowsy* Relief of:

. Nasal and Sinus Congestion Due to Colds or Allergies

Sneezing Runny Nose Itchy, Watery Eyes

Itchy Throat or Nose Due to Allergies


Indoor & Outdoor Allergies

†This product is not manufactured by Bayer HealthCare LLC, the distributor of Claritin-D® 12 Hour Tablets.

TABLETS

*When taken as directed. See Drug Facts Panel.





ALAVERT ALLERGY SINUS D-12

loratadine, pseudoephedrine sulfate tablet, film coated, extended release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0573-2660

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	5 mg
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE -	PSEUDOEPHEDRINE	120 mg
UNII:7CUC9DDI9F)	SULFATE	

Inactive Ingredients

Ingredient Name Strength

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)

ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)

MAGNESIUM STEARATE (UNII: 70097M6I30)

POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color WHITE (white to off-white) Score no score

Shape ROUND (round shape and an extra deep convex) Size 14mm

Flavor Imprint Code Alavert; D12

Contains

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-2660-12	2 in 1 CARTON	01/30/2004	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0573-2660-24	4 in 1 CARTON	01/30/2004	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076050	01/30/2004	

Labeler - GlaxoSmithKline Consumer Healthcare Holdings (US) LLC (079944263)

More about Alavert D-12 Hour Allergy and Sinus (loratadine / pseudoephedrine)

- · Check interactions
- · Compare alternatives
- Reviews (3)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: upper respiratory combinations
- En español

Patient resources

- Alavert D-12 drug information
- Alavert-D 12-Hour (Advanced Reading)

Professional resources

Other brands

Claritin-D 24 Hour

Other formulations

Alavert

Related treatment guides

- Allergic Rhinitis
- Nasal Congestion

Medical Disclaimer

Claritin Pediatric Dosage Guide

Use these dosage charts for infants and children under 12 years old: Claritin Dosage Charts (Infants & Children)

DRUG STATUS

Availability

OTC Over the counter

Pregnancy & Lactation

ላያ Risk data available

CSA Schedule*

N/A Not a controlled drug

WADA Class

റ്റ് Anti-Doping Classification

User Reviews & Ratings

3 Reviews

Images

Alavert D-12 5 mg / 120 mg (ALAVERT D12)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

X