

Home

2. Indocin SR

Indocin SR

Generic name: indomethacin (oral) [IN-doe-METH-a-sin]

Brand names: Indocin, Indocin SR

Drug class: Nonsteroidal anti-inflammatory drugs



Medically reviewed by Drugs.com on Jun 11, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Indocin SR?

Indocin SR is a nonsteroidal anti-inflammatory drug (NSAID) that is used in adults and children at least 14 years old to treat signs and symptoms caused by certain types of arthritis such as moderate to severe osteoarthritis, rheumatoid arthritis, gouty arthritis, or ankylosing spondylitis.

Indocin SR is also used to treat shoulder pain caused by bursitis or tendonitis.

Extended-release Indocin SR should not be used to treat gouty arthritis.

Indocin SR may also be used for purposes not listed in this medication guide.

Indocin SR side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Stop using Indocin SR and seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes.

Get emergency medical help if you have **signs of a heart attack or stroke:** chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath.

Indocin SR may cause serious side effects. Stop using Indocin SR and call your doctor at once if you have:

- · a skin rash, no matter how mild;
- · changes in your vision;
- · flu-like symptoms;
- neurological problems--tremors or shaking, seizures, new or worsening depression, confusion, or psychosis,

drowsiness:

- high blood pressure--severe headache, blurred vision, pounding in your neck or ears;
- kidney or heart problems--swelling, urinating less, rapid weight gain, feeling tired or short of breath;
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- **liver problems**--loss of appetite, nausea, vomiting, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- high blood potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement; or
- low red blood cells (anemia)--pale skin, tiredness, feeling light-headed or short of breath, cold hands and feet.

Indocin SR may also worsen asthma attacks in people who have asthma. Talk with your doctor about your own risk.

Older adults may be more likely to have serious stomach problems while taking medicine that contains an NSAID.

Common side effects of Indocin SR may include:

- · headache; dizziness;
- · heartburn, stomach pain, gas; or
- · nausea, vomiting.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Indocin SR side effects (more detail)

Related/similar drugs

Rinvog

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

Reviews & ratings

7.0 / 10

223 Reviews

View more

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-thecounter painkiller and fever ...

Reviews & ratings

6.1 / 10

13 Reviews

View more

FEATURED

Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used alone or with other ...

Reviews & ratings

5.7 / 10

145 Reviews

View more

Diclofenac

Diclofenac is a nonsteroidal antiinflammatory drug used to treat pain and inflammation associated ...

Reviews & ratings

7.3 / 10

695 Reviews

View more

Oxycodone

Oxycodone is an opioid analgesic used to treat moderate to severe pain; it has a high potential for ...

Reviews & ratings

6.9 / 10

1,142 Reviews

View more

Naproxen

Naproxen is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

683 Reviews

View more

Tramadol

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

Reviews & ratings

7.0 / 10

1,876 Reviews

View more

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

1,404 Reviews

View more

Duloxetine

Duloxetine is a selective serotonin and norepinephrine reuptake inhibitor antidepressant used to ...

Reviews & ratings

6.2 / 10

2,456 Reviews

View more

Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

Reviews & ratings

6.0 / 10

618 Reviews

View more

Warnings

Indocin SR can increase your risk of fatal heart attack or stroke. Do not use Indocin SR just before or after heart bypass surgery (coronary artery bypass graft, or CABG). This medicine may also cause stomach or intestinal bleeding, which can be fatal.

Before taking this medicine

Indocin SR can increase your risk of fatal heart attack or stroke. Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Indocin SR may also cause stomach or intestinal bleeding, which can be fatal. This can occur without warning, especially in older adults.

You should not use Indocin SR if you are allergic to it, or if you have ever had an asthma attack, hives or a severe allergic reaction after taking aspirin or an NSAID.

Tell your doctor if you have or have ever had:

- congestive heart failure, heart disease, high blood pressure, high cholesterol, diabetes, smoking;
- a heart attack, stroke, or blood clot;
- · bleeding problems;
- · stomach ulcers or bleeding;
- · depression, mood changes;
- seizures, epilepsy, Parkinson's disease;
- asthma;
- · fluid retention; or
- liver or kidney disease.

It may be harder for you to get pregnant while you are using this medicine.

Do not use Indocin SR during pregnancy unless your doctor tells you to. **Using an NSAID in the last 20 weeks can harm the unborn baby or cause pregnancy complications.**

Ask a doctor if it is safe to breastfeed while using this medicine.

How should I take Indocin SR?

Follow all directions on your prescription label and read all medication guides. **Use the lowest dose that is effective in treating your condition.**

Your dose needs may change if you switch to a different brand, strength, or form of Indocin SR. **Avoid medication** errors by using only the form and strength your doctor prescribes.

Take this medicine with food or milk if it upsets your stomach.

You may be given other medications to help prevent certain side effects. Keep taking these medicines for as long as your

doctor has prescribed.

Swallow the extended-release capsule whole and do not crush, chew, break, or open it.

Shake the oral suspension (liquid). Measure a dose with the supplied measuring device (not a kitchen spoon).

You may need medical tests to check your heart and kidney function. Your blood pressure will need to be checked often.

Tell your doctor if you have a planned surgery.

Call your doctor if your symptoms do not improve, or if they get worse. Indocin SR is not meant to be used long-term.

This medicine may affect the results of certain medical tests. Tell any doctor who treats you that you are using Indocin SR.

Store at room temperature away from moisture and heat. Store the liquid medicine tightly closed away from light. Do not allow the liquid medicine to freeze.

Detailed Indocin SR dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Indocin SR?

Avoid taking aspirin or other NSAIDs unless your doctor tells you to.

Avoid smoking and drinking alcohol. It may increase your risk of stomach bleeding.

Ask a doctor or pharmacist before using other medicines for pain, fever, swelling, or cold/flu symptoms. They may contain ingredients similar to Indocin SR (such as aspirin, ibuprofen, ketoprofen, or naproxen).

Avoid driving or hazardous activity until you know how Indocin SR will affect you. Your reactions could be impaired.

What other drugs will affect Indocin SR?

Ask your doctor before using Indocin SR if you take an antidepressant. Taking certain antidepressants with an NSAID may cause you to bruise or bleed easily.

Indocin SR can harm your kidneys, especially if you also use certain medicines for infections, cancer, osteoporosis, organ transplant rejection, high blood pressure, pain, or arthritis (including Advil, Motrin, and Aleve).

Tell your doctor about all your other medicines, especially:

· aspirin;

- · probenecid;
- a blood thinner--warfarin, Coumadin, Jantoven;
- medicine used to prevent blood clots--alteplase, clopidogrel, dipyridamole, ticlopidine, and others; or
- steroid medicine (such as prednisone).

This list is not complete. Other drugs may affect Indocin SR, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

i Indocin SR drug interactions (more detail)

Does Indocin SR interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

What is Ardosons called in the U.S?

More FAQ

- Why is my poop green? What does this mean?
- Can I take Tylenol while taking indomethacin and can these drugs be alternated every 3 hours?

More about Indocin SR (indomethacin)

- Check interactions
- · Compare alternatives
- Reviews (6)
- · Drug images
- Latest FDA alerts (6)
- Side effects
- Dosage information
- During pregnancy
- Drug class: Nonsteroidal anti-inflammatory drugs
- Breastfeeding

Patient resources

· Indocin SR advanced reading

Other brands

Tivorbex

Professional resources

- Indomethacin monograph
- Indomethacin Suppository (FDA)

Other brands

Tivorbex

Other formulations

• Indocin

Related treatment guides

- Back Pain
- Bursitis
- · Ankylosing Spondylitis
- Bartter Syndrome

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 19.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

প্ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

6 Reviews

Images

Indocin SR 75 MG (INDOCIN SR MSD 693)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





