

Home

2. Iloperidone

lloperidone

Generic name: iloperidone [EYE-loe-PER-i-done]

Brand name: Fanapt

Dosage form: oral tablet (1 mg; 1 mg-2 mg-4 mg-6 mg; 10 mg; 12 mg; 2 mg; 4 mg; 6 mg; 8 mg)

Drug class: Atypical antipsychotics



Medically reviewed by Drugs.com on Jan 27, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is iloperidone?

Iloperidone is an antipsychotic medication that is used to treat schizophrenia in adults.

lloperidone may also be used for purposes not listed in this medication guide.

lloperidone side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

High doses or long-term use of iloperidone can cause a serious movement disorder that may not be reversible.

The longer you use iloperidone, the more likely you are to develop this disorder, especially if you are an older adult.

lloperidone may cause serious side effects. Call your doctor at once if you have:

- · fast or pounding heartbeats, fluttering in your chest;
- a light-headed feeling, like you might pass out;
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement);
- tremors, slow muscle movement, muscles pain or stiffness;
- · confusion, agitation, thoughts of hurting yourself;
- loss of bladder control;
- · penis erection that is painful or lasts 4 hours or longer;
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor;
- low white blood cell counts--fever, chills, sore throat, mouth sores, skin sores, cough, trouble breathing; or
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven

heartbeats, tremors, feeling like you might pass out;

Older adults may be more likely to have side effects from iloperidone.

Common side effects of iloperidone may include:

- · weight gain;
- dizziness, drowsiness, tired feeling;
- dry mouth, stuffy nose; or
- fast heart rate.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i <u>lloperidone side effects</u> (more detail)

Related/similar drugs

Lybalvi

Lybalvi is used to treat adults with manic or mixed episodes associated with bipolar I disorder ...

Reviews & ratings

8.1 / 10

16 Reviews

View more

Abilify

Abilify is an antipsychotic medicine used to treat the symptoms of schizophrenia and bipolar ...

Reviews & ratings

6.1 / 10

1,187 Reviews

View more

FEATURED

Caplyta

Caplyta (lumateperone) may be used to treat schizophrenia or depression associated with bipolar ...

Reviews & ratings

6.6 / 10

View more

Seroquel

Seroquel is used to treat the symptoms of schizophrenia, bipolar disorder and major depressive ...

Reviews & ratings

7.0 / 10

694 Reviews

View more

Lamictal

Lamictal is an anti-epileptic medication used treat seizures in adults and children over 2 years ...

Reviews & ratings

8.0 / 10

792 Reviews

View more

Depakote

Depakote is used to treat various types of seizure disorders. Learn about side effects ...

Reviews & ratings

6.3 / 10

329 Reviews

View more

Lithium

Lithium is used to treat the manic episodes of manic depression - hyperactivity, rushed speech ...

Reviews & ratings

7.3 / 10

491 Reviews

Carbamazepine

Carbamazepine is used to treat epileptic seizures and nerve pain such as trigeminal neuralgia ...

Reviews & ratings

7.0 / 10

351 Reviews

View more

Risperidone

Risperidone is used to treat schizophrenia and symptoms of bipolar disorder. Learn about side ...

Reviews & ratings

5.6 / 10

731 Reviews

View more

Olanzapine

Olanzapine (brand name Zyprexa) is an atypical antipsychotic that may be used to treat adults and ...

Reviews & ratings

5.9 / 10

1,091 Reviews

View more

Warnings

Iloperidone is not approved for use in older adults with dementia-related psychosis.

Before taking this medicine

You should not use iloperidone if you are allergic to it.

Iloperidone may increase the risk of death in older adults with dementia-related psychosis and is not approved for this use.

Tell your doctor if you have ever had:

· trouble swallowing;

• long QT syndrome (in you or a family member);

a seizure;

diabetes, or risk factors such as being overweight or having a family history of diabetes;

breast cancer;

· high cholesterol or triglycerides;

an electrolyte imbalance (such as low levels of potassium or magnesium in your blood);

· low bone mineral density; or

· liver disease.

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of iloperidone on the baby.

Taking antipsychotic medicine in the last 3 months of pregnancy may cause breathing problems, feeding problems, or withdrawal symptoms in the newborn. If you get pregnant, tell your doctor right away. Do not stop taking iloperidone without your doctor's advice.

You should not breastfeed while you are taking iloperidone.

Iloperidone is not approved for use by anyone younger than 18 years old.

How should I take iloperidone?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

It may take up to 2 weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Your doctor will need to check your progress on a regular basis.

If you have stopped taking iloperidone for more than 3 days in a row, talk with your doctor before you start taking it again. You may need to restart the medicine at a lower dose.

Store at room temperature away from moisture, heat, and light.

lloperidone dosing information

Usual Adult Dose for Schizophrenia:

Initial dose: 1 mg orally twice a day

Maintenance dose: 6 to 12 mg twice a day

Maximum dose: 24 mg/day

Comments:

- -This drug must be titrated slowly to avoid orthostatic hypotension.
- -Control of symptoms may be delayed during the first 1 to 2 weeks of treatment because of the need to titrate slowly.
- -Treatment should be increased in increments of not more than 2 mg twice daily as tolerated.
- -Healthcare providers should periodically re-evaluate the long-term risks and benefits of the drug for the individual patient.

Use: Treatment of schizophrenia

1 Detailed Iloperidone dosage information

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking iloperidone?

Avoid driving or hazardous activity until you know how iloperidone will affect you. Your reactions could be impaired.

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Dizziness or severe drowsiness can cause falls, fractures, or other injuries.

While you are taking iloperidone, you may be more sensitive to very hot conditions. Avoid becoming overheated or dehydrated. Drink plenty of fluids, especially in hot weather and during exercise.

Drinking alcohol with this medicine can cause side effects.

What other drugs will affect iloperidone?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

lloperidone can cause a serious heart problem. Your risk may be higher if you also use certain other medicines for infections, asthma, heart problems, high blood pressure, depression, mental illness, cancer, malaria, or HIV.

Using iloperidone with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Many drugs can affect iloperidone. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all your current medicines and any medicine you start or stop using.

<u>Iloperidone drug interactions</u> (more detail)

Does iloperidone interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

iloperidone	
+	
Enter a drug name	Add

More about iloperidone

- Check interactions
- Compare alternatives
- Reviews (41)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: atypical antipsychotics
- Breastfeeding
- En español

Patient resources

• Iloperidone advanced reading

Other brands

Fanapt

Professional resources

- Iloperidone monograph
- Iloperidone (FDA)

Other brands

Fanapt

Related treatment guides

- Bipolar Disorder
- Psychosis
- Schizophrenia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.02.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

7.1 / 10

41 Reviews

Images

Iloperidone 6 mg (053)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×