

Home

Hycodan

Hycodan ⊲

Generic name: homatropine and hydrocodone [HOM-a-TROE-peen-and-HYE-dro-KOE-done]

Brand names: Hycodan, Hydromet, Tussigon **Drug class:** Upper respiratory combinations



Medically reviewed by Drugs.com on Aug 14, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Hycodan?

Hycodan is a combination medicine used to treat runny or stuffy nose, sneezing, cough, and sinus congestion caused by allergies or the common cold.

Hycodan contains an opioid cough medicine and may be habit-forming.

Hycodan may also be used for purposes not listed in this medication guide.

Hycodan side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up.

Stop taking Hycodan and call your doctor at once if you have:

- noisy breathing, sighing, shallow breathing;
- extreme drowsiness, confusion, feeling weak or limp;
- a seizure;
- a light-headed feeling, like you might pass out;
- severe constipation, stomach pain;
- little or no urination;
- adrenal gland problems--nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness; or
- high levels of serotonin in the body--agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle

stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea.

Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders.

Common side effects of Hycodan may include:

- · drowsiness, dizziness;
- lack of energy, coordination problems;
- headache, confusion;
- dry mouth, nausea, vomiting, constipation;
- tremors, fast or irregular heart rate; or
- feeling anxious, restless, nervous, or irritable.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Hycodan side effects (more detail)

Related/similar drugs

Benadryl

Benadryl is an antihistamine and is used to treat sneezing, runny nose, itching and other ...

Reviews & ratings

7.0 / 10

79 Reviews

View more

Mucinex

Mucinex (guaifenesin) is used to relieve the symptoms of cough and loosen mucus in the chest ...

Reviews & ratings

6.2 / 10

155 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to cavera ulcarativa colitic or Crohn's

disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Norco

Norco (acetaminophen and hydrocodone) is used to relieve moderate to severe pain. Includes Norco ...

Reviews & ratings

6.3 / 10

367 Reviews

View more

Dilaudid

Dilaudid (hydromorphone) is a narcotic pain reliever used to treat moderate to severe pain ...

Reviews & ratings

8.2 / 10

329 Reviews

View more

Vicodin

Vicodin (acetaminophen and hydrocodone) is used to relieve moderate to severe pain. Includes ...

Reviews & ratings

7.7 / 10

146 Reviews

View more

Codeine

Codeine is a opioid pain medicine used to treat mild to moderate pain. Includes codeine side ...

Reviews & ratings

View more

Dextromethorphan

Dextromethorphan is a cough suppressant used to treat coughing. Learn about side effects ...

Reviews & ratings

3.7 / 10

386 Reviews

View more

Guaifenesin

Guaifenesin is an expectorant. It helps loosen congestion in your chest and throat. Includes ...

Reviews & ratings

7.1 / 10

305 Reviews

View more

Warnings

MISUSE OF THIS MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medication in a place where others cannot get to it.

Fatal side effects can occur if you use this medicine with alcohol, or with other drugs that cause drowsiness or slow your breathing.

Before taking this medicine

You should not use Hycodan if you are allergic to homatropine or hydrocodone, or if you have:

- severe asthma or breathing problems;
- a stomach or bowel obstruction (including paralytic ileus);
- severe or uncontrolled high blood pressure;
- severe coronary artery disease (clogged arteries);
- narrow-angle glaucoma; or

• if you are unable to urinate.

Do not give this medicine to anyone younger than 6 years old.

Do not use this medicine if you have used an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine.

Tell your doctor if you have ever had:

- lung disease or breathing problems;
- a head injury, brain tumor, or seizure;
- constipation, a bowel obstruction, or stomach problems;
- problems with your bile duct, pancreas, or adrenal gland;
- an enlarged prostate;
- · urination problems;
- · liver or kidney disease;
- low blood pressure;
- heart disease, a blood vessel disorder;
- a drug addiction; or
- if you have a fever and cough with mucus.

If you use opioid medicine while you are pregnant, your baby could become dependent on the drug. This can cause life-threatening withdrawal symptoms in the baby after it is born. Babies born dependent on opioids may need medical treatment for several weeks.

Ask a doctor before using opioid medicine if you are breastfeeding. Tell your doctor if you notice severe drowsiness or slow breathing in the nursing baby.

How should I take Hycodan?

Follow the directions on your prescription label and read all medication guides. **Never use this medicine in larger amounts, or for longer than prescribed.** Cold or cough medicine is only for short-term use until your symptoms clear up.

Never share this medicine with another person, especially someone with a history of drug abuse or addiction.

MISUSE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medicine in a place where others cannot get to it. Selling or giving away this medicine is against the law.

Measure **liquid medicine** carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon). Rinse after each use.

Call your doctor if your symptoms do not improve after 5 days, or if you have a fever, rash, or headaches.

If you need surgery or medical tests, tell the surgeon or doctor ahead of time if you have taken a cough or cold medicine within the past few days.

Store at room temperature away from moisture and heat. Do not freeze.

Do not keep leftover medication. Just one dose can cause death in someone using this medicine accidentally or improperly. Read and carefully follow the instructions provided with this medicine about how to safely dispose of any unused portion.

Detailed Hycodan dosage information

What happens if I miss a dose?

Since Hycodan is used when needed, you may not be on a dosing schedule. Skip any missed dose if it's almost time for your next dose. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. An opioid overdose can be fatal, especially in a child or other person using the medicine without a prescription. Overdose symptoms may include severe drowsiness, pinpoint pupils, slow breathing, or no breathing.

Your doctor may recommend you get naloxone (a medicine to reverse an opioid overdose) and keep it with you at all times. A person caring for you can give the naloxone if you stop breathing or don't wake up. Your caregiver must still get emergency medical help and may need to perform CPR (cardiopulmonary resuscitation) on you while waiting for help to arrive.

Anyone can buy naloxone from a pharmacy or local health department. Make sure any person caring for you knows where you keep naloxone and how to use it.

What should I avoid while taking Hycodan?

Do not drink alcohol. Dangerous side effects or death could occur.

Avoid driving or operating machinery until you know how Hycodan will affect you. Dizziness or severe drowsiness can cause falls or other accidents.

What other drugs will affect Hycodan?

You may have breathing problems or withdrawal symptoms if you start or stop taking certain other medicines. Tell your doctor if you also use an antibiotic, antifungal medication, heart or blood pressure medication, seizure medication, or medicine to treat HIV or hepatitis C.

Opioid medication can interact with many other drugs and cause dangerous side effects or death. Be sure your doctor knows if you also use:

- cold or allergy medicines, bronchodilator asthma/COPD medication, or a diuretic ("water pill");
- · medicines for motion sickness, irritable bowel syndrome, or overactive bladder;
- other opioids--opioid pain medicine or prescription cough medicine;
- a sedative like Valium--diazepam, alprazolam, lorazepam, Xanax, Klonopin, Versed, and others;

- drugs that make you sleepy or slow your breathing--a sleeping pill, muscle relaxer, medicine to treat mood disorders or mental illness;
- drugs that affect serotonin levels in your body--a stimulant, or medicine for depression, Parkinson's disease, migraine headaches, serious infections, or nausea and vomiting.

This list is not complete. Other drugs may affect Hycodan, including prescription and over-the-counter medicines, vitamins, and herbal products. **Not all possible interactions are listed here.**

<u>Hycodan drug interactions</u> (more detail)

Does Hycodan interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Hycodan	
+	
Enter a drug name	Add

More about Hycodan (homatropine / hydrocodone)

- · Check interactions
- · Compare alternatives
- · Pricing & coupons
- Reviews (37)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Drug class: upper respiratory combinations
- En español

Patient resources

· Hycodan advanced reading

Other brands

Hydromet, Tussigon

Professional resources

• Hycodan prescribing information

• Hycodan Oral Solution (FDA)

Other brands

Hydromet, Tussigon

Related treatment guides

Cough

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ം Risk data available

CSA Schedule*

2 High potential for abuse

Approval History

10+ years FDA approved 1943

User Reviews & Ratings

8.6 / 10

37 Reviews

Images

Hycodan 1.5 mg-5 mg (HYCODAN)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











P

Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

X