

Home

Opcicon One-Step

Opcicon One-Step

Generic name: levonorgestrel emergency contraceptive [LEE-voe-nor-jes-trel]

Brand names: AfterPill, Curae, EContra EZ, EContra One-Step, Her Style, ... show all 15 brands

Drug class: Contraceptives



Medically reviewed by Drugs.com on Apr 4, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Opcicon One-Step?

Opcicon One-Step is used to prevent pregnancy after unprotected sex or failure of other forms of birth control (such as condom breakage, or missing 2 or more birth control pills).

Opcicon One-Step may also be used for purposes not listed in this medication guide.

Opcicon One-Step side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor or seek emergency medical help if you have severe pain in your lower stomach or side. This could be a sign of a tubal pregnancy (a pregnancy that implants in the fallopian tube instead of the uterus). A tubal pregnancy is a medical emergency.

Common side effects of Opcicon One-Step may include:

- · stomach pain;
- · breast pain or tenderness;
- nausea, vomiting, diarrhea;
- headache, dizziness;
- · feeling tired; or
- · changes in your menstrual periods.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Opcicon One-Step side effects (more detail)

Related/similar drugs

Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

Reviews & ratings

6.1 / 10

2,624 Reviews

View more

Aviane

Aviane is used as contraception to prevent pregnancy. Includes side effects, interactions and ...

Reviews & ratings

6.1 / 10

554 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Kyleena

The Kyleena IUD is placed in your uterus by your healthcare provider to prevent pregnancy for up to ...

Reviews & ratings

6.6 / 10

1,637 Reviews

View more

ella

ella (ulipristal) is an emergency contraceptive used to prevent pregnancy after having sex without ...

Reviews & ratings

7.1 / 10

771 Reviews

View more

Liletta

Liletta (levonorgestrel) is an intrauterine device (IUD) used to prevent pregnancy and also to ...

Reviews & ratings

5.5 / 10

647 Reviews

View more

Skyla

Skyla (levonorgestrel intrauterine system) is flexible plastic T-shaped device used for prevention ...

Reviews & ratings

6.9 / 10

1,537 Reviews

View more

Warnings

Opcicon One-Step will not terminate pregnancy if the fertilized egg has already attached to the uterus.

Ask a doctor or pharmacist before taking levonorgestrel if you regularly use medication for seizures, tuberculosis, or HIV/AIDS. Certain other medicines can make levonorgestrel less effective.

Before taking this medicine

Opcicon One-Step is not intended for use as a routine form of birth control. Talk with your doctor about the many forms of birth control available.

Do not use this medicine if you are already pregnant. Opcicon One-Step will not terminate a pregnancy that has

already begun (the fertilized egg has attached to the uterus).

Not approved for use by anyone younger than 17 years old.

You should not use levonorgestrel if you are allergic to it.

Ask a doctor or pharmacist before taking levonorgestrel if you regularly use medication for seizures, tuberculosis, or HIV/AIDS. Certain medications can make levonorgestrel less effective as an emergency form of contraception.

Levonorgestrel may slow breast milk production. Tell your doctor if you are breastfeeding.

How should I take Opcicon One-Step?

Use exactly as directed on the label, or as prescribed by your doctor.

Opcicon One-Step must be taken as soon as possible after unprotected sex (no later than 72 hours afterward).

Call your doctor right away if you vomit within 2 hours after taking Opcicon One-Step. **Do not take a second dose without first asking your doctor.**

If your period is late by 1 week or longer after the expected date, you may be pregnant. Get a pregnancy test and contact your doctor if you are pregnant. Opcion One-Step will not terminate pregnancy if the fertilized egg has attached to the uterus.

Within 3 weeks after taking Opcicon One-Step, a doctor should confirm that you are not pregnant, and that this medicine has not caused any harmful effects.

Store at room temperature away from moisture and heat.

Detailed Opcicon One-Step dosage information

What happens if I miss a dose?

Since Opcicon One-Step is used as a single dose, it does not have a daily dosing schedule.

What happens if I overdose?

Because Opcicon One-Step is supplied as a single tablet in an exact strength, an overdose is unlikely to occur when the levonorgestrel is used as directed. **Do not take more than one tablet at the same time.**

What should I avoid while taking Opcicon One-Step?

Opcicon One-Step will not protect you from sexually transmitted diseases--including HIV and AIDS. Using a condom is the only way to protect yourself from these diseases. Avoid having unprotected sex.

What other drugs will affect Opcicon One-Step?

Certain other medications can make Opcicon One-Step less effective, which may result in pregnancy. Ask a

doctor or pharmacist if Opcicon One-Step is safe to use if you are using any of the following medications:

- · efavirenz;
- rifampin; or
- seizure medication--carbamazepine, felbamate, fosphenytoin, phenobarbital, phenytoin, primidone.

This list is not complete. Other drugs may affect levonorgestrel, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

① Opcicon One-Step drug interactions (more detail)

Does Opcicon One-Step interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

Can Plan B make your period late or cause bleeding?	~
Can you drink alcohol after taking Plan B?	~
How effective is Plan B and how late can you take it?	~
How many times can you take Plan B?	~
What's the weight limit for Plan B?	~
What are my birth control options and how effective are they?	~

More FAQ

• Mirena, Kyleena, Skyla & Liletta - What's the difference?

More about Opcicon One-Step (levonorgestrel)

- · Check interactions
- Compare alternatives
- Pricing & coupons

- Reviews (4)
- Drug images
- Latest FDA alerts (1)
- Side effects
- Dosage information
- During pregnancy
- Drug class: contraceptives
- En español

Patient resources

Other brands

Mirena, Kyleena, Liletta, Skyla, ... +15 more

Professional resources

Levonorgestrel Emergency Contraceptive prescribing information

Other brands

Mirena, Kyleena, Liletta, Skyla, ... +3 more

Related treatment guides

• Emergency Contraception

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.02.

Emergency Contraception 101

Emergency Contraception Overview

Emergency Contraceptives Available in the U.S.

DRUG STATUS

Availability OTC Over the counter Pregnancy & Lactation Risk data available CSA Schedule* N/A Not a controlled drug Approval History

Drug history at FDA

User Reviews & Ratings

4 Reviews

Related News

Certain Birth Control Pills, Devices Linked to Higher Heart Risks

Images

Opcicon One-Step levonorgestrel 1.5mg (718)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

 \times