

#### Home

2. Loteprednol/tobramycin

# Loteprednol and tobramycin (ophthalmic)

Generic name: loteprednol and tobramycin (ophthalmic) [ LOE-te-PRED-nol-and-TOE-bra-MYE-sin-off-THAL-mik ]

Brand name: Zylet

**Dosage form:** ophthalmic suspension (0.5%-0.3%) **Drug class:** Ophthalmic steroids with anti-infectives



Medically reviewed by Drugs.com on Nov 18, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

## What is loteprednol and tobramycin ophthalmic?

Loteprednol and tobramycin ophthalmic (for the eyes) is a combination antibiotic and steroid medicine that is used to treat eye inflammation caused by allergies, shingles (herpes zoster), severe acne, iritis, uveitis, eye injury, radiation, chemical burns, or certain other conditions.

loteprednol and tobramycin is used when there is a risk of bacterial infection in or around the eye.

Loteprednol and tobramycin ophthalmic may also be used for purposes not listed in this medication guide.

## Loteprednol and tobramycin ophthalmic side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Loteprednol and tobramycin may cause serious side effects. Call your doctor at once if you have:

- severe pain, burning or stinging when using the eye drops;
- blurred vision, tunnel vision, seeing halos around lights;
- pain behind your eyes, sudden vision changes;
- slow healing after eye surgery;
- signs of eye infection--redness, severe discomfort, crusting or drainage.

#### Common side effects of loteprednol and tobramycin may include:

- · blurred vision; or
- minor burning or stinging.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.

1 Loteprednol / tobramycin ophthalmic side effects (more detail)

# Related/similar drugs

### Humira

Humira is a tumor necrosis factor blocker that is used to treat many inflammatory conditions such ...

## **Reviews & ratings**

6.4 / 10

698 Reviews

View more

#### FEATURED

## **Ozempic**

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

## **Reviews & ratings**

6.6 / 10

1,504 Reviews

View more

## **Triamcinolone**

Triamcinolone is used to treat allergies, skin conditions, ulcerative colitis, and arthritis. Learn ...

## **Reviews & ratings**

6.7 / 10

226 Reviews

View more

## **Warnings**

You should not use loteprednol and tobramycin if you have a fungal or viral infection in your eyes (including herpes simplex).

## Before taking this medicine

You should not use this medicine if you are allergic to loteprednol or tobramycin, or if you have a fungal or viral infection in your eyes (including herpes simplex).

Tell your doctor if you have ever had:

- glaucoma; or
- cataracts, or if you need cataract surgery.

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant.

It may not be safe to breast-feed while using this medicine. Ask your doctor about any risk.

## How should I use loteprednol and tobramycin ophthalmic?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

**Do not use while wearing soft contact lenses.** A preservative in loteprednol and tobramycin could permanently stain the lenses. Use the medicine at least 15 minutes before inserting your contact lenses.

Wash your hands before using eye medication.

Shake the eye drops well just before each use.

**To use this medicine:** Tilt your head back slightly and pull down your lower eyelid to create a small pocket. Hold the dropper above the eye and squeeze a drop into this pocket. Close your eyes for 1 or 2 minutes.

Use only the number of drops your doctor has prescribed.

Do not touch the tip of the eye dropper or place it directly on your eye. A contaminated dropper can infect your eye, which could lead to serious vision problems.

If you use this medicine for longer than 10 days, you may need frequent vision tests to check the pressure inside your eyes.

Call your doctor if your symptoms do not improve after 2 days of treatment.

You should not stop using this medicine suddenly. Follow your doctor's instructions about tapering your dose.

Store this medicine in an upright position at room temperature. Do not freeze.

1 Detailed Loteprednol / tobramycin ophthalmic dosage information

## What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

## What happens if I overdose?

An overdose of loteprednol and tobramycin ophthalmic is not expected to be dangerous. Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

## What should I avoid while using loteprednol and tobramycin ophthalmic?

Do not share loteprednol and tobramycin with another person, even if they have the same symptoms you have.

## What other drugs will affect loteprednol and tobramycin ophthalmic?

Medicine used in the eyes is not likely to be affected by other drugs you use. But many drugs can interact with each other. Tell each of your healthcare providers about all medicines you use, including prescription and over-the-counter medicines, vitamins, and herbal products.

1 Loteprednol / tobramycin ophthalmic drug interactions (more detail)

## Does loteprednol/tobramycin ophthalmic interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

loteprednol/tobramycin ophthalmic
+
Enter a drug name
Add

## More about loteprednol / tobramycin ophthalmic

- Check interactions
- · Compare alternatives
- Reviews (5)
- Side effects
- Dosage information
- During pregnancy
- · Drug class: ophthalmic steroids with anti-infectives
- En español

#### Patient resources

· Loteprednol and tobramycin advanced reading

## Other brands

**Zylet** 

## **Professional resources**

#### Other brands

**Zylet** 

## Related treatment guides

- · Conjunctivitis, Bacterial
- Cyclitis
- Iritis
- Keratitis
- Uveitis

## **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

#### **Antibiotics 101**

List of Common Antibiotics & Types

**Antibiotics and Alcohol** 

**Probiotics and Antibiotics** 

**Antibiotics for Strep Throat** 

Antibiotics for a Tooth Infection

Antibiotics for UTI's

## **DRUG STATUS**

# Availability Rx Prescription only Pregnancy & Lactation Risk data available CSA Schedule\* N/A Not a controlled drug

**Approval History**Drug history at FDA

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com Advertising policy

Attribution & citations

#### Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

#### Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







