

#### Home

Apriso

# Apriso 🖘

Generic name: mesalamine (oral) [ me-SAL-a-meen ]

Drug class: 5-aminosalicylates

Medically reviewed by Sanjai Sinha, MD. Last updated on Apr 8, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

# What is Apriso?

Apriso affects a substance in the body that causes inflammation, tissue damage, and diarrhea.

Apriso is used to treat mild to moderate ulcerative colitis in patients 18 years of age and older.

Apriso is used to prevent the symptoms of ulcerative colitis from recurring.

# Warnings

Stop using Apriso and call your doctor at once if you have severe stomach pain, stomach cramping, bloody diarrhea (may occur with fever, headache, and skin rash).

# Before taking this medicine

You should not use Apriso if you are allergic to mesalamine, aspirin, sulfasalazine, or salicylates (such as Disalcid, Doans Pills, Dolobid, Kaopectate, Nuprin Backache, Pepto-Bismol, Tricosal, Trilisate, and others).

To make sure Apriso is safe for you, tell your doctor if you have ever had:

- a kidney stone or kidney disease;
- liver disease;
- a blockage in your stomach or intestines (such as pyloric stenosis); or
- a skin condition such as eczema.

Apriso capsules contain phenylalanine. Tell your doctor if you have phenylketonuria (PKU).

Tell your doctor if you are pregnant or plan to become pregnant. It is not known whether mesalamine will harm an unborn baby. However, having untreated or uncontrolled ulcerative colitis during pregnancy may cause complications such as low birth weight or premature birth. The benefit of treating ulcerative colitis may outweigh any risks to the baby.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk. If you are breastfeeding, tell your doctor if you notice diarrhea in the nursing baby.

Do not give this medicine to a child. Apriso is not approved for use in anyone younger than 18 years old.

Apriso pregnancy and breastfeeding warnings (more detail)

# How should I take Apriso?

Take Apriso exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using mesalamine.

You may take Apriso with or without food.

Drink plenty of liquids to help prevent kidney stones.

Swallow the capsule whole and do not open, chew, or break it. **Tell your doctor if you have trouble swallowing the capsule**.

Call your doctor if your ulcerative colitis symptoms do not improve, or if they get worse.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Apriso.

You may need frequent blood tests to check your blood cells or kidney function.

Store at room temperature away from moisture and heat.

# **Dosing information**

Usual Adult Dose for Ulcerative Colitis -- Maintenance:

Apriso 375 mg extended release capsules: 1.5 g (4 capsules) orally once a day in the morning

Comments:

Do not co-administer Apriso 375 mg extended release capsules with antacids.

Use: Maintenance of remission of ulcerative colitis in adults.

Detailed Apriso dosage information

# What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

# What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

# What should I avoid while taking Apriso?

If you have a skin condition such as eczema, mesalamine could make you sunburn more easily. Avoid sunlight or tanning beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Ask your doctor before using an antacid, and use only the type your doctor recommends. Some antacids can make it harder for your body to absorb mesalamine.

# Apriso side effects

Get emergency medical help if you have **signs of an allergic reaction to Apriso:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Stop using Apriso and call your doctor at once if you have:

- · severe stomach pain, stomach cramping, bloody diarrhea;
- fever, headache, general ill feeling;
- rash, itching, eye redness;
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- low white blood cell counts fever, chills, mouth sores, skin sores, sore throat, cough, feeling light-headed, trouble breathing;
- signs of a kidney stone severe pain in your side and back, frequent need to urinate, foul-smelling urine, dark or cloudy urine;
- kidney problems increased or decreased urination, swelling, weight gain; or
- **liver problems** loss of appetite, upper stomach pain, tiredness, easy bruising or bleeding, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Low white blood cell counts may be more likely in older adults.

Common Apriso side effects may include:

- burping, constipation, nausea, vomiting, stomach pain, diarrhea, gas;
- dizziness;
- cold symptoms such as stuffy nose, sneezing, sore throat;
- back pain;
- · headache;
- · rash; or
- · abnormal liver function tests.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

# Related/similar drugs

# **Skyrizi**

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

# **Reviews & ratings**

7.9 / 10

56 Reviews

View more

### **Omvoh**

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

# Reviews & ratings

1 Reviews

View more

### FEATURED

## **Entyvio**

Entyvio (vedolizumab) is used to treat ulcerative colitis (UC) and Crohn's disease, reducing ...

# **Reviews & ratings**

6.2 / 10

162 Reviews

View more

# Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used alone or with other ...

### **Reviews & ratings**

5.7 / 10

145 Reviews

View more

### **Stelara**

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

### **Reviews & ratings**

7.6 / 10

252 Reviews

View more

### Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

# Reviews & ratings

7.0 / 10

223 Reviews

View more

### Humira

Humira is a tumor necrosis factor blocker that is used to treat many inflammatory conditions such ...

### **Reviews & ratings**

6.4 / 10

698 Reviews

View more

# **Tremfya**

Tremfya (guselkumab) injection is used to treat plaque psoriasis, psoriatic arthritis, ulcerative ...

### **Reviews & ratings**

7.2 / 10

99 Reviews

View more

# **Sulfasalazine**

Sultasalazine is used to treat ulcerative colitis, rheumatoid arthritis, and polyarticular-course ...

### **Reviews & ratings**

6.0 / 10

198 Reviews

View more

# What other drugs will affect Apriso?

Apriso can harm your kidneys, especially if you also use certain medicines for infections, cancer, osteoporosis, organ transplant rejection, bowel disorders, high blood pressure, or pain or arthritis (including Advil, Motrin, and Aleve).

Tell your doctor about all your other medicines, especially:

• azathioprine or 6-mercaptopurine.

This list is not complete. Other drugs may interact with mesalamine, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Apriso drug interactions (more detail)

# Does Apriso interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Apriso
+
Enter a drug name
Add

# **Popular FAQ**

Is mesalamine an immunosuppressant?

What are 7 mesalamine side effects to be aware of?	~
How long does it take for mesalamine to work?	~
When is the best time to take mesalamine?	~
Why is mesalamine so expensive?	~
How long can you take mesalamine?	~

#### References

1. Apriso Product Label

# More about Apriso (mesalamine)

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (29)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- · FDA approval history
- Drug class: 5-aminosalicylates
- Breastfeeding
- En español

### **Patient resources**

### Other brands

Lialda, Pentasa, Canasa, Rowasa, ... +2 more

# **Professional resources**

- Apriso prescribing information
- Mesalamine (AHFS Monograph)

## Other brands

Lialda, Pentasa, Asacol, Canasa, ... +4 more

# Related treatment guides

- Ulcerative Colitis, Maintenance
- Ulcerative Colitis

- Inflammatory Bowel Disease
- Ulcerative Proctitis

# **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Apriso only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 16.01.

#### **DRUG STATUS**

### **Availability**

Rx Prescription only

### **Pregnancy & Lactation**

প্ Risk data available

### **CSA Schedule\***

N/A Not a controlled drug

### **Approval History**

Drug history at FDA

### **User Reviews & Ratings**

6.7 / 10

29 Reviews

### **Images**

Apriso 0.375 g (G M)

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com

Advertising policy

Attribution & citations

### Terms & privacy

Terms of use

Editorial policy

Privacy policy

### Support

Help center

**Sitemap** 

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

