

Home

2. Asacol HD

Asacol HD

Generic name: mesalamine (oral) [me-SAL-a-meen]

Drug class: 5-aminosalicylates

Medically reviewed by Sanjai Sinha, MD. Last updated on Apr 8, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is Asacol HD?

Asacol HD affects a substance in the body that causes inflammation, tissue damage, and diarrhea.

Asacol HD delayed-release 800mg tablets are used to treat moderately active ulcerative colitis in adults.

Warnings

Stop using Asacol HD and call your doctor at once if you have severe stomach pain, stomach cramping, bloody diarrhea (may occur with fever, headache, and skin rash).

Before taking this medicine

You should not use Asacol HD if you are allergic to mesalamine, aspirin, sulfasalazine, or salicylates (such as Disalcid, Doans Pills, Dolobid, Kaopectate, Nuprin Backache, Pepto-Bismol, Tricosal, Trilisate, and others).

To make sure Asacol HD is safe for you, tell your doctor if you have ever had:

- · a kidney stone or kidney disease;
- · liver disease;
- a blockage in your stomach or intestines (such as pyloric stenosis); or
- a skin condition such as eczema.

Tell your doctor if you are pregnant or plan to become pregnant. It is not known whether mesalamine will harm an unborn baby. However, having untreated or uncontrolled ulcerative colitis during pregnancy may cause complications such as low birth weight or premature birth. The benefit of treating ulcerative colitis may outweigh any risks to the baby.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk. If you are breastfeeding, tell your doctor if you notice diarrhea in the nursing baby.

Do not give this medicine to a child. Asacol HD is not approved for use in anyone younger than 18 years old.

How should I take Asacol HD?

Take Asacol HD exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Asacol HD.

Take Asacol HD on an empty stomach, at least 1 hour before or 2 hours after a meal.

Drink plenty of liquids to help prevent kidney stones.

Swallow the tablet whole and do not crush, chew, or break it. **Tell your doctor if you have trouble swallowing the tablet.**

Tell your doctor if you find undissolved tablets in your stool.

Call your doctor if your ulcerative colitis symptoms do not improve, or if they get worse.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using this medicine.

You may need frequent blood tests to check your blood cells or kidney function.

Store at room temperature away from moisture and heat.

Dosing information

Usual Adult Dose for Ulcerative Colitis:

800 mg delayed release tablets: 1600 mg orally 3 times a day

Duration of therapy: 6 weeks.

Detailed Asacol HD dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Asacol HD?

If you have a skin condition such as eczema, Asacol HD could make you sunburn more easily. Avoid sunlight or tanning



beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Asacol HD side effects

Get emergency medical help if you have **signs of an allergic reaction to Asacol HD:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Stop using Asacol HD and call your doctor at once if you have:

- severe stomach pain, stomach cramping, bloody diarrhea;
- fever, headache, general ill feeling;
- rash, itching, eye redness;
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- low white blood cell counts fever, chills, mouth sores, skin sores, sore throat, cough, feeling light-headed, trouble breathing;
- signs of a kidney stone severe pain in your side and back, frequent need to urinate, foul-smelling urine, dark or cloudy urine;
- kidney problems increased or decreased urination, swelling, weight gain; or
- liver problems loss of appetite, upper stomach pain, tiredness, easy bruising or bleeding, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Low white blood cell counts may be more likely in older adults.

Common Asacol HD side effects may include:

- burping, constipation, nausea, vomiting, stomach pain, diarrhea, gas;
- · dizziness:
- · cold symptoms such as stuffy nose, sneezing, sore throat;
- back pain;
- · headache:
- · rash: or
- abnormal liver function tests.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Asacol HD side effects (more detail)

Related/similar drugs

Skyrizi

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis,

ulcerative colitis ...

Reviews & ratings

7.9 / 10

56 Reviews

View more

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

FEATURED

Entyvio

Entyvio (vedolizumab) is used to treat ulcerative colitis (UC) and Crohn's disease, reducing ...

Reviews & ratings

6.2 / 10

162 Reviews

View more

Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used alone or with other ...

Reviews & ratings

5.7 / 10

145 Reviews

View more

Stelara

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

Reviews & ratings

7.6 / 10

View more

Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

Reviews & ratings

7.0 / 10

223 Reviews

View more

Hydrocortisone topical

Hydrocortisone cream is used to treat eczema, dermatitis, psoriasis, hemorrhoids, insect bites or ...

Reviews & ratings

7.0 / 10

63 Reviews

View more

Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

Reviews & ratings

7.1 / 10

484 Reviews

View more

Prednisone

Prednisone is used to treat allergic disorders, ulcerative colitis, psoriasis and arthritis. Learn ...

Reviews & ratings

7.5 / 10

867 Reviews

View more

What other drugs will affect Asacol HD?

Mesalamine can harm your kidneys, especially if you also use certain medicines for infections, cancer, osteoporosis, organ transplant rejection, bowel disorders, high blood pressure, or pain or arthritis (including Advil, Motrin, and Aleve).

Tell your doctor about all your other medicines, especially:

• azathioprine or 6-mercaptopurine.

This list is not complete. Other drugs may interact with mesalamine, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Asacol HD drug interactions (more detail)

Does Asacol HD interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Asacol HD	
+	
Enter a drug name	Add

Popular FAQ

How long does it take for mesalamine to work?	~
When is the best time to take mesalamine?	~
Why is mesalamine so expensive?	~
How long can you take mesalamine?	~
Is mesalamine an immunosuppressant?	~
Can I stop taking mesalamine?	~
What is the maximum dosage of mesalamine?	~
What are 7 mesalamine side effects to be aware of?	~

More about Asacol HD (mesalamine)

· Check interactions

- · Compare alternatives
- Reviews (9)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: 5-aminosalicylates
- Breastfeeding

Patient resources

• Asacol HD advanced reading

Other brands

Lialda, Pentasa, Apriso, Canasa, ... +2 more

Professional resources

- · Asacol HD prescribing information
- Mesalamine (AHFS Monograph)

Other brands

Lialda, Pentasa, Apriso, Canasa, ... +3 more

Other formulations

Asacol

Related treatment guides

- · Crohn's Disease, Maintenance
- Ulcerative Colitis
- Ulcerative Colitis, Maintenance
- Inflammatory Bowel Disease
- Ulcerative Colitis, Active
- Ulcerative Proctitis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Asacol HD only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal

circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 16.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

8.9 / 10

9 Reviews

Images

Asacol HD 800 mg (PG 800)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center

Sitemap

Contact us













<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







×

Copyright © 2000-2025 Drugs.com. All rights reserved.