



[Home](#)

2. Lipitor

Lipitor

Generic name: [atorvastatin](#) [a-TOR-vä-stä-tin]

Dosage form: oral tablet (10 mg; 20 mg; 40 mg; 80 mg)

Drug class: [Statins](#)

Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on Oct 12, 2023.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is Lipitor?

Lipitor is used together with diet, weight loss, and exercise to reduce the risk of heart attack and stroke and to decrease the chance that heart surgery will be needed in people who have heart disease or who are at risk of developing heart disease.

Lipitor is also used to lower the amount of fatty substances such as low-density lipoprotein (LDL) cholesterol ('bad cholesterol') and triglycerides in the blood and to increase the amount of high-density lipoprotein (HDL) cholesterol ('good cholesterol') in the blood.

Lipitor may also be used to decrease the amount of cholesterol and other fatty substances in the blood in children and teenagers 10 to 17 years of age who have familial heterozygous hypercholesterolemia (an inherited condition in which cholesterol cannot be removed from the body normally).

Lipitor (atorvastatin) belongs to a class of medications called HMG-CoA reductase inhibitors (statins). It works by slowing the production of cholesterol in the body to decrease the amount of cholesterol that may build up on the walls of the arteries and block blood flow to the heart, brain, and other parts of the body.

Accumulation of cholesterol and fats along the walls of your arteries (a process known as atherosclerosis) decreases blood flow and, therefore, the oxygen supply to your heart, brain, and other parts of your body. Lowering your blood level of cholesterol and fats with Lipitor has been shown to prevent heart disease, angina (chest pain), strokes, and heart attacks.

Warnings

You should not take Lipitor if you have liver disease or [cirrhosis](#).

Lipitor can cause the breakdown of muscle tissue, which can lead to [kidney failure](#). Call your doctor right away if you have unexplained [muscle pain](#), tenderness, or weakness especially if you also have [fever](#), unusual tiredness, or dark urine.

Serious drug interactions can occur when certain medicines are used together with Lipitor. Tell each of your healthcare

providers about all medicines you use now, and any medicine you start or stop using.

Avoid eating foods that are high in fat or cholesterol. Lipitor will not be as effective in lowering your cholesterol if you do not follow a cholesterol-lowering diet plan.

Lipitor is only part of a complete program of treatment that also includes diet, exercise, and weight control. Follow your diet, medication, and exercise routines very closely.

Before taking this medicine

You should not use Lipitor if you are allergic to atorvastatin, or if you have liver failure or cirrhosis.

To make sure Lipitor is safe for you, tell your doctor if you have ever had:

- muscle pain or weakness;
- diabetes;
- stroke;
- a thyroid disorder;
- a habit of drinking more than 2 alcoholic beverages per day; or
- kidney disease.

Lipitor can cause the breakdown of muscle tissue, which can lead to kidney failure. This happens more often in women, in older adults, or people who have kidney disease or poorly controlled hypothyroidism ([underactive thyroid](#)).

Atorvastatin may harm an unborn baby. Tell your doctor if you are pregnant.

Ask a doctor if it is safe to breastfeed while using Lipitor.

 [Lipitor pregnancy and breastfeeding warnings](#) (more detail)

How should I take Lipitor?

Take Lipitor exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Do not change your dose or stop taking any of your medications without your doctor's advice.

Lipitor is usually taken once per day. Follow your doctor's instructions.

You may take Lipitor tablets with or without food.

It may take up to 2 weeks before your cholesterol levels improve, and you may need frequent blood tests. Even if you have no symptoms, tests can help your doctor determine if Lipitor is effective.

Your treatment may also include diet, exercise, weight control, and blood tests.

Store at room temperature away from moisture, heat, and light.

 [Detailed Lipitor dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if you are more than 12 hours late for the dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Lipitor?

Avoid eating foods high in fat or cholesterol, or Lipitor will not be as effective.

Drinking alcohol may increase your risk of liver damage.

Grapefruit may interact with atorvastatin and cause side effects. Avoid consuming grapefruit products and drinking more than 1.2 liters of [grapefruit juice](#) each day.

Lipitor side effects

Get emergency medical help if you have **signs of an allergic reaction to Lipitor** (hives, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, [sore throat](#), burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Lipitor can cause the breakdown of muscle tissue, which can lead to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark urine.

Muscle problems may be more likely in older adults and those who have kidney problems, thyroid problems, or take certain other medicines.

Also call your doctor at once if you have:

- muscle weakness in your hips, shoulders, neck, and back;
- trouble lifting your arms, trouble climbing or standing;
- **liver problems** - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, [jaundice](#) (yellowing of the skin or eyes);
- **kidney problems** - swelling, urinating less, feeling tired or short of breath; or
- **high blood sugar** - increased thirst, increased urination, dry mouth, fruity breath odor.

Common Lipitor side effects may include:

- pain in your bones, spine, joints, or muscles;
- pain and burning when you urinate, painful urination;
- muscle spasms;
- [upset stomach](#);

- trouble with sleeping;
- stuffy nose, runny nose, sore throat;
- [diarrhea](#), [nausea](#); or
- pain in arms or legs (limbs).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Lipitor side effects](#) (more detail)

Related/similar drugs

Crestor

Crestor (rosuvastatin) is used to treat high cholesterol and high triglycerides in the blood ...

Reviews & ratings

5.6 / 10

[138 Reviews](#)

[View more](#)

Zetia

Zetia (ezetimibe) is used to treat high cholesterol. Includes Zetia side effects, interactions and ...

Reviews & ratings

4.3 / 10

[75 Reviews](#)

[View more](#)

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

[685 Reviews](#)

[View more](#)

Gemfibrozil

Gemfibrozil is used to treat high cholesterol and triglyceride levels in people with pancreatitis ...

Reviews & ratings

6.1 / 10

[27 Reviews](#)

[View more](#)

Pravastatin

Pravastatin (Pravachol) is a statin medicine used to lower cholesterol and triglycerides in the ...

Reviews & ratings

4.5 / 10

[78 Reviews](#)

[View more](#)

Fenofibrate

Fenofibrate is used to treat high cholesterol and high triglyceride levels. Learn about side ...

Reviews & ratings

5.7 / 10

[69 Reviews](#)

[View more](#)

Simvastatin

Simvastatin (Zocor) is a cholesterol-lowering medication that blocks the production of cholesterol ...

Reviews & ratings

4.8 / 10

[131 Reviews](#)

[View more](#)

What other drugs will affect Lipitor?

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which can increase risk of serious muscle problems or make the medicines less effective.

Tell your doctor about all your current medicines. Many drugs can interact with atorvastatin, especially:

- **other cholesterol lowering medicine** - [gemfibrozil](#), [niacin](#), [fenofibrate](#), [fenofibric acid](#), and others;
- [colchicine](#);
- **antibiotic or antifungal medicine** - [rifampin](#), [erythromycin](#), [clarithromycin](#), [itraconazole](#), [ketoconazole](#), [posaconazole](#), and [voriconazole](#);
- [birth control pills](#);
- medicine to prevent [organ transplant](#) rejection; or
- antiviral medicine to treat [hepatitis C](#) or HIV.

This list is not complete and many other drugs may interact with atorvastatin. This includes prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible drug interactions are listed here.

 [Lipitor drug interactions](#) (more detail)

Does Lipitor interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Lipitor

+

Enter a drug name

Add

Ingredients

Active ingredient: atorvastatin calcium

Inactive ingredients: calcium carbonate, USP; candelilla wax, croscarmellose sodium, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, Opadry White YS-1-7040 (hypromellose, polyethylene glycol, talc, titanium dioxide); polysorbate 80, simethicone emulsion.

Manufacturer

Distributed by: Parke-Davis, Division of Pfizer Inc., NY, NY 10017, USA.

Popular FAQ

Does Green Tea interact with any drugs?



Can you use limes or lime juice while taking statin drugs (i.e. atorvastatin)?



More FAQ

- [Do I need to eat food when taking atorvastatin?](#)
- [What are the side effects of statins?](#)
- [Does Lipitor work better if you take it at night?](#)
- [Can Lipitor affect libido?](#)
- [How long does atorvastatin stay in the system after stopping the drug?](#)

[View more FAQ...](#)

More about Lipitor (atorvastatin)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(102\)](#)
- [Drug images](#)
- [Latest FDA alerts \(9\)](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Support group](#)
- [Drug class: statins](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

[Atorvaliq](#)

Professional resources

- [Lipitor prescribing information](#)
- [Atorvastatin Calcium \(AHFS Monograph\)](#)

Other brands

[Atorvaliq](#)

Related treatment guides

- [High Cholesterol, Familial Homozygous](#)
- [Hyperlipoproteinemia Type IIa, Elevated LDL](#)
- [High Cholesterol, Familial Heterozygous](#)
- [Hyperlipoproteinemia](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Lipitor only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 23.02.H2

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	FDA approved 1996

User Reviews & Ratings

4.3 / 10

[102 Reviews](#)

Related News

[Two-Drug Combo of a Statin Plus Ezetimibe, Lowers Cholesterol Better Than Statins Alone](#)

Images

[Lipitor 10 mg \(PD 155 10\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.