

Home

Silenor

Silenor 🖘

Generic name: doxepin [DOX-e-pin]

Drug classes: Miscellaneous anxiolytics, sedatives and hypnotics, Tricyclic antidepressants

Medically reviewed by Philip Thornton, DipPharm. Last updated on Feb 19, 2025.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is Silenor?

Silenor is a tricyclic antidepressant.

Silenor tablets are used to treat insomnia in people who have trouble staying asleep.

This medication guide provides information about the Silenor brand of doxepin used to treat insomnia. Sinequan and other generic brands of doxepin used to treat depression or anxiety are not covered in this medication guide. Silenor tablets are not for use in treating depression.

Warnings

You should not use Silenor if you have untreated narrow-angle glaucoma or severe problems with urination.

Do not use doxepin if you have used an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine.

Some people using this medicine have engaged in activity such as driving, eating, or making phone calls and later having no memory of the activity. If this happens to you, stop taking Silenor and talk with your doctor.

Before taking this medicine

You should not use Silenor if you are allergic to doxepin, amoxapine, or loxapine, or if you have:

- untreated narrow-angle glaucoma; or
- · severe urination problems.

Do not use Silenor if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others.

Tell your doctor if you have used an "SSRI" antidepressant in the past 5 weeks, such as citalopram, escitalopram,

fluoxetine (Prozac), fluvoxamine, paroxetine, sertraline (Zoloft), trazodone, or vilazodone.

To make sure Silenor is safe for you, tell your doctor if you have ever had:

- sleep apnea (breathing stops during sleep);
- · depression, mental illness, or addiction to drugs or alcohol;
- · kidney or liver disease;
- · glaucoma; or
- · urination problems.

Ask your doctor about taking Silenor **if you are pregnant.** Taking an antidepressant during late pregnancy may cause medical problems in the baby. However, you may have a relapse of depression if you stop taking your antidepressant. Tell your doctor right away if you become pregnant. **Do not start or stop taking this medicine without your doctor's advice.**

This medicine may affect fertility (your ability to have children), whether you are a man or a woman.

You should not breastfeed while using Silenor.

Silenor is not approved for use by anyone younger than 18 years old.

Silenor pregnancy and breastfeeding warnings (more detail)

How should I take Silenor?

Take Silenor exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

Take this medicine within 30 minutes before bedtime. Do not take within 3 hours after eating.

It may take 7 to 10 days before your insomnia symptoms improve. Keep using as directed and call your doctor if your symptoms do not improve after 10 days of treatment.

Store at room temperature away from moisture, heat, and light. Keep the bottle tightly closed when not in use.

Detailed Silenor dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

Do not take Silenor if you do not have time for a full night's sleep before being active again.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of doxepin can be fatal.**

What to avoid

Do not drink alcohol. Dangerous side effects or death could occur.

Avoid taking Silenor within 3 hours after eating a meal.

Avoid driving or hazardous activity until you know how this medicine will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Silenor side effects

Get emergency medical help if you have **signs of an allergic reaction to Silenor:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, or making phone calls. If this happens to you, call your doctor right away.

Call your doctor at once if you have:

- · unusual thoughts or behavior;
- · confusion, hallucinations;
- depressed mood, thoughts about hurting yourself;
- · uncontrolled muscle movements;
- a seizure:
- a light-headed feeling, like you might pass out;
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing;
- · eye pain or redness;
- chest pain;
- low red blood cells (anemia) pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet; or
- signs of infection flu-like symptoms, swelling, skin redness, diarrhea, skin sores or white patches, trouble breathing, pain or burning when you urinate.

Common Silenor side effects may include:

- · drowsiness, tiredness;
- nausea; or
- cold symptoms such as stuffy nose, sneezing, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Silenor side effects (more detail)

Related/similar drugs

Ativan

Ativan is a benzodiazepine used to treat anxiety disorders or anxiety associated with depression ...

Reviews & ratings

8.0 / 10

479 Reviews

View more

Benadryl

Benadryl is an antihistamine and is used to treat sneezing, runny nose, itching and other ...

Reviews & ratings

7.0 / 10

79 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Ambien

Ambien (zolpidem) is a prescription sedative used to treat insomnia, helping you fall asleep and ...

Reviews & ratings

7.0 / 10

558 Reviews

View more

Lunesta

Lunesta is a sedative and is used to treat insomnia. Lunesta's effects typically begin within 30 ...

Reviews & ratings

5.9 / 10

341 Reviews

View more

Eszopiclone

Eszopiclone (brand name Lunesta) is a sedative and is used to treat insomnia. Eszopiclone's effects ...

Reviews & ratings

5.7 / 10

442 Reviews

View more

Temazepam

Temazepam is used to treat insomnia (trouble sleeping). Learn about side effects, interactions and ...

Reviews & ratings

6.5 / 10

448 Reviews

View more

Melatonin

Melatonin is used to combat jet lag and ease sleep problems like insomnia. Learn about uses ...

Reviews & ratings

5.7 / 10

239 Reviews

View more

Diphenhydramine

Diphenhydramine is an antihistamine used to treat sneezing, runny nose, itching, hives and other ...

Reviews & ratings

5.5 / 10

541 Reviews

View more

Zolpidem

Zolpidem is a sedative-hypnotic medicine used for the short-term treatment of insomnia (trouble ...

Reviews & ratings

6.9 / 10

1,055 Reviews

View more

What other drugs will affect Silenor?

Using Silenor with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- cimetidine (Tagamet); or
- tolazamide (Tolinase).

This list is not complete. Other drugs may interact with doxepin, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Silenor drug interactions (more detail)

Does Silenor interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Silenor
+
Enter a drug name
Add

Popular FAQ

More FAQ

• What are some common side effects of antidepressants?

References

1. Silenor Product Label

More about Silenor (doxepin)

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (127)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- · Generic availability
- FDA approval history
- Drug class: miscellaneous anxiolytics, sedatives and hypnotics
- Breastfeeding
- En español

Patient resources

Other brands

Sinequan

Professional resources

- Silenor prescribing information
- Doxepin (AHFS Monograph)

Other brands

Sinequan

Related treatment guides

• Insomnia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Silenor only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্য Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

The Drug history at FDA

User Reviews & Ratings

6.5 / 10

127 Reviews

Images

Silenor 3 mg (3 SP)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×