

Home

2. Livalo

Livalo 🖘

Generic name: pitavastatin [pit-AV-a-STAT-in]

Brand names: Livalo, Zypitamag

Drug class: Statins

Medically reviewed by Sanjai Sinha, MD. Last updated on Apr 15, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is Livalo?

Livalo belongs to a group of drugs called HMG CoA reductase inhibitors, or "statins."

Livalo is used together with diet to lower blood levels of "bad" cholesterol (low-density lipoprotein, or LDL) and triglycerides, to increase levels of "good" cholesterol (high-density lipoprotein, or HDL), and to lower triglycerides (a type of fat in the blood).

Livalo is used to treat high cholesterol in adults. Lowering your cholesterol may help prevent heart disease and hardening of the arteries, conditions that can lead to heart attack, stroke, and vascular disease.

Livalo is also used to treat an inherited form of high cholesterol in children at least 8 years old.

Warnings

You should not take Livalo if you have liver disease (including significantly abnormal liver enzyme tests), or if you also take cyclosporine.

Do not take Livalo if you are pregnant.

Do not breastfeed while you are taking Livalo.

Before taking this medicine

You should not use Livalo if you are allergic to pitavastatin, or if you have:

- liver disease (including significantly abnormal liver enzyme tests);
- if you are pregnant or breastfeeding; or
- if you are taking cyclosporine.

Do not use Livalo if you are pregnant. Pitavastatin can harm an unborn baby. Use effective birth control to prevent

pregnancy. Stop taking this medicine and tell your doctor at once if you become pregnant.

Pitavastatin can pass into breast milk and may harm a nursing baby. Do not breastfeed while you are taking Livalo.

To make sure Livalo is safe for you, tell your doctor if you have ever had:

- · kidney disease;
- liver disease;
- a thyroid disorder; or
- if you drink large amounts of alcohol.

Livalo can cause a condition that results in the breakdown of skeletal muscle tissue, potentially leading to kidney failure. This condition may be more likely to occur in older adults and in people who have kidney disease or poorly controlled hypothyroidism (underactive thyroid).

Do not give this medicine to a child without medical advice.

1 Livalo pregnancy and breastfeeding warnings (more detail)

How should I take Livalo?

Take Livalo exactly as prescribed by your doctor. Follow all directions on your prescription label. Your doctor may occasionally change your dose to make sure you get the best results. Do not take this medicine in larger or smaller amounts or for longer than recommended.

You may take Livalo with or without food, but take it at the same time each day.

While using Livalo, you may need frequent blood tests to check your liver function.

You may need to stop using Livalo for a short time if you have:

- uncontrolled seizures;
- an electrolyte imbalance (such as high or low potassium levels in your blood);
- · severely low blood pressure;
- · a severe infection or illness;
- · dehydration; or
- surgery or a medical emergency.

Livalo is only part of a complete program of treatment that also includes diet, exercise, and weight control. Follow your diet, medication, and exercise routines very closely.

Store at room temperature away from moisture, heat, and light.

i Livalo patient tips (more detail)

Dosing information

Usual Adult Dose for Hyperlipidemia:

Initial dose: 2 mg orally once a day with or without food

Maintenance dose: 1 to 4 mg once a day with or without food

Maximum dose: 4 mg once a day with or without food

Detailed Livalo dosage information

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What to avoid

Avoid eating foods that are high in fat or cholesterol. Livalo will not be as effective in lowering your cholesterol if you do not follow a cholesterol-lowering diet plan.

Drinking alcohol can can raise triglyceride levels and may increase your risk of liver damage.

Livalo side effects

Get emergency medical help if you have any **signs of an allergic reaction to Livalo**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

In rare cases, Livalo can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. **Call your doctor right away if** you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark urine.

Also call your doctor at once if you have:

- · confusion, memory problems;
- muscle weakness in your hips, shoulders, neck, and back;
- · trouble lifting your arms, trouble climbing or standing;
- kidney problems vomiting, pain in your side or lower back, little or no urinating, swelling, rapid weight gain; or
- **liver problems** loss of appetite, stomach pain (upper right side), tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Common Livalo side effects may include:

- muscle pain;
- pain in your arms or legs;

- · back pain; or
- diarrhea, constipation.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Livalo side effects (more detail)

Related/similar drugs

Lipitor

Lipitor is used to treat high cholesterol. Learn about side effects, interactions and indications.

Reviews & ratings

4.3 / 10

102 Reviews

View more

Crestor

Crestor (rosuvastatin) is used to treat high cholesterol and high triglycerides in the blood ...

Reviews & ratings

5.6 / 10

138 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Zetia

Zetia (ezetimibe) is used to treat high cholesterol. Includes Zetia side effects, interactions and ...

Reviews & ratings

4.3 / 10

75 Reviews

View more

Zocor

Zocor (simvastatin) reduces lowdensity lipoprotein cholesterol and total cholesterol in the blood ...

Reviews & ratings

5.0 / 10

22 Reviews

View more

Lescol

Lescol (fluvastatin) is used to treat high cholesterol which helps prevent heart disease, stroke ...

Reviews & ratings

2 Reviews

View more

What other drugs will affect Livalo?

Tell your doctor about all other medicines you use. Like pitavastatin, certain other drugs can increase your risk of serious muscle problems, and it is very important that your doctor knows if you are using any of them:

- colchicine;
- · gemfibrozil;
- · fenofibric acid;
- · fenofibrate; or
- medicines that contain niacin (Advicor, Niaspan, Niacor, Simcor, Slo-Niacin, and others).

This list is not complete. Other drugs may interact with pitavastatin, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

1 Livalo drug interactions (more detail)

Does Livalo interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Livalo	
+	
Enter a drug name	Add

Frequently asked questions

- · How is Livalo different from other statins?
- When is the best time of day to take Livalo?
- Can you drink alcohol while taking Livalo?
- · What are the side effects of statins?
- Do statins help treat COVID-19?

References

1. Livalo Product Label

More about Livalo (pitavastatin)

- Check interactions
- Compare alternatives
- · Pricing & coupons
- Reviews (92)
- Drug images
- Latest FDA alerts (2)
- Side effects
- Dosage information
- Patient tips
- During pregnancy
- · Generic availability
- FDA approval history
- · Drug class: statins
- Breastfeeding
- En español

Patient resources

Other brands

Zypitamag, Nikita

Professional resources

- · Livalo prescribing information
- Pitavastatin Calcium, Pitavastatin Magnesium (AHFS Monograph)

Other brands

Zypitamag

Related treatment guides

- Dyslipidemia
- High Cholesterol

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Livalo only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ঝ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 2009

User Reviews & Ratings

5.4 / 10

92 Reviews

Images

Livalo 2 mg (KC 2)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







