

Home

Anaprox-DS

Anaprox-DS

Generic name: naproxen [na-PROX-en]

Brand names: Aleve, Aleve Back and Muscle Pain, Aleve Easy Open Arthritis, Aleve Liquid Gels, Anaprox-DS, ... show all 10 brands

Drug class: Nonsteroidal anti-inflammatory drugs



Medically reviewed by Drugs.com on Aug 3, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Anaprox-DS?

Anaprox-DS is a nonsteroidal anti-inflammatory drug (NSAID).

Anaprox-DS is used to treat pain or inflammation caused by conditions such as arthritis, ankylosing spondylitis, tendinitis, bursitis, gout, or menstrual cramps.

The **delayed-release** or **extended-release** tablets are slower-acting forms of naproxen that are used only for treating chronic conditions such as arthritis or ankylosing spondylitis. **These forms of naproxen will not work fast enough to treat acute pain.**

Anaprox-DS may also be used for purposes not listed in this medication guide.

Anaprox-DS side effects

Get emergency medical help if you have **signs of an allergic reaction** (runny or stuffy nose, wheezing or trouble breathing, hives, swelling in your face or throat) **or a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Stop using Anaprox-DS and seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes.

Get emergency medical help if you have **signs of a heart attack or stroke**: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath.

Anaprox-DS may cause serious side effects. Stop using Anaprox-DS and call your doctor at once if you have:

- shortness of breath (even with mild exertion);
- · swelling or rapid weight gain;
- the first sign of any skin rash or blister, no matter how mild;

- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- **liver problems**--nausea, upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- kidney problems--little or no urination, painful urination, swelling in your feet or ankles; or
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet.

Common side effects of Anaprox-DS may include:

- · headache:
- · indigestion, heartburn, stomach pain; or
- flu symptoms;

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Anaprox-DS side effects (more detail)

Related/similar drugs

Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

Reviews & ratings

7.0 / 10

223 Reviews

View more

Propranolol

Propranolol is a beta-blocker that is used to treat tremors, chest pain, high blood pressure, heart ...

Reviews & ratings

8.2 / 10

1,376 Reviews

View more

FEATURED

Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used

alone or with other ...

Reviews & ratings

5.7 / 10

145 Reviews

View more

Amitriptyline

Amitriptyline is a tricyclic antidepressant used to treat depression and, off-label, conditions ...

Reviews & ratings

7.5 / 10

1,803 Reviews

View more

Tramadol

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

Reviews & ratings

7.0 / 10

1,876 Reviews

View more

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

1,404 Reviews

View more

Duloxetine

Duloxetine is a selective serotonin and norepinephrine reuptake inhibitor antidepressant used to ...

Reviews & ratings

6.2 / 10

View more

Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

Reviews & ratings

6.0 / 10

618 Reviews

View more

Meloxicam

Meloxicam is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.3 / 10

600 Reviews

View more

Aspirin

Aspirin is used to treat mild to moderate pain and to reduce fever or inflammation. Learn about ...

Reviews & ratings

7.8 / 10

51 Reviews

View more

Warnings

Anaprox-DS can increase your risk of fatal heart attack or stroke. Do not use Anaprox-DS just before or after heart bypass surgery (coronary artery bypass graft, or CABG). This medicine may also cause stomach or intestinal bleeding, which can be fatal.

Before taking this medicine

Anaprox-DS can increase your risk of fatal heart attack or stroke, even if you don't have any risk factors. Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Anaprox-DS may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are using this medicine, especially in older adults.

You should not use Anaprox-DS if you are allergic to it, or if you have ever had an asthma attack or severe allergic reaction after taking aspirin or an NSAID.

Ask a doctor before giving Anaprox-DS to a child younger than 12 years old.

Ask a doctor or pharmacist if this medicine is safe to use if you have:

- heart disease, high blood pressure, high cholesterol, diabetes, or if you smoke;
- a heart attack, stroke, or blood clot;
- · stomach ulcers or bleeding;
- asthma;
- liver or kidney disease;
- fluid retention; or
- if you take aspirin to prevent heart attack or stroke.

If you are pregnant, you should not take Anaprox-DS unless your doctor tells you to. Taking an NSAID during the last 20 weeks of pregnancy can cause serious heart or kidney problems in the unborn baby and possible complications with your pregnancy.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk.

How should I take Anaprox-DS?

Use exactly as directed on the label, or as prescribed by your doctor. Use the lowest dose that is effective in treating your condition.

Shake the **oral suspension** (liquid) before you measure a dose. Measure a dose with the supplied measuring device (not a kitchen spoon).

Take Anaprox-DS with food or milk if it upsets your stomach.

Always follow directions on the medicine label about giving this medicine to a child. Anaprox-DS doses are based on weight in children. Your child's dose needs may change if the child gains or loses weight.

If you use Anaprox-DS long-term, you may need frequent medical tests.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Anaprox-DS.

Store at room temperature away from moisture, heat, and light. Keep the bottle tightly closed when not in use.

Detailed Anaprox-DS dosage information

What happens if I miss a dose?

Since Anaprox-DS is used when needed, you may not be on a dosing schedule. Skip any missed dose if it's almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Anaprox-DS?

Avoid drinking alcohol. It may increase your risk of stomach bleeding.

Avoid taking aspirin or other NSAIDs unless your doctor tells you to.

Ask a doctor or pharmacist before using other medicines for pain, fever, swelling, or cold/flu symptoms. They may contain ingredients similar to Anaprox-DS (such as aspirin, ibuprofen, or ketoprofen).

Ask your doctor before using an antacid, and use only the type your doctor recommends. Some antacids can make it harder for your body to absorb Anaprox-DS.

What other drugs will affect Anaprox-DS?

Ask your doctor before using Anaprox-DS if you take an antidepressant. Taking certain antidepressants with an NSAID may cause you to bruise or bleed easily.

Ask a doctor or pharmacist before using Anaprox-DS with any other medications, especially:

- other NSAIDs or salicylates (diflunisal, salsalate);
- antacids and sucralfate;
- cholestyramine;
- cyclosporine;
- digoxin;
- · lithium;
- · methotrexate;
- pemetrexed;
- · probenecid;
- warfarin (Coumadin, Jantoven) or similar blood thinners;
- a diuretic or "water pill"; or
- · heart or blood pressure medication.

This list is not complete. Other drugs may affect Anaprox-DS, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Does Anaprox-DS interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Anaprox-DS	
+	
Enter a drug name	Add

Popular FAQ

Naproxen vs ibuprofen: What's the difference?	~
What is Caridoxen used for?	~
How long does naproxen stay in your system?	~
Does Feverfew interact with any drugs?	~
What is Apronax called in the USA?	V

More FAQ

- Which painkiller should you use?
- What is the best way to reduce swelling in your face?
- Can NSAIDs be used to treat a COVID-19 fever?

More about Anaprox-DS (naproxen)

- · Check interactions
- Compare alternatives
- Reviews (4)
- Drug images
- Latest FDA alerts (8)
- Side effects
- Dosage information
- During pregnancy
- · Generic availability

- Drug class: Nonsteroidal anti-inflammatory drugs
- Breastfeeding
- En español

Patient resources

Anaprox DS advanced reading

Other brands

Aleve, Naprosyn, Anaprox, Aleve Back and Muscle Pain, ... +5 more

Professional resources

- Naproxen monograph
- Naproxen Tablets (FDA)

Other brands

Aleve, Naprosyn, Naprelan, All Day Pain Relief, All Day Relief

Related treatment guides

- Ankylosing Spondylitis
- Aseptic Necrosis
- Back Pain
- Bursitis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 22.01.

DRUG STATUS

Availability

of Rx and/or OTC


Pregnancy & Lactation

Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History



User Reviews & Ratings

4 Reviews

Images

Anaprox DS naproxen sodium 550 mg (NPS 550)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









