



[Home](#)

2. [Levoxyl](#)

Levoxyl

Generic name: [levothyroxine](#) [LEE-voe-thye-ROX-een]

Drug class: [Thyroid drugs](#)

Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on Sep 26, 2023.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is Levoxyl?

Levoxyl is a [thyroid medicine](#) that replaces a hormone normally produced by your thyroid gland to regulate the body's energy and metabolism. Levothyroxine is given when your thyroid does not produce enough of this hormone on its own.

Levoxyl is a prescription medicine used to treat [hypothyroidism \(low thyroid hormone\)](#).

Levoxyl is also used to treat or prevent [goiter \(enlarged thyroid gland\)](#), which can be caused by hormone imbalances, radiation treatment, surgery, or cancer.

Levoxyl is also used in patients who need surgery and radioiodine therapy to manage a type of thyroid cancer called well-differentiated thyroid cancer.

Warnings

You may not be able to take Levoxyl if you have certain medical conditions. Tell your doctor if you have an untreated or uncontrolled adrenal gland disorder, a thyroid disorder called [thyrotoxicosis](#), or if you have any recent or current symptoms of a [heart attack](#).

Levoxyl should not be used to treat obesity or weight problems. Dangerous side effects or death can occur from the misuse of this medicine, especially if you are taking any other weight-loss medications or appetite suppressants.

Before taking this medicine

Levoxyl should not be used to treat obesity or weight problems. Dangerous side effects or death can occur from the misuse of levothyroxine, especially if you are taking any other weight-loss medications or appetite suppressants.

Since thyroid hormone occurs naturally in the body, almost anyone can take Levoxyl. However, you may not be able to take this medicine if you have certain medical conditions.

To make sure Levoxyl is safe for you, tell your doctor if you have ever had:

- an untreated or uncontrolled adrenal gland disorder;
- a thyroid disorder called thyrotoxicosis; or
- symptoms of a heart attack (chest pain or heavy feeling, pain spreading to the jaw or shoulder, [nausea](#), sweating, general ill feeling).
- a thyroid nodule;
- heart problems such as a heart attack, stroke, blood clot or blood clotting disorder;
- diabetes (your diabetes medicine may need to be adjusted);
- anemia (low red blood cells);
- weak bones (osteoporosis), or low bone mineral density;
- problems with your pituitary gland;
- any food or drug allergies;
- plan to have surgery;
- recently received radiation therapy with iodine (such as I-131); or
- kidney disease.

If you become pregnant while taking Levoxyl, do not stop taking the medicine without your doctor's advice. Having low thyroid hormone levels during pregnancy could harm both mother and baby. **Your dose needs may be different during pregnancy.**

Tell your doctor if you are breast-feeding. **Your dose needs may be different while you are nursing.**

Do not give Levoxyl to a child without medical advice.

 [Levoxyl pregnancy and breastfeeding warnings](#) (more detail)

How should I take Levoxyl?

Take Levoxyl exactly as it was prescribed for you. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Levoxyl is taken by mouth.

Levoxyl works best if you take it on an empty stomach, 30 to 60 minutes before breakfast. Follow your doctor's dosing instructions and try to take the medicine at the same time each day.

Swallow the tablet whole, with a full glass (8 ounces) of water. The Levoxyl tablet may dissolve very quickly and could swell in your throat.

When giving Levoxyl to infants and children who cannot swallow tablets the tablets may be crushed. Place the freshly crushed tablet in a small amount (5 mL to 10 mL or 1 to 2 teaspoonfuls) of water and immediately give the medicine by spoon or dropper. Do not store the mixed medicine or mix it with foods that decrease the absorption of levothyroxine, such as soybean-based infant formula.

It may take several weeks before your body starts to respond to levothyroxine. **Keep using Levoxyl even if you feel well.** You may need to use for the rest of your life.

You may need frequent medical tests. Tell any doctor, dentist, or surgeon who treats you that you are using .

Store at room temperature away from moisture and heat.

Do not share Levoxyl with another person, even if they have the same symptoms you have.

 [Detailed Levoxyl dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include [headache](#), leg cramps, tremors, feeling nervous or irritable, chest pain, shortness of breath, and fast or pounding heartbeats.

What to avoid

Avoid the following food products, which can make your body absorb less levothyroxine: [grapefruit juice](#), infant soy formula, soybean flour, cotton seed meal, walnuts, and high-fiber foods.

side effects

Get emergency medical help if you have **signs of an allergic reaction to Levoxyl**: [hives](#); difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- fast or irregular heartbeats;
- chest pain, pain spreading to your jaw or shoulder;
- shortness of breath;
- fever, [hot flashes](#), sweating;
- tremors, or if you feel unusually cold;
- weakness, tiredness, sleep problems ([insomnia](#));
- memory problems, feeling depressed or irritable;
- headache, leg cramps, muscle aches;
- feeling nervous or irritable;
- dryness of your skin or hair, [hair loss](#);
- irregular menstrual periods; or

- [vomiting](#), [diarrhea](#), appetite changes, weight changes.

Certain side effects may be more likely in older adults.

Common Levoxyl side effects may include:

- chest pain, irregular heartbeats;
- shortness of breath;
- headache, leg cramps, muscle pain or weakness;
- tremors, feeling nervous or irritable, trouble sleeping;
- increased appetite;
- feeling hot;
- [weight loss](#);
- changes in your menstrual periods;
- diarrhea; or
- [skin rash](#), partial hair loss.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Levoxyl side effects](#) (more detail)

Related/similar drugs

Synthroid

Synthroid (levothyroxine) treats hypothyroidism (low thyroid hormone) and different types of ...

Reviews & ratings

6.1 / 10

[191 Reviews](#)

[View more](#)

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

[685 Reviews](#)

[View more](#)

Euthyrox

Euthyrox (levothyroxine) is used to treat hypothyroidism and to treat or prevent goiter. Includes ...

Reviews & ratings

1.9 / 10

[164 Reviews](#)

[View more](#)

What other drugs will affect ?

Many other medicines can be affected by your thyroid hormone levels. Certain other medicines may also increase or decrease the effects of levothyroxine.

Certain medicines can make levothyroxine less effective if taken at the same time. If you use any of the following drugs, avoid taking them within 4 hours before or 4 hours after you take :

- [calcium carbonate](#) ([Alka-Mints](#), Caltrate, [Os-Cal](#), Oyster Shell Calcium, [Rolaids Soft Chew](#), Tums, and others);
- [cholestyramine](#), [colesevelam](#), [colestipol](#);
- [ferrous sulfate](#) iron supplement;

- [sucralfate](#);
- [sodium polystyrene sulfonate](#) (Kalexate, [Kayexalate](#), [Kionex](#));
- **stomach acid reducers** - [esomeprazole](#), [lansoprazole](#), [omeprazole](#), [rabeprazole](#), [Nexium](#), [Prilosec](#), [Prevacid](#), [Protonix](#), [Zegerid](#), and others; or
- **antacids that contain aluminum or magnesium** - Gaviscon, Maalox, [Milk of Magnesia](#), [Mintox](#), [Mylanta](#), [Pepcid Complete](#), and others.

Many drugs can interact with levothyroxine. This includes prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible interactions are listed here. Tell your doctor about all your current medicines and any medicine you start or stop using.

 [Levoxyl drug interactions](#) (more detail)









Does Levoxyl interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

+

Add

Popular FAQ

- What is the difference between Levoxyl and Synthroid? 
- Who makes Levoxyl and where is it manufactured? 
- Is Levoxyl a brand or a generic? 
- Does levothyroxine cause weight gain or loss? 
- What can I eat for breakfast after taking levothyroxine? 
- Can I take other medications with levothyroxine? 
- Does levothyroxine cause hair loss? 
- How long after taking levothyroxine can you drink milk? 

More FAQ

- [Should you take your thyroid medication at night?](#)

[View more FAQ...](#)

References

1. [Levoxyl Product Label](#)

More about Levoxyl (levothyroxine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(38\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: thyroid drugs](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

[Synthroid](#), [Tirosint](#), [Euthyrox](#), [Unithroid](#), ... [+3 more](#)

Professional resources

- [Levoxyl prescribing information](#)
- [Levothyroxine \(AHFS Monograph\)](#)

Other brands

[Synthroid](#), [Tirosint](#), [Euthyrox](#), [Unithroid](#), ... [+5 more](#)

Related treatment guides

- [Hashimoto's disease](#)
- [TSH Suppression](#)
- [Hypothyroidism, After Thyroid Removal](#)
- [Myxedema Coma](#)
- [Thyroid Suppression Test](#)
- [Underactive Thyroid](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Levoxyl only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 17.01.

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	Drug history at FDA


User Reviews & Ratings

6.5 / 10

[38 Reviews](#)

Images

[Levoxyl 50 mcg \(0.05 mg\) \(LEVOXYL dp 50\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.