



[Home](#)

2. Hysingla ER

Hysingla ER

Generic name: [hydrocodone](#) (oral) [*HYE-droe-KOE-done*]

Drug class: [Opioids](#) (narcotic analgesics)

Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on Sep 25, 2023.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is Hysingla ER?

Hysingla ER is an opioid pain medication.

Hysingla ER is an extended-release tablet that is used for around-the-clock treatment of [severe pain](#).

Hysingla ER is not for use on an as-needed basis for pain.

Warnings

Hydrocodone can slow or stop your breathing. Never use Hysingla ER in larger amounts, or for longer than prescribed. Do not crush, break, or open an extended-release pill. Swallow it whole to avoid exposure to a potentially fatal dose.

Hydrocodone may be habit-forming, even at regular doses. Never share Hysingla ER with another person, especially someone with a history of drug abuse or addiction. Keep the medication in a place where others cannot get to it.

MISUSE OF OPIOID MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep Hysingla ER in a place where others cannot get to it.

Tell your doctor if you are pregnant. **Hydrocodone may cause life-threatening withdrawal symptoms in a newborn if the mother has taken this medicine during pregnancy.**

Fatal side effects can occur if you use opioid medicine with alcohol, or with other drugs that cause drowsiness or slow your breathing.

Before taking this medicine

You should not use Hysingla ER if you are allergic to hydrocodone, or if you have:

- severe [asthma](#) or breathing problems; or
- a blockage in your stomach or intestines.

To make sure Hysingla ER is safe for you, tell your doctor if you have ever had:

- breathing problems, sleep apnea;
- a [head injury](#), [brain tumor](#), or [seizures](#);
- drug or alcohol addiction, or mental illness;
- urination problems;
- liver or kidney disease;
- problems with your gallbladder, pancreas, or thyroid; or
- a heart rhythm disorder called [long QT syndrome](#).

If you use opioid medicine while you are pregnant, your baby could become dependent on the drug. This can cause life-threatening withdrawal symptoms in the baby after it is born. Babies born dependent on [opioids](#) may need medical treatment for several weeks.

Ask a doctor before using Hysingla ER if you are breastfeeding. Tell your doctor if you notice severe drowsiness or slow breathing in the nursing baby.

 [Hysingla ER pregnancy and breastfeeding warnings](#) (more detail)

How should I take Hysingla ER?

Hysingla ER may be habit-forming, even at regular doses. Follow the directions on your prescription label and read all medication guides. Never use Hysingla ER in larger amounts, or for longer than prescribed. Tell your doctor if you feel an increased urge to take more of this medicine.

Never share opioid medicine with another person, especially someone with a history of drug abuse or addiction. MISUSE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medication in a place where others cannot get to it. Selling or giving away opioid medicine is against the law.

Your dose needs may be different if you have recently used a similar opioid pain medicine and your body is tolerant to it. Talk with your doctor if you are not sure you are opioid-tolerant.

Do not crush, break, or open an **extended-release tablet**. Swallow the tablet whole to avoid exposure to a potentially fatal dose. **Never crush or break a Hysingla ER tablet to inhale the powder or mix it into a liquid to inject the drug into your vein.** This can cause death.

Do not stop using this medicine suddenly after long-term use, or you could have serious withdrawal symptoms. Ask your doctor how to safely stop using Hysingla ER.

Store at room temperature, away from heat, moisture, and light. Keep track of your medicine. You should be aware if anyone is using it improperly or without a prescription.

Do not keep leftover opioid medication. **Just one dose can cause death in someone using this medicine accidentally or improperly.** Ask your pharmacist where to locate a drug take-back disposal program. If there is no take-back program, flush the unused medicine down the toilet.

Dosing information

Usual Adult Dose for Chronic Pain:

The following dosing recommendations can only be considered suggested approaches to what is actually a series of clinical decisions over time; each patient should be managed individually.

As First Opioid Analgesic and For Patients who are NOT Opioid Tolerant:

Extended-Release Hysingla ER Tablets: Initial dose: 20 mg orally every 24 hours

Comments:

-Use of higher starting doses in patients who are not opioid tolerant may cause fatal respiratory depression; monitor patients closely for respiratory depression, especially during the first 24 to 72 hours.

-An opioid tolerant patient is one who has been receiving for 1-week or longer at least: oral morphine 60 mg/day, fentanyl transdermal patch 25 mcg per hour, oral oxycodone 30 mg/day, oral hydromorphone 8 mg/day, oral oxymorphone 25 mg/day, or an equianalgesic dose of another opioid.

 [Detailed Hysingla ER dosage information](#)

What happens if I miss a dose?

Since Hysingla ER is used for pain, you are not likely to miss a dose. Skip any missed dose if it is almost time for your next dose. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **A hydrocodone overdose can be fatal, especially in a child or other person using the medicine without a prescription.** Overdose symptoms may include severe drowsiness, pinpoint pupils, slow breathing, or no breathing.

Your doctor may recommend you get naloxone (a medicine to reverse an opioid overdose) and keep it with you at all times. A person caring for you can give the naloxone if you stop breathing or don't wake up. Your caregiver must still get emergency medical help and may need to perform CPR (cardiopulmonary resuscitation) on you while waiting for help to arrive.

Anyone can buy naloxone from a pharmacy or local health department. Make sure any person caring for you knows where you keep naloxone and how to use it.

What to avoid

Do not drink alcohol. Dangerous side effects or death could occur.

Avoid driving or hazardous activity until you know how Hysingla ER will affect you. [Dizziness](#) or drowsiness can cause falls, accidents, or severe injuries.

Hysingla ER side effects

Get emergency medical help if you have **signs of an allergic reaction to Hysingla ER**: [hives](#); difficult breathing; swelling of your face, lips, tongue, or throat.

Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up.

Stop using Hysingla ER and call your doctor at once if you have:

- noisy breathing, sighing, shallow breathing, breathing that stops during sleep;
- a slow heart rate or weak pulse;
- pain or burning when you urinate;
- confusion, tremors, severe drowsiness;
- a light-headed feeling, like you might pass out;
- **low cortisol levels**--nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness; or
- **high levels of serotonin in the body** - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea.

Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: [agitation](#), hallucinations, fever, sweating, shivering, [fast heart rate](#), muscle stiffness, twitching, loss of coordination, nausea, vomiting, or [diarrhea](#).

Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent.

Common Hysingla ER side effects may include:

- [constipation](#), nausea, vomiting;
- dizziness, drowsiness, feeling tired;
- [headache](#); or
- [cold symptoms](#) such as stuffy nose, sneezing, [sore throat](#).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Hysingla ER side effects](#) (more detail)

Related/similar drugs

Oxycodone

Oxycodone is an opioid analgesic used to treat moderate to severe pain; it has a high potential for ...

Reviews & ratings

6.9 / 10

[1,142 Reviews](#)

[View more](#)

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

5.8 / 10

[30 Reviews](#)

[View more](#)

FEATURED

Omvox

Omvox is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Naproxen

Naproxen is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

[683 Reviews](#)

[View more](#)

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-the-counter painkiller and fever ...

Reviews & ratings

6.1 / 10

[13 Reviews](#)

[View more](#)

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

[1,404 Reviews](#)

[View more](#)

Tramadol

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

Reviews & ratings

7.0 / 10

[1,876 Reviews](#)

[View more](#)

Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

Reviews & ratings

6.0 / 10

[618 Reviews](#)

[View more](#)

Meloxicam

Meloxicam is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.3 / 10

[600 Reviews](#)

[View more](#)

Aspirin

Aspirin is used to treat mild to moderate pain and to reduce fever or

inflammation. Learn about ...

Reviews & ratings

7.8 / 10

[51 Reviews](#)

[View more](#)

What other drugs will affect Hysingla ER?

You may have breathing problems or withdrawal symptoms if you start or stop taking certain other medicines. Tell your doctor if you also use an [antibiotic](#), antifungal medication, heart or blood pressure medication, seizure medication, or medicine to treat [HIV](#) or [hepatitis C](#).

Opioid medication can interact with many other drugs and cause dangerous side effects or death. Be sure your doctor knows if you also use:

- cold or allergy medicines, bronchodilator asthma/COPD medication, or a diuretic ("water pill");
- medicines for motion sickness, irritable bowel syndrome, or overactive bladder;
- **other opioids** - opioid pain medicine or prescription cough medicine;
- **a sedative like Valium** - diazepam, alprazolam, lorazepam, Xanax, Klonopin, Versed, and others;
- **drugs that make you sleepy or slow your breathing** - a sleeping pill, muscle relaxer, medicine to treat mood disorders or mental illness; or
- **drugs that affect serotonin levels in your body** - a stimulant, or medicine for depression, Parkinson's disease, migraine headaches, serious infections, or nausea and vomiting.

This list is not complete. Other drugs may interact with hydrocodone, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible interactions are listed here.

 [Hysingla ER drug interactions](#) (more detail)

Does Hysingla ER interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Hysingla ER

+

Enter a drug name

Add

Popular FAQ

Which drugs cause opioid-induced constipation?



More FAQ

- [Oxycodone vs Hydrocodone - How do they compare?](#)
- [Which painkiller should you use?](#)
- [Norco vs Vicodin - How do they compare?](#)

References

1. [Hysingla ER Product Label](#)

More about Hysingla ER (hydrocodone)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(98\)](#)
- [Drug images](#)
- [Latest FDA alerts \(3\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [FDA approval history](#)
- [Drug class: Opioids \(narcotic analgesics\)](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

[Zohydro ER](#), [Vantrela ER](#)

Professional resources

- [Hysingla ER prescribing information](#)
- [HYDROcodone Bitartrate \(Antitussive\) \(AHFS Monograph\)](#)

Related treatment guides

- [Pain](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Hysingla ER only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
2	High potential for abuse
Approval History	
	Drug history at FDA

User Reviews & Ratings

4.2 / 10

[98 Reviews](#)

Images

[Hysingla ER 20 mg \(HYD 20\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.