

Home

2. Linezolid

Linezolid (oral/injection) ⊲ ¬

Generic name: linezolid (oral/injection) [lin-EZ-oh-lid]

Brand name: Zyvox

Dosage forms: intravenous solution (2 mg/mL-D5%; 2 mg/mL-NaCl 0.9%), oral powder for reconstitution (100 mg/5 mL), ... show all 3

dosage forms

Drug class: Oxazolidinone antibiotics



Medically reviewed by Drugs.com on Jun 4, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is linezolid?

Linezolid is an antibiotic that fights bacteria in the body. Linezolid is also an MAO (monoamine oxidase) inhibitor.

Linezolid is used to treat different types of bacterial infections, such as pneumonia, skin infections, and infections that are resistant to other antibiotics.

Linezolid may also be used for purposes not listed in this medication guide.

Linezolid side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Linezolid may cause serious side effects. Call your doctor at once if you have:

- vision problems, changes in color vision;
- severe stomach pain, diarrhea that is watery or bloody;
- a seizure;
- sweating, feeling anxious or shaky (may be signs of low blood sugar);
- lactic acidosis--unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired; or
- low blood cell counts--fever, chills, tiredness, weakness, confusion, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Common side effects of linezolid may include:

- · nausea, vomiting, diarrhea;
- · rash;
- anemia (low red blood cells); or
- · headache, dizziness.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Linezolid side effects (more detail)

Related/similar drugs

Augmentin

Augmentin is a prescription antibiotic combining amoxicillin and clavulanate to treat bacterial ...

Reviews & ratings

5.4 / 10

447 Reviews

View more

Vancomycin

Vancomycin is used to treat infections of the intestines that cause colitis.

Learn about side ...

Reviews & ratings

6.5 / 10

57 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Amoxicillin/clavulanate

Amoxicillin and clavulanate potassium is a combination antibiotic used to treat bacterial ...

Reviews & ratings

5.3 / 10

910 Reviews

View more

Clindamycin

Clindamycin (Cleocin) is used to treat serious infections caused by bacteria. Includes clindamycin ...

Reviews & ratings

5.7 / 10

806 Reviews

View more

Levofloxacin

Levofloxacin is a fluoroquinolone antibiotic used to treat serious bacterial infections and prevent ...

Reviews & ratings

5.1 / 10

741 Reviews

View more

Metronidazole

Metronidazole is an antibiotic used to fight bacteria in your body. Learn about side effects ...

Reviews & ratings

6.0 / 10

2,431 Reviews

View more

Ciprofloxacin is an antibiotic belong to a group of drugs called fluoroquinolones. Learn about side ...

Reviews & ratings

5.0 / 10

935 Reviews

View more

Warnings

Some medicines can cause unwanted or dangerous effects when used with linezolid. Tell your doctor about all other medicines you use.

Do not use linezolid if you have used an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, or tranylcypromine.

Before taking this medicine

You should not use linezolid if you are allergic to it.

Do not use linezolid if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, tranylcypromine, and others.

Tell your doctor if you also take stimulant medicine, opioid medicine, herbal products, or medicine for depression, mental illness, Parkinson's disease, migraine headaches, serious infections, or prevention of nausea and vomiting. **An interaction with linezolid could cause a serious condition called serotonin syndrome.**

Tell your doctor if you have ever had:

- high blood pressure;
- · a thyroid disorder;
- a carcinoid tumor;
- bone marrow suppression or a weak immune system;
- kidney or liver disease;
- pheochromocytoma (adrenal gland tumor);
- · diabetes;
- seizures; or
- if you use a catheter.

Tell your doctor if you are pregnant or breastfeeding.

If you are breastfeeding, tell your doctor if you notice diarrhea or vomiting in the nursing baby.

Linezolid liquid may contain phenylalanine and could be harmful if you have phenylketonuria (PKU).

How is linezolid used?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Linezolid oral is taken by mouth.

Linezolid tablets or liquid can be taken with or without food.

Gently mix the **oral suspension** (liquid) by turning the bottle upside down 3 to 5 times. **Do not shake.** Measure a dose with the supplied syringe or a dose-measuring device (not a kitchen spoon).

Linezolid injection is given in a vein.

Read and follow all Instructions for Use. Ask your doctor or pharmacist if you need help.

Prepare an injection only when you are ready to give it. Call your pharmacist if the medicine has particles in it.

Do not reuse a needle or syringe. Place them in a puncture-proof "sharps" container and dispose of it following state or local laws. Keep out of the reach of children and pets.

You will need frequent medical tests. Your vision and blood pressure may also need to be checked often.

Keep using linezolid even if your symptoms quickly improve. Skipping doses could make your infection resistant to medication. Linezolid will not treat a viral infection (flu or a common cold).

Store all forms of linezolid at room temperature away from moisture, heat, and light. Do not freeze. Throw away any **liquid** not used within 21 days.

Detailed Linezolid dosage information

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while using linezolid?

Avoid taking anti-diarrhea medicine without first asking your doctor. Diarrhea may be a sign of a new infection.

Eating tyramine while you are using linezolid can raise your blood pressure to dangerous levels. Avoid foods that have a high level of tyramine, such as:

· aged cheeses or meats;

- pickled or fermented meats, smoked or air-dried meats;
- · sauerkraut;
- · soy sauce;
- tap beer (alcoholic and nonalcoholic);
- · red wine; or
- any meat, cheese, or other protein-based food that has been improperly stored.

You should be very familiar with the list of foods you must avoid while you are using linezolid.

What other drugs will affect linezolid?

Many drugs can affect linezolid, and some drugs should not be used at the same time. Tell your doctor about all other medicines you use. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here.

Linezolid drug interactions (more detail)

Does linezolid interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



More about linezolid

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (32)
- · Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: oxazolidinone antibiotics
- Breastfeeding
- En español

Patient resources

- · Linezolid (Intravenous) advanced reading
- Linezolid (Oral) (Advanced Reading)
- Linezolid Tablets
- · Linezolid Injection
- Linezolid Suspension

Other brands

Zyvox

Professional resources

- Linezolid monograph
- Linezolid Injection (FDA)
- Linezolid Oral Suspension (FDA)
- Linezolid Tablets (FDA)

Other brands

Zyvox

Related treatment guides

- Methicillin-Resistant Staphylococcus Aureus Infection
- Bacterial Infection
- Nosocomial Pneumonia
- Bacteremia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.02.

Antibiotics 101

List of Common Antibiotics & Types

Antibiotics and Alcohol

Probiotics and Antibiotics

Antibiotics for Strep Throat

Antibiotics for a Tooth Infection

Antibiotics for UTI's

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

∾° Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

8.6 / 10

32 Reviews

Images

Linezolid 600 mg (I 22)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us














<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







 \times

Copyright © 2000-2025 Drugs.com. All rights reserved.