



[Home](#)

2. Wellbutrin

Wellbutrin

Generic name: [bupropion](#) [byoo-PRO-pee-on]

Brand names: [Wellbutrin SR](#), [Wellbutrin XL](#)

Drug class: [Miscellaneous antidepressants](#)

Medically reviewed by [Melisa Puckey, BPharm](#). Last updated on Aug 15, 2023.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)



The current formulations of this medicine are Wellbutrin XL and Wellbutrin SR. If generic versions of this product have been approved by the FDA, there may be [generic equivalents available](#).

See also: [Wellbutrin SR](#), [Wellbutrin XL](#)

What is Wellbutrin?

Wellbutrin is an antidepressant medicine that is thought to work in the brain and nerves on the chemical messengers norepinephrine and dopamine. Wellbutrin XL is used to treat major depressive disorder (MDD) and seasonal affective disorder, and Wellbutrin SR is only used to treat major depressive disorder (MDD).

Another brand of bupropion called Zyban is used to help to stop smoking, Wellbutrin is not indicated to help quit smoking.

Warnings

You should not take Wellbutrin if you have seizures, an eating disorder, or if you have suddenly stopped using alcohol, seizure medications, or sedatives. **If you take Wellbutrin for depression, do not also take Zyban to quit smoking.**

Do not use Wellbutrin if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine.

Wellbutrin may cause seizures, especially in people with certain medical conditions or when using certain drugs. Tell your doctor about all of your medical conditions and the drugs you use.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor will need to check your progress at regular visits while you are using this medicine. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks,

trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Before taking this medicine

Do not use Wellbutrin if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others.

You should not take Wellbutrin if you are allergic to bupropion, or if you have:

- a seizure disorder;
- an eating disorder such as anorexia or bulimia; or
- if you have suddenly stopped using alcohol, seizure medications, or a sedative (such as Xanax, Valium, Fiorinal, Klonopin, and others).

Do not use an MAO inhibitor within 14 days before or 14 days after you take bupropion. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Do not take Wellbutrin to treat more than one condition at a time. **If you take bupropion for depression, do not also take this medicine to quit smoking.**

Wellbutrin may cause seizures, especially if you have certain medical conditions or use certain drugs. Tell your doctor about all of your medical conditions and the drugs you use.

To make sure this medicine is safe for you, tell your doctor if you have ever had:

- a head injury, seizures, or brain or spinal cord tumor;
- narrow-angle glaucoma;
- heart disease, high blood pressure, or a heart attack;
- diabetes;
- kidney or liver disease (especially cirrhosis); or
- depression, bipolar disorder or other mental illness; or
- if you drink alcohol.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor will need to check your progress at regular visits while you are using Wellbutrin. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Ask your doctor about taking Wellbutrin **if you are pregnant.** It is not known whether bupropion will harm an unborn baby. However, you may have a relapse of depression if you stop taking your antidepressant. Tell your doctor right away if you become pregnant. **Do not start or stop taking bupropion without your doctor's advice.**

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of bupropion on the baby.

It may not be safe to breastfeed while using Wellbutrin. Ask your doctor about any risk.

 [Wellbutrin pregnancy and breastfeeding warnings](#) (more detail)

How should I take Wellbutrin?

Take Wellbutrin exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended. Too much of this medicine can increase your risk of a seizure.

Do not crush, chew, or break an extended-release tablet. Swallow it whole.

You should not change your dose or stop using Wellbutrin suddenly, unless you have a seizure while taking this medicine. Stopping suddenly can cause unpleasant withdrawal symptoms.

Do not stop using Wellbutrin suddenly, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine.

Some people taking Wellbutrin have had high blood pressure that is severe, especially when also using a nicotine replacement product (patch or gum). Your blood pressure may need to be checked before and during treatment with bupropion.

This medicine can cause you to have a false positive drug screening test. If you provide a urine sample for drug screening, tell the laboratory staff that you are taking bupropion.

Store at room temperature away from moisture and heat.

 [Detailed Wellbutrin dosage information](#)

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of bupropion can be fatal.** Overdose symptoms may include muscle stiffness, hallucinations, fast or uneven heartbeat, shallow breathing, or fainting.

What to avoid

Drinking alcohol with bupropion may increase your risk of seizures. If you drink alcohol regularly, talk with your doctor before changing the amount you drink. Bupropion can also cause seizures in people who drink a lot of alcohol and then suddenly quit drinking when they start using the medication.

Bupropion may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

Wellbutrin side effects

Get emergency medical help if you have any of these **signs of an allergic reaction to Wellbutrin**: (hives, itching, fever,

swollen glands, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, depression, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself.

Call your doctor at once if you have:

- a seizure (convulsions);
- confusion, unusual changes in mood or behavior;
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights;
- fast or irregular heartbeats; or
- **a manic episode** - racing thoughts, increased energy, reckless behavior, feeling extremely happy or irritable, talking more than usual, severe problems with sleep.

Common Wellbutrin side effects may include:

- dry mouth, sore throat, stuffy nose;
- ringing in the ears;
- blurred vision;
- nausea, vomiting, stomach pain, loss of appetite, constipation;
- sleep problems (insomnia);
- tremors, sweating, feeling anxious or nervous;
- fast heartbeats;
- confusion, agitation, hostility;
- rash;
- weight loss;
- increased urination;
- headache, dizziness; or
- muscle or joint pain.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Wellbutrin side effects](#) (more detail)

Related/similar drugs

Lexapro

Lexapro is used to treat anxiety and major depressive disorder. [Learn](#)

about side effects ...

Reviews & ratings

7.6 / 10

[2,376 Reviews](#)

[View more](#)

Zoloft

Zoloft is an antidepressant used to treat major depression, obsessive-compulsive disorder, panic ...

Reviews & ratings

7.3 / 10

[1,972 Reviews](#)

[View more](#)

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

[685 Reviews](#)

[View more](#)

Seroquel

Seroquel is used to treat the symptoms of schizophrenia, bipolar disorder and major depressive ...

Reviews & ratings

7.0 / 10

[694 Reviews](#)

[View more](#)

Prozac

Prozac (fluoxetine) is an SSRI antidepressant used to treat depression, OCD, panic disorder ...

Reviews & ratings

7.2 / 10

1.2 / 10

[1,239 Reviews](#)

[View more](#)

Cymbalta

Cymbalta (duloxetine) is used to treat major depressive disorder, general anxiety disorder and ...

Reviews & ratings

6.3 / 10

[1,899 Reviews](#)

[View more](#)

Celexa

Celexa (citalopram) is an antidepressant in a group of drugs called selective serotonin reuptake ...

Reviews & ratings

7.3 / 10

[866 Reviews](#)

[View more](#)

Concerta

Concerta is used to treat attention deficit disorder (ADD) and attention deficit hyperactivity ...

Reviews & ratings

6.9 / 10

[424 Reviews](#)

[View more](#)

Citalopram

Citalopram is an antidepressant (selective serotonin reuptake inhibitor) and is used to treat ...

Reviews & ratings

7.4 / 10

[2,231 Reviews](#)

[View more](#)

Escitalopram

Escitalopram is an SSRI antidepressant used to treat major depressive disorder in adults and teens ...

Reviews & ratings

7.5 / 10

[3,962 Reviews](#)

[View more](#)

What other drugs will affect Wellbutrin?

You may have a higher risk of seizures if you use certain other medicines while taking Wellbutrin.

Many drugs can interact with bupropion. Tell your doctor about all medicines you use, and those you start or stop using during your treatment with Wellbutrin. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

 [Wellbutrin drug interactions](#) (more detail)

Does Wellbutrin interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).






Wellbutrin

+

Enter a drug name

Add

Popular FAQ

- Does Wellbutrin XL/SR cause weight gain or loss? 
- Can I stop taking Wellbutrin suddenly or should I taper off? 
- How long does it take Wellbutrin XL/SR (bupropion) to work? 
- What's the difference between Wellbutrin XL and SR? 
- Does Auvelity work better than Wellbutrin? 

[View more FAQ...](#)

More about Wellbutrin (bupropion)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(756\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Support group](#)
- [Drug class: miscellaneous antidepressants](#)
- [Breastfeeding](#)

Patient resources

Other brands

[Zyban](#), [Forfivo XL](#), [Aplenzin](#), [Budeprion XL](#), ... +2 more

Professional resources

- [Wellbutrin prescribing information](#)
- [Wellbutrin Tablets \(FDA\)](#)
- [Bupropion Hydrobromide \(AHFS Monograph\)](#)

Other brands

[Zyban](#), [Forfivo XL](#), [Aplenzin](#)

Other formulations

- [Wellbutrin XL](#)
- [Wellbutrin SR](#)

Related treatment guides

- [Major Depressive Disorder](#)

- [Depression](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Wellbutrin only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 26.01.

DRUG STATUS

Availability

 Discontinued

Pregnancy & Lactation

 Risk data available

CSA Schedule*

N/A Not a controlled drug



Approval History

 FDA approved 1985




User Reviews & Ratings

6.8 / 10

[756 Reviews](#)

Images

[Wellbutrin 100 mg \(WELLBUTRIN 100\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.