

Home

Tofranil

Tofranil ⊲

Generic name: imipramine [*im-IP-ra-meen*] **Drug class:** Tricyclic antidepressants



Medically reviewed by Drugs.com on Aug 3, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Tofranil?

Tofranil is a tricyclic antidepressant used to treat symptoms of depression. This medicine is sometimes used to treat bedwetting in children ages 6 and older.

Tofranil may also be used for purposes not listed in this medication guide.

Tofranil side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Tofranil may cause serious side effects. Call your doctor at once if you have:

- easy bruising, unusual bleeding, purple or red spots under your skin;
- tunnel vision, eye pain or swelling, or seeing halos around lights;
- a light-headed feeling, like you might pass out;
- · new or worsening chest pain, pounding heartbeats or fluttering in your chest;
- sudden numbness or weakness, problems with vision, speech, or balance;
- · fever, sore throat;
- · confusion, hallucinations, unusual thoughts or behavior;
- · painful or difficult urination;
- seizure (convulsions); or
- jaundice (yellowing of the skin or eyes).

Common side effects of Tofranil may include:

- · increased blood pressure;
- tingly feeling, weakness, lack of coordination;
- dry mouth, nausea, vomiting; or
- breast swelling (in men or women).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Tofranil side effects (more detail)

Related/similar drugs

Naproxen

Naproxen is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

683 Reviews

View more

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-thecounter painkiller and fever ...

Reviews & ratings

6.1 / 10

13 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

1,404 Reviews

View more

Tramadol

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

Reviews & ratings

7.0 / 10

1,876 Reviews

View more

Meloxicam

Meloxicam is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.3 / 10

600 Reviews

View more

Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

Reviews & ratings

6.0 / 10

618 Reviews

View more

Sertraline (Zoloft) is an SSRI used to treat depression, anxiety, OCD, PTSD, and PMDD by increasing ...

Reviews & ratings

7.3 / 10

4,561 Reviews

View more

Trazodone

Trazodone is a tetracyclic antidepressant used to treat depression and anxiety disorders. Includes ...

Reviews & ratings

6.3 / 10

1,432 Reviews

View more

Aspirin

Aspirin is used to treat mild to moderate pain and to reduce fever or inflammation. Learn about ...

Reviews & ratings

7.8 / 10

51 Reviews

View more

Warnings

Some young people have thoughts about suicide when first taking an antidepressant. Stay alert to changes in your mood or symptoms. **Report any new or worsening symptoms to your doctor**.

Before taking this medicine

You should not use Tofranil if you are allergic to it, or if:

- you recently had a heart attack; or
- you are allergic to other antidepressants (amitriptyline, amoxapine, clomipramine, desipramine, doxepin, nortriptyline, protriptyline, or trTofranil).

Do not use Tofranil if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, tranylcypromine, and others.

Tell your doctor if you have used an "SSRI" antidepressant in the past 5 weeks, such as citalopram, escitalopram, fluoxetine (Prozac), fluoxamine, paroxetine, sertraline (Zoloft), trazodone, or vilazodone.

Tell your doctor if you have ever had:

- heart disease, stroke, or seizures;
- bipolar disorder (manic-depression), schizophrenia or other mental illness;
- kidney or liver disease;
- · overactive thyroid;
- · narrow-angle glaucoma; or
- problems with urination.

Some young people have thoughts about suicide when first taking an antidepressant. Stay alert to changes in your mood or symptoms. Report any new or worsening symptoms to your doctor.

It is not known whether Tofranil will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

You should not breastfeed while using Tofranil.

Do not give this medicine to a child without medical advice. To franil is not approved to treat depression in anyone younger than 18 years old. This medicine should not be used to treat bed-wetting in a child younger than 6 years old.

How should I take Tofranil?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Tell your doctor if you have a planned surgery.

Call your doctor if your symptoms do not improve, or if they get worse.

You should not stop using Tofranil suddenly. Follow your doctor's instructions about tapering your dose.

Store at room temperature away from moisture, heat, and light. Keep the bottle tightly closed when not in use.

Detailed Tofranil dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of Tofranil can be fatal.**

Overdose symptoms may include irregular heartbeats, feeling like you might pass out, seizures, or coma.

What should I avoid while taking imipramine?

Do not drink alcohol. Dangerous side effects or death can occur when alcohol is combined with Tofranil.

Avoid driving or hazardous activity until you know how Tofranil will affect you. Your reactions could be impaired.

Avoid exposure to sunlight or tanning beds. Tofranil can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

What other drugs will affect Tofranil?

Taking this medicine with other drugs that make you sleepy can worsen this effect. Ask your doctor before taking Tofranil with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

Many drugs can affect Tofranil. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all other medicines you use.

1 Tofranil drug interactions (more detail)

Does Tofranil interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Tofranil	
+	
Enter a drug name	Add

More about Tofranil (imipramine)

- · Check interactions
- Compare alternatives
- Reviews (16)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: tricyclic antidepressants

Breastfeeding

Patient resources

· Tofranil advanced reading

Other brands

Tofranil-PM

Professional resources

- Tofranil prescribing information
- Tofranil injection (FDA)
- Imipramine (AHFS Monograph)

Related treatment guides

- Enuresis
- Depression
- Pain
- Primary Nocturnal Enuresis

Further information


Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 14.01.

DRUG STATUS



User Reviews & Ratings

7.6 / 10

16 Reviews

Images

Tofranil 10 mg (M 10)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





