

Home

2. Doxorubicin

Doxorubicin ⊲₀

Generic name: doxorubicin [DOX-oh-ROO-bi-sin]

Brand names: Adriamycin, Adriamycin RDF, Rubex, Adriamycin PFS

Dosage forms: intravenous powder for injection (10 mg; 50 mg), intravenous solution (2 mg/mL)

Drug class: Antibiotics / antineoplastics



Medically reviewed by Drugs.com on Mar 14, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is doxorubicin?

Doxorubicin is used to treat different types of cancers that affect the breast, bladder, kidneys, ovaries, thyroid, stomach, lungs, bones, nerve tissues, joints, and soft tissues. Doxorubicin is also used to treat Hodgkin lymphoma, non-Hodgkin lymphoma, and certain types of leukemia.

Doxorubicin may also be used for purposes not listed in this medication guide.

Doxorubicin side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Some side effects may occur during the injection. Tell your caregiver right away if you feel dizzy, nauseated, light-headed, sweaty, or have a headache, chest tightness, back pain, trouble breathing, or swelling in your face.

Doxorubicin may cause serious side effects. Call your doctor at once if you have:

- pain, blisters, or skin sores where the injection was given;
- · missed menstrual periods;
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin;
- low white blood cell counts--fever, swollen gums, painful mouth sores, pain when swallowing, skin sores, cold or flu symptoms, cough, trouble breathing; or
- signs of heart problems--fast heartbeats, shortness of breath (even with mild exertion), feeling very weak or tired, swelling in your ankles or feet.

Doxorubicin may cause your urine to turn a reddish-orange color. This side effect is usually not harmful.

Common side effects of doxorubicin may include:

- nausea, vomiting; or
- hair loss.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Doxorubicin side effects (more detail)

Related/similar drugs

Abecma

Abecma is a CAR T cell therapy used to treat relapsed refractory multiple myeloma (RRMM); it is a ...

Reviews & ratings

Add a review

View more

Tecvayli

Tecvayli (teclistamab) is a treatment for relapsed or refractory multiple myeloma(cancer of the ...

Reviews & ratings

5 Reviews

View more

FEATURED

Carvykti

Carvykti (ciltacabtagene autoleucel) is used for the treatment of relapsed or refractory multiple ...

Reviews & ratings

Add a review

View more

Verzenio

Verzenio is used to treat certain HRpositive and HER2-negative breast cancers in adults. It is an ...

Reviews & ratings

8.0 / 10

View more

Trodelvy

Trodelvy is a targeted anticancer medication used to treat types of breast cancer and bladder ...

Reviews & ratings

4 Reviews

View more

Opdivo

Opdivo is used to treat many cancers, such as melanoma, non-small cell lung cancer, malignant ...

Reviews & ratings

5.8 / 10

93 Reviews

View more

Darzalex

Darzalex (daratumumab) is a target therapy used to treat multiple myeloma. Includes DARZALEX uses

Reviews & ratings

6 Reviews

View more

Talvey

Talvey is used to treat relapsed or refractory multiple myeloma. Includes Talvey side effects ...

Reviews & ratings

1 Reviews

View more

Keytruda is used to treat multiple types of cancer such as melanoma, nonsmall cell lung cancer ...

Reviews & ratings

4.9 / 10

252 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Warnings

You should not use doxorubicin if you have severe liver disease, severe heart problems, very low blood cell counts caused by prior chemotherapy, or if you recently had a heart attack.

Doxorubicin can weaken your immune system. Your blood may need to be tested often. Tell your doctor if you have unusual bruising or bleeding, or signs of infection (fever, chills, body aches).

Doxorubicin may cause dangerous effects on your heart. Call your doctor at once if you feel very weak or tired, or have fast heartbeats, shortness of breath (even with mild exertion), or swelling in your ankles or feet.

Before taking this medicine

You should not use this medicine if you are allergic to doxorubicin or similar medications (doxorubicin, daunorubicin, epirubicin, idarubicin, mitoxantrone), or if you have:

- · very low blood cell counts caused by chemotherapy you received in the past;
- · severe liver disease;
- · severe heart problems; or
- if you have recently had a heart attack.

Tell your doctor if you have ever had:

- severe liver disease;
- · heart disease or heart failure; or

· radiation treatment to your chest.

Tell your doctor about all other cancer medicines or radiation treatments you have received in the past.

Using doxorubicin may increase your risk of developing a bone marrow disease or other types of leukemia later in life. Ask your doctor about your specific risk.

You may need to have a negative pregnancy test before starting this treatment.

Both men and women using this medicine should use effective birth control to prevent pregnancy. Doxorubicin can harm an unborn baby or cause birth defects if the mother or father is using this medicine.

Keep using birth control for at least 6 months after your last dose. Tell your doctor right away if a pregnancy occurs while either the mother or the father is using doxorubicin.

This medicine may affect fertility (ability to have children) in both men and women. However, it is important to use birth control to prevent pregnancy because doxorubicin can harm an unborn baby.

Doxorubicin may also cause early menopause, depending on your age when you receive this medicine. Ask your doctor about this risk.

You should not breastfeed while you are using doxorubicin.

How is doxorubicin given?

Doxorubicin is given as an infusion into a vein. A healthcare provider will give you this injection.

Tell your caregivers if you feel any burning, pain, or swelling around the IV needle when doxorubicin is injected.

Doxorubicin is sometimes given together with other cancer medications. You may be given other medications to prevent nausea, vomiting, or infections.

If any of this medication accidentally gets on your skin, wash it thoroughly with soap and warm water.

Doxorubicin can lower your blood cell counts. **Your blood will need to be tested often.** Your cancer treatments may be delayed based on the results.

Detailed Doxorubicin dosage information

What happens if I miss a dose?

Call your doctor for instructions if you miss an appointment for your doxorubicin injection.

What happens if I overdose?

Since this medication is given by a healthcare professional in a medical setting, an overdose is unlikely to occur.

What should I avoid while taking doxorubicin?

Avoid being near people who are sick or have infections. Tell your doctor at once if you develop signs of infection.

Avoid activities that may increase your risk of bleeding or injury. Use extra care to prevent bleeding while shaving or brushing your teeth.

What other drugs will affect doxorubicin?

Many drugs can affect doxorubicin. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all your current medicines and any medicine you start or stop using.

1 Doxorubicin drug interactions (more detail)

Does doxorubicin interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

What is the chemo drug called the Red Devil?	~
What is the strongest chemotherapy drug for breast cancer?	~
How is Lartruvo administered?	~

More FAQ

- What is the lifetime or cumulative dose for Adriamycin?
- How does AC (Adriamycin and Cytoxan [cyclophosphamide]) work for Breast Cancer?
- What are the long-term side effects of Adriamycin?
- Does Adriamycin cause heart failure?

View more FAQ...

More about doxorubicin

- · Check interactions
- Compare alternatives
- · Pricing & coupons

- Reviews (8)
- Side effects
- Dosage information
- During pregnancy
- Drug class: antibiotics/antineoplastics
- Breastfeeding
- En español

Patient resources

- Doxorubicin advanced reading
- Doxorubicin (Conventional)

Other brands

Adriamycin

Professional resources

- DOXOrubicin monograph
- Doxorubicin (FDA)
- Doxorubicin Liposomal (FDA)

Other brands

Adriamycin

Related treatment guides

- Bladder Cancer
- Breast Cancer
- Acute Lymphoblastic Leukemia
- Acute Myeloblastic Leukemia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.02.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ড Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

5.4 / 10

8 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









