

Home

2. Levetiracetam

Levetiracetam (oral/injection) ⊲ ¬

Generic name: levetiracetam (oral/injection) [LEE-ve-tye-RA-se-tam]

Brand names: Elepsia XR, Keppra, Keppra XR, Roweepra, Spritam, Roweepra XR

Dosage forms: intravenous solution (100 mg/mL; 1000 mg/100 mL-NaCl 0.75%; 1500 mg/100 mL-NaCl 0.54%; 250 mg/50 mL-NaCl 0.82%;

500 mg/100 mL-NaCl 0.82%), ... show all 5 dosage forms

Drug class: Pyrrolidine anticonvulsants



Medically reviewed by Drugs.com on Apr 29, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is levetiracetam?

Levetiracetam is used alone or together with other medicines to treat certain types of seizures in people with epilepsy, including partial onset seizures, myoclonic seizures, and tonic-clonic seizures.

Levetiracetam is used to treat myoclonic seizures in people who are at least 12 years old and also used to treat primary generalized tonic-clonic seizures in people with certain types of epilepsy who are at least 6 years old.

When used for partial onset seizures:

- the levetiracetam extended-release tablet is for adults and children at least 12 years old.
- the **levetiracetam tablet for oral suspension** is for children at least 4 years old who weigh at least 44 pounds (20 kilograms).
- other levetiracetam brand forms may be used in adults and children at least 1 month old.

Levetiracetam **injection** is usually given only if you are unable to take the medicine by mouth. Some brand forms of levetiracetam **injection** are for use only in people who are at least 16 years old.

Levetiracetam may also be used for purposes not listed in this medication guide.

Levetiracetam side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes.

Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself.

Levetiracetam may cause serious side effects. Call your doctor at once if you have:

- confusion, hallucinations, extreme drowsiness, feeling very weak or tired;
- loss of balance or coordination, problems with walking or movement;
- · a skin rash, no matter how mild; or
- low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath.

Common side effects of levetiracetam may include:

- dizziness, drowsiness, tiredness, weakness;
- · feeling aggressive or irritable;
- loss of appetite, nausea;
- · stuffy nose, sore throat, runny nose; or
- infections, including the flu.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

<u>Levetiracetam side effects</u> (more detail)

Related/similar drugs

Lamictal

Lamictal is an anti-epileptic medication used treat seizures in adults and children over 2 years ...

Reviews & ratings

8.0 / 10

792 Reviews

View more

Depakote

Depakote is used to treat various types of seizure disorders. Learn about side effects ...

Reviews & ratings

6.3 / 10

329 Reviews

View more

FEATURED

Caplyta

Caplyta (lumateperone) may be used to treat schizophrenia or depression associated with bipolar ...

Reviews & ratings

6.6 / 10

184 Reviews

View more

Lyrica

Lyrica is used to control seizures, treat nerve pain and fibromyalgia. Learn about side effects ...

Reviews & ratings

6.4 / 10

925 Reviews

View more

Keppra

Keppra is an anti-epileptic drug used to treat partial onset seizures in people with epilepsy ...

Reviews & ratings

5.7 / 10

307 Reviews

View more

Diazepam

Diazepam is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

• • •

Reviews & ratings

8.4 / 10

884 Reviews

View more

* 10 TF 11101 0

Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

Reviews & ratings

7.9 / 10

1,054 Reviews

View more

Pregabalin

Pregabalin may be used to treat certain types of pain and used in combination with other ...

Reviews & ratings

6.8 / 10

1,668 Reviews

View more

Clonazepam

Clonazepam is used to treat seizures and panic disorder. Learn about side effects, interactions and ...

Reviews & ratings

8.2 / 10

2,176 Reviews

View more

Warnings

Some people have thoughts about suicide while taking seizure medicine. Tell your doctor right away if you have any sudden changes in mood or behavior, or thoughts about suicide.

Seizures may increase if you stop using levetiracetam suddenly. Ask your doctor before stopping the medicine.

Before taking this medicine

You should not use levetiracetam if you are allergic to it.

Tell your doctor if you have or have ever had:

- · depression, a mood disorder, or suicidal thoughts or actions
- high blood pressure;
- mental illness or psychosis; or
- kidney disease (or if you are on dialysis).

Do not give levetiracetam to a child without medical advice. Certain brand forms of this medicine are for use only in children of certain ages.

Some people have thoughts about suicide while taking seizure medicine. Stay alert to changes in your mood or symptoms. Your family or caregivers should also watch for sudden changes in your behavior.

Do not start or stop seizure medication during pregnancy without your doctor's advice. It is not known if levetiracetam will harm an unborn baby, but having a seizure during pregnancy could harm both mother and baby. Ask your doctor about this risk. Tell your doctor if you become pregnant or plan to become pregnant.

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of levetiracetam on the baby.

Ask a doctor if it is safe to breastfeed while using this medicine.

How should I use levetiracetam?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You may take levetiracetam oral with or without food.

Levetiracetam injection is given into a vein by a healthcare provider.

Your dose needs may change if you switch to a different brand, strength, or form of this medicine. **Avoid medication** errors by using exactly as directed on the label, or as prescribed by your doctor.

Doses are based on weight in children and teenagers. Your child's dose may change if the child gains or loses weight.

Measure liquid medicine with the supplied measuring device (not a kitchen spoon).

Swallow the **tablet** whole and do not crush or chew it. Ask your doctor or pharmacist if you are unable to swallow a whole tablet. **Do not use a tablet that is damaged in any way.**

Read and carefully follow any Instructions for Use provided to you on how to prepare and take levetiracetam tablet for oral suspension. Ask your doctor or pharmacist if you do not understand these instructions.

Part of the extended-release tablet may appear in your stool, but this will not make the medicine less effective.

Your kidney function and blood pressure may need to be checked often. **Keep all follow-up visits with your healthcare provider.**

Seizures may increase if you stop using levetiracetam suddenly. Ask your doctor before stopping the medicine. Follow your doctor's instructions about tapering your dose.

Store at room temperature away from moisture, heat, and light.

Detailed Levetiracetam dosage information

What happens if I miss a dose?

Take levetiracetam **oral** as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

In a medical setting you are not likely to miss a dose of levetiracetam injection.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include extreme drowsiness, agitation, aggression, shallow breathing, weakness, or fainting.

What should I avoid while using levetiracetam?

Avoid driving or hazardous activity until you know how levetiracetam will affect you. Dizziness or loss of balance or coordination can cause falls, accidents, severe injuries, or may impair your reactions.

What other drugs will affect levetiracetam?

Other drugs may affect levetiracetam, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all other medicines you use.

Levetiracetam drug interactions (more detail)

Does levetiracetam interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

levetiracetam	
+	
Enter a drug name	Add

More about levetiracetam

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (429)

- Drug images
- Latest FDA alerts (6)
- Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: pyrrolidine anticonvulsants
- Breastfeeding
- En español

Patient resources

- Levetiracetam Extended-Release Tablets patient information
- Levetiracetam Injection
- Levetiracetam Tablets
- Levetiracetam Oral Solution

Other brands

Keppra, Keppra XR, Spritam, Elepsia XR, ... +2 more

Professional resources

- · Levetiracetam monograph
- Levetiracetam (FDA)
- Levetiracetam Extended Release Tablets (FDA)
- Levetiracetam Injection (FDA)
- Levetiracetam Injection, Concentrate (FDA)
- Levetiracetam Solution (FDA)

Other brands

Keppra, Keppra XR, Spritam, Elepsia XR, ... +2 more

Related treatment guides

- Seizures
- Bipolar Disorder
- Epilepsy
- Neuralgia

- Hyperekplexia
- New Daily Persistent Headache

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 15.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

প্ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

User Reviews & Ratings

5.9 / 10

429 Reviews

Related News

Alert: Dr. Reddy's Issues a Nationwide Recall of Levetiracetam in 0.75% Sodium Chloride Injection 1,000 mg/100 mL, in the U.S., Due to Mislabeling of Infusion Bag

Images

Levetiracetam 500 mg (H 88)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





