

Home

Lacosamide

Lacosamide (oral/injection) ⊲ ¬

Generic name: lacosamide (oral/injection) [la-KOE-sa-mide]

Brand names: Motpoly XR, Vimpat

Dosage forms: intravenous solution (200 mg/20 mL), oral capsule, extended release (100 mg; 150 mg; 200 mg), ... show all 4 dosage

forms

Drug class: Miscellaneous anticonvulsants



Medically reviewed by Drugs.com on Feb 4, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is lacosamide?

Lacosamide is used to treat partial onset seizures in adults and children who are at least 1 month old.

Lacosamide is also used to treat **primary generalized tonic-clonic seizures** in adults and children who are at least 4 years old.

Lacosamide may also be used for purposes not listed in this medication guide.

Lacosamide side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself.

Lacosamide may cause serious side effects. Call your doctor at once if you have:

- a light-headed feeling, like you might pass out;
- severe dizziness;
- problems with balance or muscle movement;
- · chest pain, shortness of breath;

- · fast or pounding heartbeats;
- · dark urine; or
- · very slow heartbeats.

Common side effects of lacosamide may include:

- headache, dizziness;
- · drowsiness:
- · double vision; or
- nausea.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1 800 FDA 1088.

A

Lacosamide side effects (more detail)

Related/similar drugs

Depakote

Depakote is used to treat various types of seizure disorders. Learn about side effects ...

Reviews & ratings

6.3 / 10

329 Reviews

View more

Lyrica

Lyrica is used to control seizures, treat nerve pain and fibromyalgia. Learn about side effects ...

Reviews & ratings

6.4 / 10

925 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

View more

Keppra

Keppra is an anti-epileptic drug used to treat partial onset seizures in people with epilepsy ...

Reviews & ratings

5.7 / 10

307 Reviews

View more

Diazepam

Diazepam is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn ...

Reviews & ratings

8.4 / 10

884 Reviews

View more

Lamictal

Lamictal is an anti-epileptic medication used treat seizures in adults and children over 2 years ...

Reviews & ratings

8.0 / 10

792 Reviews

View more

Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

Reviews & ratings

7.9 / 10

View more

Pregabalin

Pregabalin may be used to treat certain types of pain and used in combination with other ...

Reviews & ratings

6.8 / 10

1,668 Reviews

View more

Warnings

Do not stop using lacosamide suddenly or you may have increased seizures.

Some people have thoughts about suicide while taking lacosamide. **Tell your doctor right away if you have any sudden changes in mood or behavior, or thoughts about suicide.**

You may feel faint while using lacosamide. Call your doctor right away if you have fast or pounding heartbeats, shortness of breath, and sudden dizziness.

Before taking this medicine

You should not use lacosamide if you are allergic to it.

Tell your doctor if you have ever had:

- · heart problems;
- depression, a mood disorder, suicidal thoughts or actions;
- · drug or alcohol addiction;
- nerve problems caused by diabetes;
- · kidney disease; or
- liver disease.

Lacosamide oral solution may contain phenylalanine. Tell your doctor if you have phenylketonuria (PKU).

Some people have thoughts about suicide while using lacosamide. Your doctor will need to check your progress at regular visits. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Do not start or stop using seizure medication during pregnancy without your doctor's advice. Having a seizure during pregnancy could harm both mother and baby. Tell your doctor right away if you become pregnant.

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of lacosamide on the baby.

Ask a doctor before using this medicine if you are breastfeeding. Tell your doctor if you notice severe drowsiness in the nursing baby.

How should I use lacosamide?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Lacosamide **oral** is taken by mouth. Lacosamide **injection** is given as an infusion into a vein. A healthcare provider will give you this injection if you are unable to take the medicine by mouth.

Doses are based on weight in children. Your child's dose may change if the child gains or loses weight.

Lacosamide may be habit-forming. **Misuse can cause addiction, overdose, or death.** Keep the medication in a place where others cannot get to it. Selling or giving away lacosamide is against the law.

Lacosamide tablets or oral solution can be taken with or without food.

Swallow the tablet/capsule whole and do not crush, chew, or break it.

Measure the **oral solution** with the supplied measuring device (not a kitchen spoon).

Call your doctor if your seizures get worse or you have them more often while using lacosamide.

Seizures may increase if you stop using lacosamide suddenly. Ask your doctor before stopping the medicine.

Wear a medical alert tag or carry an ID card to let others know you use lacosamide.

Store at room temperature away from moisture and heat. Do not freeze the liquid form of lacosamide. Throw away any **oral solution** not used within 7 weeks after opening the bottle.

Keep track of your medicine. You should be aware if anyone is using it improperly or without a prescription.

Detailed Lacosamide dosage information

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while using lacosamide?

Avoid driving or hazardous activity until you know how lacosamide will affect you. Your reactions could be impaired.

Drinking alcohol with this medicine can cause side effects such as drowsiness and dizziness.

What other drugs will affect lacosamide?

Tell your doctor about all your other medicines, especially:

• heart or blood pressure medication.

This list is not complete. Other drugs may affect lacosamide, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Lacosamide drug interactions (more detail)

Does lacosamide interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

lacosamide	
+	
Enter a drug name	Add

More about lacosamide

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (178)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: miscellaneous anticonvulsants
- Breastfeeding
- En español

Patient resources

- Lacosamide (Intravenous) advanced reading
- Lacosamide (Oral) (Advanced Reading)
- Lacosamide Tablets

- Lacosamide Injection
- Lacosamide Oral Solution

Other brands

Vimpat, Motpoly XR

Professional resources

- · Lacosamide monograph
- Lacosamide (FDA)
- Lacosamide Injection (FDA)
- Lacosamide Oral Solution (FDA)

Other brands

Vimpat, Motpoly XR

Related treatment guides

- Epilepsy
- Seizures

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 13.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্য Risk data available

CSA Schedule*

5 Some potential for abuse

Approval History

The Drug history at FDA

User Reviews & Ratings

6.8 / 10

178 Reviews

Images

Lacosamide 100 mg (SG 193)





Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





