

Home

2. Dronabinol

Dronabinol

Generic name: dronabinol [*droe-NAH-bih-nol*]

Brand names: Marinol, Syndros

Dosage forms: oral capsule (10 mg; 2.5 mg; 5 mg), oral solution (5 mg/mL)

Drug class: Miscellaneous antiemetics



Medically reviewed by Drugs.com on Jan 13, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is dronabinol?

Dronabinol is a man-made form of tetrahydrocannabinol (THC), a psychoactive substance found in cannabis (also known as marijuana).

Dronabinol is used to treat loss of appetite that causes weight loss in people with AIDS.

Dronabinol is also used to treat severe nausea and vomiting caused by cancer chemotherapy. Dronabinol is usually given after medicines to control nausea and vomiting have been tried without success.

Dronabinol may also be used for purposes not listed in this medication guide.

Dronabinol side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, skin rash or burning; mouth sores; warmth, redness, or tingly feeling; difficult breathing; swelling of your face, lips, tongue, or throat.

Dronabinol may cause serious side effects. Call your doctor at once if you have:

- a seizure;
- · fast or pounding heartbeats;
- a light-headed feeling, like you might pass out;
- confusion, trouble sleeping, problems with memory or concentration;
- unusual changes in mood or behavior;
- restlessness, feeling nervous or irritable;
- slurred speech, drowsiness;
- · severe or ongoing nausea, vomiting, or stomach pain; or

• increased blood pressure--severe headache, blurred vision, pounding in your neck or ears.

Common side effects of dronabinol may include:

- · feeling "high";
- · dizziness, drowsiness, thinking problems;
- · unusual thoughts or fears;
- · feelings of extreme happiness; or
- nausea, vomiting, stomach pain.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i Dronabinol side effects (more detail)

Related/similar drugs

Reglan

Reglan (metoclopramide) is used to treat heartburn caused by gastric reflux. Includes Reglan side ...

Reviews & ratings

3.7 / 10

352 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Metoclopramide

Metoclopramide is used short-term to treat heartburn caused by gastroesophageal reflux. Learn about

Reviews & ratings

00/40

View more

Ativan

Ativan is a benzodiazepine used to treat anxiety disorders or anxiety associated with depression ...

Reviews & ratings

8.0 / 10

479 Reviews

View more

Diphenhydramine

Diphenhydramine is an antihistamine used to treat sneezing, runny nose, itching, hives and other ...

Reviews & ratings

5.5 / 10

541 Reviews

View more

Meclizine

Meclizine is used to treat or prevent nausea, vomiting, and dizziness caused by motion sickness ...

Reviews & ratings

7.4 / 10

234 Reviews

View more

Promethazine

Promethazine is a phenothiazine and antihistamine used to treat allergies, motion sickness, nausea ...

Reviews & ratings

7.2 / 10

388 Reviews

View more

Zofran

Zofran (ondansetron) is used to prevent nausea and vomiting caused by surgery or cancer medicines ...

Reviews & ratings

8.0 / 10

270 Reviews

View more

Olanzapine

Olanzapine (brand name Zyprexa) is an atypical antipsychotic that may be used to treat adults and ...

Reviews & ratings

5.9 / 10

1,091 Reviews

View more

Warnings

Dronabinol may cause new or worsening psychosis (unusual thoughts or behavior), especially if you have ever had depression or mental illness.

You should not use dronabinol **capsules** if you are allergic to sesame oil. You should not use dronabinol **oral solution** if you have had an allergic reaction to alcohol or if you also use disulfiram (Antabuse) or metronidazole (Flagyl).

Dronabinol can raise or lower blood pressure, especially in older adults or in people with heart problems.

Call your doctor at once if you have new or worsening mood symptoms, changes in behavior, headaches, vision problems, rapid heartbeats, or severe dizziness.

Before taking this medicine

You should not use dronabinol if you are allergic to it.

You should not take dronabinol capsules if you are allergic to sesame oil.

Dronabinol oral solution (liquid medicine) contains alcohol. You should not take dronabinol liquid if you have ever had an allergic reaction to alcohol or if you also use disulfiram (Antabuse) or metronidazole (Flagyl). **Do not take dronabinol liquid within 14 days after or 7 days before using disulfiram or metronidazole.**

Tell your doctor if you have ever had:

- · an allergy to any medication;
- epilepsy or other seizure disorder;
- heart problems, high or low blood pressure, fainting spells, fast heartbeats;
- · alcoholism or drug addiction; or
- depression, mental illness, or psychosis.

It is not known whether dronabinol will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

HIV can be passed to your baby if you are not properly treated during pregnancy. **If you have HIV or AIDS**, take all of your medicines as directed to control your infection.

Women with HIV or AIDS should not breastfeed a baby. Even if your baby is born without HIV, the virus may be passed to the baby in your breast milk. However, if you are being treated for nausea and vomiting caused by cancer chemotherapy and you are breastfeeding, your doctor should check the weight of your baby often.

Dronabinol is not approved for use by anyone younger than 18 years old.

How should I take dronabinol?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed. **Tell your doctor if you feel an increased urge to use more of this medicine**.

Dronabinol may be habit-forming. **Misuse can cause addiction, overdose, or death.** Keep the medication in a place where others cannot get to it. Selling or giving away this medicine is against the law.

To stimulate appetite in people with AIDS, dronabinol is usually taken 1 hour before lunch and 1 hour before dinner.

To prevent nausea and vomiting caused by chemotherapy:

- Take your **first dose on an empty stomach**, at least 30 minutes before eating.
- After your first dose, you may take dronabinol with or without food but take it at the same time in relation to your mealtimes for each treatment for chemotherapy.
- Dronabinol is usually given 1 to 3 hours before your chemotherapy treatment, and then every 2 to 4 hours (up to 6 doses per day).

Older adults may need to take this medicine less often. Follow your doctor's dosing instructions very carefully.

Swallow the capsule whole and do not crush, chew, break, or open it.

Measure **liquid medicine** carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

Take the liquid medicine with a full glass of water.

You may need to follow a special diet while using dronabinol. Follow all instructions of your doctor or dietitian. Learn

about the foods you should eat to help boost your food intake.

Call your doctor if your symptoms do not improve, or if they get worse while using dronabinol.

Do not stop using dronabinol suddenly after long-term use, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine.

Store the capsules or unopened liquid medicine in a tightly-closed container in the refrigerator. Do not freeze.

You may also store the **capsules** at cool room temperature, away from moisture and heat. **After opening the liquid medicine**, store it at room temperature. Throw away any unused liquid 42 days after opening the bottle.

Detailed Dronabinol dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include changes in mood, memory problems, little or no urinating, constipation, loss of energy, problems with speech or coordination, or feeling light-headed.

What should I avoid while taking dronabinol?

Do not use marijuana while taking dronabinol.

Avoid driving or operating machinery until you know how dronabinol will affect you. Dizziness or severe drowsiness can cause falls or other accidents.

Do not drink alcohol while taking dronabinol.

What other drugs will affect dronabinol?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Using dronabinol with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Other drugs may affect dronabinol, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

Dronabinol drug interactions (more detail)

Does dronabinol interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

dronabinol	
+	
Enter a drug name	Add

More about dronabinol

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (61)
- Drug images
- Latest FDA alerts (1)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: miscellaneous antiemetics
- Breastfeeding
- En español

Patient resources

- Dronabinol advanced reading
- Dronabinol Capsules

Other brands

Marinol, Syndros

Professional resources

- Dronabinol monograph
- Dronabinol (FDA)

Other brands

Marinol, Syndros

Related treatment guides

• Post Traumatic Stress Disorder

- Unintentional Weight Loss (Underweight)
- Anorexia
- AIDS Related Wasting
- Nausea/Vomiting
- Nausea/Vomiting, Chemotherapy Induced

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 12.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্% Risk data available

CSA Schedule*

M Multiple: 2,3

Approval History

□ Drug history at FDA

User Reviews & Ratings

8.3 / 10

61 Reviews

Images

Dronabinol 2.5 mg (RP 867)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





