

#### Home

2. Naphazoline

# Naphazoline ophthalmic

Generic name: naphazoline ophthalmic [ na-FAZ-oh-leen-off-THAL-mik ]

Brand names: Clear Eyes, Clear Eyes + Redness Relief, Redness Relief Eye Drops, AK-Con, Albalon, ... show all 18 brands

Dosage forms: ophthalmic gel forming solution (0.012%), ophthalmic solution (0.012% with glycerin)

Drug class: Ophthalmic antihistamines and decongestants



Medically reviewed by Drugs.com on Dec 21, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking What to avoid Interactions

# What is naphazoline ophthalmic?

Naphazoline is a vasoconstrictor. It works by narrowing swollen blood vessels in the eyes to reduce eye redness.

Naphazoline ophthalmic (for the eye) is for temporary relief of minor eye redness or discomfort caused by minor irritants.

Naphazoline ophthalmic may also be used for purposes not listed in this medication guide.

# Naphazoline ophthalmic side effects

Get emergency medical help if you have any of these **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Naphazoline ophthalmic may cause serious side effects. Stop using naphazoline ophthalmic and call your doctor at once if you have:

- · ongoing or worsening eye redness;
- · eye pain;
- · changes in your vision;
- · chest pain, fast or uneven heart rate; or
- severe headache, buzzing in your ears, anxiety, confusion, or feeling short of breath.

#### Common side effects of naphazoline ophthalmic may include:

- mild burning or stinging of the eye;
- · blurred vision, watery eyes; or
- mild headache, dizziness, nervousness.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**a** 

Naphazoline ophthalmic side effects (more detail)

# Related/similar drugs

### Alphagan P

Alphagan P is used to treat glaucoma or ocular hypertension (high pressure inside the eye). Learn ...

#### **Reviews & ratings**

5.0 / 10

8 Reviews

View more

#### FEATURED

#### Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

### Reviews & ratings

5.3 / 10

685 Reviews

View more

### Lumify

Lumify Eye Drops and Lumify Preservative Free are used to reduce eye redness.lncludes Lumify side ...

#### **Reviews & ratings**

3.2 / 10

174 Reviews

View more

# Warnings

Naphazoline ophthalmic is for temporary relief of minor eye redness or discomfort caused by minor irritants.

You should not use naphazoline ophthalmic if you have narrow-angle glaucoma.

Stop using naphazoline ophthalmic and call your doctor at once if you have ongoing or worsening eye redness, eye pain, vision changes, severe dizziness, or headache, buzzing in your ears, or feeling short of breath.

# Before taking this medicine

You should not use naphazoline ophthalmic if you are allergic to it, or if you have narrow-angle glaucoma.

Ask a doctor or pharmacist if it is safe for you to take this medicine if you have other medical conditions, especially:

- · heart disease, high blood pressure;
- · diabetes;
- a thyroid disorder; or
- an eye injury or infection.

FDA pregnancy category C. It is not known whether naphazoline ophthalmic will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medication.

It is not known whether naphazoline ophthalmic passes into breast milk or if it could harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

# How should I use naphazoline ophthalmic?

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended.

Using the medication too long or too often may worsen your symptoms and cause damage to the blood vessels in your eyes.

Wash your hands before using the eye drops.

To apply the eye drops:

- Tilt your head back slightly and pull down your lower eyelid to create a small pocket. Hold the dropper above the eye with the tip down. Look up and away from the dropper and squeeze out a drop.
- Close your eye and gently press your finger to the inside corner of the eye for about 1 minute, to keep the liquid from draining into your tear duct.
- Use only the number of drops recommended.

Do not touch the tip of the eye dropper or place it directly on your eye. A contaminated dropper can infect your eye, which could lead to serious vision problems.

Do not use the eye drops if the liquid has changed colors or has particles in it.

Store at room temperature away from moisture and heat. Do not freeze. Keep the bottle tightly closed when not in use.

# What happens if I miss a dose?

Since naphazoline ophthalmic is used when needed, you may not be on a dosing schedule. If you are on a schedule,

use the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** use extra medicine to make up the missed dose.

# What happens if I overdose?

An overdose of naphazoline ophthalmic is not expected to be dangerous. Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

Keep naphazoline ophthalmic out of the reach of children. Certain eye medications can cause serious medical problems in a young child who accidentally sucks on or swallows medicine from the eye dropper.

# What should I avoid while using naphazoline ophthalmic?

Do not use this medication while wearing contact lenses. Naphazoline ophthalmic may contain a preservative that can discolor soft contact lenses. Wait at least 15 minutes after using this medicine before putting in your contact lenses.

# What other drugs will affect naphazoline ophthalmic?

Ask a doctor or pharmacist if it is safe for you to use naphazoline ophthalmic if you are also using any of the following drugs:

- an antidepressant--amitriptyline, clomipramine, desipramine, desvenlafaxine, doxepin, duloxetine, imipramine, maprotiline, milnacipran, nortriptyline, venlafaxine;
- ergot medicine--ergotamine, dihydroergotamine, ergonovine, methylergonovine; or
- an MAO inhibitor--furazolidone, isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, tranylcypromine.

This list is not complete and other drugs may interact with naphazoline ophthalmic. Tell your doctor about all medications you use. This includes prescription, over-the-counter, vitamin, and herbal products. Do not start a new medication without telling your doctor.

Naphazoline ophthalmic drug interactions (more detail)

# Does naphazoline ophthalmic interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

naphazoline ophthalmic	
+	
Enter a drug name	Add

# More about naphazoline ophthalmic

- · Check interactions
- Compare alternatives

- Pricing & coupons
- Reviews (3)
- · Side effects
- · Drug class: ophthalmic antihistamines and decongestants
- En español

### **Patient resources**

· Naphazoline advanced reading

#### Other brands

Naphcon, Clear Eyes Redness Relief, Albalon, AK-Con, ... +3 more

### **Professional resources**

- Naphazoline monograph
- Naphazoline (FDA)

#### Other brands

Nazil Ofteno

### Related treatment guides

• Eye Redness

### **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.02.

#### **DRUG STATUS**


Availability
Rx ont
OTC

CSA Schedule\*

N/A

Not a controlled drug

Approval History



3 Reviews

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com
Advertising policy
Attribution & citations

#### Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

#### Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.