

[Home](#)

2. Remeron

Remeron

Generic name: [mirtazapine](#) [*mir-TAZ-a-peen*]**Brand names:** Remeron, [Remeron SolTab](#)**Drug class:** [Tetracyclic antidepressants](#)Medically reviewed by [Kaci Durbin, MD](#). Last updated on Aug 23, 2023.[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is Remeron?

Remeron is an antidepressant. The way mirtazapine works is still not fully understood. It is thought to positively affect communication between nerve cells in the central nervous system and/or restore chemical balance in the brain.

Remeron is prescription medicine used to treat a certain type of depression called [Major Depressive Disorder \(MDD\)](#) in adults.

It is not known if Remeron is safe and effective for use in treating MDD in children.

Warnings

You should not take Remeron if you are also taking tryptophan (sometimes called L-tryptophan).

Do not use this medicine if you have used an MAO inhibitor in the past 14 days, such as [isocarboxazid](#), [linezolid](#), [methylene blue](#) injection, [phenelzine](#), [tranylcypromine](#) and others.

Some people have thoughts about suicide when first taking an antidepressant. Your doctor will need to check your progress at regular visits while you are using Remeron. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Do not give this medicine to anyone younger than 18 years old without the advice of a doctor. Mirtazapine is not approved for use in children.

It may take up to several weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve after 4 weeks of treatment.

Drinking alcohol can increase certain side effects of mirtazapine.

Remeron may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

Do not stop taking mirtazapine abruptly. Talk to your doctor before stopping mirtazapine.

Before taking this medicine

You should not take Remeron if you are allergic to mirtazapine or any of the ingredients in Remeron or Remeron SolTab.

Do not use Remeron if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, tranylcypromine, and others.

To make sure Remeron is safe for you, tell your doctor if you have ever had:

- low white blood cell counts;
- [glaucoma](#);
- liver or kidney disease;
- low sodium levels in your blood;
- a seizure;
- [depression](#), suicidal thoughts or actions;
- [bipolar disorder](#) (manic depression) in you or a family member;
- heart problems or stroke;
- [high cholesterol](#) or triglycerides;
- [long QT syndrome](#) (in you or a family member); or
- [low blood pressure](#).

Be sure your doctor knows if you also take stimulant medicine, opioid medicine, [herbal products](#), or medicine for depression, mental illness, [Parkinson's disease](#), [migraine](#) headaches, serious infections, or prevention of [nausea and vomiting](#). These medicines may interact with mirtazapine and cause a serious condition called serotonin syndrome.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor should check your progress at regular visits. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Tell your doctor if you are pregnant or breastfeeding.

The orally disintegrating tablet may contain phenylalanine and could be harmful if you have [phenylketonuria](#) (PKU).

Remeron is not approved for use by anyone younger than 18 years old.

 [Remeron pregnancy and breastfeeding warnings](#) (more detail)

How should I take Remeron?

Take Remeron exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Take the medicine at the same time each day, usually at bedtime.

Take the **regular Remeron tablet** with water.

You may take this medicine with or without food.

Remove a Remeron SolTab orally disintegrating tablet from the package only when you are ready to take the medicine. Place the tablet on your tongue and allow it to dissolve, without chewing. Swallow several times as the tablet dissolves.

It may take a few weeks for your symptoms to improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve after 4 weeks of treatment.

Do not stop using Remeron suddenly, or you could have unpleasant symptoms (such as [dizziness](#), [vomiting](#), [agitation](#), sweating, confusion, numbness, tingling, or electric shock feelings). Ask your doctor how to safely stop using this medicine.

Store at room temperature away from moisture, heat, and light.

 [Remeron patient tips](#) (more detail)

Dosing information

Usual Adult Dose for Depression:

Initial dose: 15 mg orally once a day at bedtime

Maintenance dose: 15 to 45 mg orally once a day

Maximum dose: 45 mg/day

Comments:

-May increase the dose every 1 to 2 weeks to a maximum 45 mg/day according to patient response.

-Patients should be periodically reassessed to determine the need for continued use of this drug.

Use: Treatment of major depressive disorder

 [Detailed Remeron dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include confusion, memory problems, drowsiness, and [fast heart rate](#).

What to avoid

Drinking alcohol with this medicine can cause side effects.

Avoid driving or hazardous activity until you know how this medicine will affect you. Your reactions could be impaired.

Remeron side effects

Get emergency medical help if you have **signs of an allergic reaction to Remeron** ([hives](#), joint pain, fever, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, [sore throat](#), burning eyes, skin pain, red or purple [skin rash](#) with blistering and peeling).

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, [anxiety](#), [panic attacks](#), trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Call your doctor at once if you have:

- racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual;
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights;
- a light-headed feeling, like you might pass out;
- severe rash, blisters, or swelling on the palms of your hands or the soles of your feet;
- a seizure;
- **low white blood cell counts** - fever, chills, sore throat, cough, sores in your mouth or nose, flu-like symptoms, trouble breathing; or
- **low sodium level** - [headache](#), confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, [nausea](#), vomiting, or [diarrhea](#).

Common Remeron side effects include:

- drowsiness, dizziness;
- increased appetite; or
- weight gain.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Remeron side effects](#) (more detail)

Related/similar drugs

Lexapro

Lexapro is used to treat anxiety and major depressive disorder. Learn about side effects ...

Reviews & ratings

7.6 / 10

[2,376 Reviews](#)

[View more](#)

Zoloft

Zoloft is an antidepressant used to treat major depression, obsessive-compulsive disorder, panic ...

Reviews & ratings

7.3 / 10

[1,972 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Seroquel

Seroquel is used to treat the symptoms of schizophrenia, bipolar disorder and major depressive ...

Reviews & ratings

7.0 / 10

[694 Reviews](#)

[View more](#)

Prozac

Prozac (fluoxetine) is an SSRI antidepressant used to treat depression, OCD, panic disorder ...

Reviews & ratings

7.2 / 10

[1,239 Reviews](#)

[View more](#)

Cymbalta

Cymbalta (duloxetine) is used to treat major depressive disorder, general anxiety disorder and ...

Reviews & ratings

6.3 / 10

[1,899 Reviews](#)

[View more](#)

Wellbutrin

Wellbutrin (bupropion) is used to treat major depressive disorder and seasonal affective disorder ...

Reviews & ratings

6.8 / 10

[756 Reviews](#)

[View more](#)

Celexa

Celexa (citalopram) is an antidepressant in a group of drugs called selective serotonin reuptake ...

Reviews & ratings

7.3 / 10

[866 Reviews](#)

[View more](#)

Concerta

Concerta is used to treat attention deficit disorder (ADD) and attention deficit hyperactivity ...

Reviews & ratings

6.9 / 10

[424 Reviews](#)

[View more](#)

Citalopram

Citalopram is an antidepressant
(selective serotonin reuptake inhibitor)
and is used to treat ...

Reviews & ratings

7.4 / 10

[2,231 Reviews](#)

[View more](#)

What other drugs will affect Remeron?

Using Remeron with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or [seizures](#).

Tell your doctor about all your current medicines. Many drugs can interact with mirtazapine, especially:

- [cimetidine](#);
- [diazepam](#);
- [St. John's wort](#);
- [tramadol](#);
- [tryptophan](#) (sometimes called L-tryptophan);
- **an antibiotic** - [clarithromycin](#), [rifampin](#), rifampicin, telithromycin;
- **antifungal medicine** - [itraconazole](#), [ketoconazole](#);
- **antiviral medicine to treat HIV/AIDS** - indinavir, nelfinavir, [ritonavir](#), saquinavir;
- **medicine to treat mood disorders, thought disorders, or mental illness** - such as [lithium](#), other [antidepressants](#), or [antipsychotics](#);
- **migraine headache medicine** - [sumatriptan](#), [rizatriptan](#), [Imitrex](#), Maxalt, and others; or
- **seizure medicine** - [carbamazepine](#), [phenytoin](#).

This list is not complete and many Other drugs may interact with mirtazapine. This includes prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible drug interactions are listed here.

 [Remeron drug interactions](#) (more detail)

Does Remeron interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Remeron

+

Popular FAQ

Does Remeron help with sleep?



More about Remeron (mirtazapine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(313\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Support group](#)
- [Drug class: tetracyclic antidepressants](#)
- [Breastfeeding](#)
- [En español](#)

Professional resources

- [Remeron prescribing information](#)
- [Mirtazapine \(AHFS Monograph\)](#)

Other brands

[Remeron SolTab](#)

Related treatment guides

- [Major Depressive Disorder](#)
- [Depression](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and

use Remeron only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 9.03.

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
10+ years	FDA approved 1996

User Reviews & Ratings

7.1 / 10

313 Reviews

Images

Remeron 30 mg (Organon TZ 5 TZ 5)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

- About Drugs.com
- Advertising policy
- Attribution & citations

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)


[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.