



[Home](#)

2. Revlimid

Revlimid

Generic name: [lenalidomide](#)

Dosage form: capsules (2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg, 25 mg)

Drug classes: [Miscellaneous antineoplastics](#), [Other immunosuppressants](#)

Medically reviewed by [Melisa Puckey, BPharm](#). Last updated on Apr 22, 2024.

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

What is Revlimid?

Revlimid is a medication used to treat types of [multiple myeloma](#), [mantle cell lymphoma](#), [follicular lymphoma](#), [marginal zone lymphoma](#), and anemia in [myelodysplastic syndromes](#) (MDS) in adults. Revlimid helps to slow cancer growth, block new blood vessel growth in tumors, and help the immune system fight cancer. Revlimid is available as capsules that are taken once a daily.

Revlimid received FDA approval on December 27, 2005. The active ingredient, sometimes called the generic name, is lenalidomide.

This medicine is only available through a restricted distribution program called the Lenalidomide REMS program, as it may cause birth defects or embryo-fetal death.

What is Revlimid used for?

Revlimid is used for specific adult patients with:

Multiple Myeloma in combination with dexamethasone. Also, for multiple myeloma maintenance therapy in patients who have had an autologous hematopoietic stem cell transplantation (auto-HSCT).

Myelodysplastic Syndromes with transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes (MDS) associated with a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities.

Mantle Cell Lymphoma (MCL) that has relapsed or progressed after two prior therapies, one of which included bortezomib.

Previously treated Follicular Lymphoma (FL), used in combination with a rituximab product.

Previously treated Marginal Zone Lymphoma (MZL), used in combination with combination with a rituximab product.

Revlimid is **not** indicated and is not recommended for the treatment of patients with CLL outside of controlled clinical trials.

Revlimid side effects

Common Revlimid side effects may include:

- [cold symptoms](#) such as stuffy nose, sneezing, sore throat;
- sleep problems, tiredness;
- weakness;
- headache;
- tremors;
- nosebleed;
- muscle cramps;
- joint pain;
- shortness of breath;
- fever, cough, tiredness;
- itching, rash, swelling; or
- stomach pain, nausea, vomiting, diarrhea, [constipation](#).

Serious Revlimid side effects.

Get emergency medical help if you have signs of an allergic reaction (hives, difficulty breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Lenalidomide may cause serious side effects. Call your doctor at once if you have:

- signs of a stroke or blood clot - sudden numbness or weakness, severe headache, problems with speech or vision, [shortness of breath](#), swelling or redness in your arm or leg;
- [heart attack](#) symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, sweating;
- liver problems - upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- low blood cell counts - fever, chills, swollen gums, mouth sores, skin sores, easy bruising, unusual bleeding;
- signs of a tumor getting worse - swollen glands, low fever, rash, or pain; or
- signs of tumor cell breakdown - lower back pain, blood in your urine, little or no urinating, numbness or tingly feeling around your mouth, muscle weakness or tightness, feeling short of breath, confusion, fainting.

This is not a complete list of side effects, and others may occur. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

 [Revlimid side effects](#) (more detail)

Related/similar drugs

Velcade

Velcade injection is used to treat multiple myeloma and mantle cell lymphoma. Includes Velcade side ...

Reviews & ratings

8.6 / 10

[9 Reviews](#)

[View more](#)

Gazyva

Gazyva is a targeted immunotherapy that works with the immune system to treat chronic lymphocytic ...

Reviews & ratings

[4 Reviews](#)

[View more](#)

Treanda

Treanda (bendamustine) is used to treat chronic lymphocytic leukemia and non-Hodgkin lymphoma ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Carvykti

Carvykti (ciltacabtagene autoleucel) is used for the treatment of relapsed or refractory multiple ...

Reviews & ratings

[Add a review](#)

[View more](#)

Abecma

Abecma is a CAR T cell therapy used to treat relapsed refractory multiple myeloma (RRMM); it is a ...

Reviews & ratings

[Add a review](#)

[View more](#)

Tecvayli

Tecvayli (teclistamab) is a treatment for relapsed or refractory multiple myeloma(cancer of the ...

Reviews & ratings

[5 Reviews](#)

[View more](#)

Reblozyl

Reblozyl is used to treat anemia in adults with beta-thalassemia or myelodysplastic syndromes ...

Reviews & ratings

[6 Reviews](#)

[View more](#)

Darzalex

Darzalex (daratumumab) is a target therapy used to treat multiple myeloma. Includes DARZALEX uses ...

Reviews & ratings

[6 Reviews](#)

[View more](#)

Talvey

Talvey is used to treat relapsed or refractory multiple myeloma. Includes Talvey side effects ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Warnings

Never use Revlimid if you are pregnant. Even one dose of lenalidomide can cause severe, life-threatening birth

defects or death of a baby if the mother or the father is taking this medicine at the time of conception or during pregnancy.

Use birth control to prevent pregnancy, whether you are a man or a woman.

For women: Use two forms of birth control, beginning 4 weeks before you start taking Revlimid and ending 4 weeks after you stop taking it. See the 'Before taking this medicine' section for more information.

For men: Use a condom to prevent pregnancy during your treatment and for up to 4 weeks after your treatment ends. See the 'Before taking this medicine' section for more information.

Revlimid may cause blood clots. Stop using this medicine and call your doctor at once if you have symptoms such as sudden numbness, severe headache, problems with vision or speech, chest pain, shortness of breath, coughing up blood, or swelling in your arm or leg.

Revlimid can lower blood cells that help your body fight infections and help your blood to clot. Call your doctor if you have unusual bruising or bleeding or signs of infection (fever, chills, body aches). You will need frequent blood tests while you are taking Revlimid.

Before taking this medicine

You should not use Revlimid if you are allergic to lenalidomide.

To make sure Revlimid is safe for you, tell your doctor if you have ever had:

- an allergic reaction to thalidomide;
- kidney disease (or if you are on dialysis);
- liver disease;
- a blood clots or stroke;
- high blood pressure, high cholesterol or triglycerides;
- a thyroid disorder;
- lactose intolerance; or
- if you smoke.

Using Revlimid may increase your risk of developing other types of cancer, such as leukemia or lymphoma. Talk with your doctor about your specific risk.

Pregnancy

Revlimid can cause severe, life-threatening birth defects or death of a baby if the mother or the father is taking this medicine at the time of conception or during pregnancy. Even one dose of lenalidomide can cause major birth defects in the baby's arms and legs, bones, ears, eyes, face, and heart. Never use Revlimid if you are pregnant. Tell your doctor right away if your period is late while taking Revlimid.

For Women: If you have not had a hysterectomy, you will be required to use two reliable forms of birth control, beginning 4 weeks before you start taking Revlimid and ending 4 weeks after you stop taking it. Even women with fertility problems are required to use birth control while taking this medicine. You must also have a negative pregnancy test 10 to 14 days

before treatment and again 24 hours before. While you are taking Revlimid, you will have a pregnancy test every 2 to 4 weeks.

The birth control method you use must be proven highly effective, such as birth control pills, an intrauterine device (IUD), a tubal ligation, or a sexual partner's vasectomy. The extra form of birth control you use must be a barrier method such as a latex condom, a diaphragm, or a cervical cap.

Stop using Revlimid and call your doctor at once if you quit using birth control, if your period is late, or if you think you might be pregnant. Not having sexual intercourse (abstinence) is the most effective method of preventing pregnancy.

For Men: If a man fathers a baby while using Revlimid, the baby may have birth defects. Use a condom to prevent pregnancy during your treatment and for up to 4 weeks after your treatment ends. You must agree in writing to always use latex condoms when having sex with a woman who is able to get pregnant, even if you have had a vasectomy. Contact your doctor if you have had unprotected sex, even once, or if you think your female sexual partner may be pregnant.

Breastfeeding

You should not breastfeed while using Revlimid.

 [Revlimid pregnancy and breastfeeding warnings](#) (more detail)

How should I take Revlimid?

Take the medicine at the same time each day. You may take Revlimid with or without food.

Take each dose with a full glass of water. Swallow the capsule whole without breaking it open.

Lenalidomide can increase your risk of bleeding or infection. Your blood will need to be tested often.

Medicine from an open capsule can be dangerous if it gets on your skin. If this occurs, wash your skin with soap and water. Ask your doctor or pharmacist how to handle and dispose of a broken capsule safely.

Take Revlimid exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or longer than recommended. Never share this medicine with another person, even if they have the same disorder you have.

Revlimid Dosing information

Usual Adult Revlimid Dose for Multiple Myeloma

Use: Multiple Myeloma in combination with dexamethasone

Dose: 25 mg orally once a day on Days 1 to 21 of repeated 28-day cycles until disease progression or unacceptable toxicity

Use: Maintenance therapy for Multiple Myeloma following auto-HSCT

Dose: 10 mg once a day continuously (Days 1 to 28 of repeated 28-day cycles) for 3 cycles, then increase to 15 mg once a day if tolerated until disease progression or unacceptable toxicity.

Comments:

For patients who are not eligible for auto-HSCT, therapy should continue until disease progression or unacceptable toxicity.

For patients who are eligible for auto-HSCT, hematopoietic stem cell mobilization should occur within 4 cycles.

Following auto-HSCT, initiate maintenance therapy after adequate hematologic recovery (ANC 1000/mcL or more AND/OR platelet count 75,000/mcL or more).

Consult the manufacturer's product information for dexamethasone dosing recommendations.

Usual Adult Dose for Myelodysplastic Disease:

Dose: 10 mg orally once a day; therapy is continued or modified based on clinical and laboratory findings until disease progression or unacceptable toxicity

Use: Treatment of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes (MDS) associated with a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities.

Usual Adult Dose for Lymphoma:

Follicular Lymphoma or Marginal Zone Lymphoma:

Dose: 20 mg orally once a day on Days 1 through 21 of repeated 28-day cycles for up to 12 cycles in combination with a rituximab product.

Use: In combination with a rituximab product for the treatment of previously treated follicular lymphoma (FL). In combination with a rituximab product for the treatment of previously treated marginal zone lymphoma (MZL).

Mantel Cell Lymphoma:

Dose: 25 mg orally once a day on Days 1 to 21 of repeated 28-day cycles until disease progression or unacceptable toxicity; treatment is continued, modified, or discontinued based on clinical and laboratory findings.

Use: The treatment of patients with mantle cell lymphoma (MCL) whose disease has relapsed or progressed after 2 prior therapies, one of which included bortezomib.

Revlimid is available in 2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg, and 25 mg capsules.

 [Detailed Revlimid dosage information](#)

What happens if I miss a dose?

Take the missed dose as soon as you remember. If you are more than 12 hours late, skip the missed dose. Do not take extra medicine to make up for the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What to avoid

You must not donate blood or sperm while you are using Revlimid, and for at least 4 weeks after your last dose. Avoid exposing another person to your blood or semen through casual or sexual contact.

This medicine can pass into body fluids (urine, feces, vomit). Caregivers should wear rubber gloves while cleaning up a patient's body fluids, handling contaminated trash or laundry or changing diapers. Wash hands before and after removing gloves. Wash soiled clothing and linens separately from other laundry.

What other drugs will affect Revlimid?

Tell your doctor if you also use pembrolizumab (Keytruda).

If you use hormonal birth control (pills, implants, injections) to prevent pregnancy: There are certain drugs that can make hormonal birth control less effective in your body. Tell your doctor about all other medicines you use. You may need to replace your hormonal birth control method with another effective form of contraception.

Other drugs may interact with lenalidomide, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

 [Revlimid drug interactions](#) (more detail)

Does Revlimid interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Revlimid

+

Enter a drug name

Add

Revlimid Package Insert

Review the [Revlimid Package Insert](#) for more detailed information about this medicine. Discuss any medical questions you have with your doctor or other health care provider. This is not all the information you need to know about this medicine for safe and effective use, and it does not take the place of talking to your doctor about your treatment.

Handling and Disposal

Care should be exercised in the handling of this medicine. Capsules should not be opened or broken. If powder from capsules contacts the skin, wash the skin immediately and thoroughly with soap and water. If Revlimid contacts the mucous membranes, flush thoroughly with water. Procedures for the proper handling and disposal of anticancer drugs should be considered. Several guidelines on the subject have been published. Dispense no more than a 28-day supply.

Ingredients

Active ingredient: lenalidomide

Inactive ingredients: lactose anhydrous, microcrystalline cellulose, croscarmellose sodium, and magnesium stearate.

The 5 mg and 25 mg capsule shells contain gelatin, titanium dioxide, and black ink.

The 2.5 mg and 10 mg capsule shells contain gelatin, FD&C blue #2, yellow iron oxide, titanium dioxide, and black ink.

The 15 mg capsule shell contains gelatin, FD&C blue #2, titanium dioxide, and black ink.

The 20 mg capsule shell contains gelatin, FD&C blue #2, yellow iron oxide, titanium dioxide and black ink.

Storage

Store at 20°C - 25°C (68°F - 77°F)

Company

Bristol-Myers Squibb Company Princeton, NJ 08543 USA.

REVLIMID® is a trademark of Celgene Corporation, a Bristol-Myers Squibb company.

Popular FAQ

Is Revlimid a chemotherapy drug?



What is the success rate of Monjuvi?



How does Revlimid work?



What is the best time of day to take Revlimid?



How long does Revlimid stay in your system?



How long do you take Revlimid for?



Can Revlimid cause weight gain?



More FAQ

- [What are biologic drugs and how do they work?](#)

[View more FAQ...](#)

References

1. [Revlimid Product Label](#)
2. [Thalidomide and lenalidomide: Mechanism-based potential drug combinations](#)

More about Revlimid (lenalidomide)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(28\)](#)
- [Drug images](#)
- [Side effects](#)

- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [FDA approval history](#)
- [Drug class: miscellaneous antineoplastics](#)
- [En español](#)

Professional resources

- [Revlimid prescribing information](#)
- [Lenalidomide \(AHFS Monograph\)](#)

Related treatment guides

- [Follicular Lymphoma](#)
- [Anemia](#)
- [Lymphoma](#)
- [Mantle Cell Lymphoma](#)

Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

DRUG STATUS		
Availability		
Rx	Prescription only	
Pregnancy & Lactation		
	Risk data available	
CSA Schedule*		
N/A	Not a controlled drug	▼
Approval History		
	FDA approved 2005	▼

User Reviews & Ratings

7.0 / 10

[28 Reviews](#)

Related News

Images

Revlimid 10 mg (REV 10 mg)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT

Copyright © 2000-2025 Drugs.com. All rights reserved.

