

[Home](#)

2. Lialda

Lialda

Generic name: [mesalamine](#) (oral) [*me-SAL-a-meen*]**Drug class:** [5-aminosalicylates](#)Medically reviewed by [Kaci Durbin, MD](#). Last updated on Aug 23, 2023.[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is Lialda?

Lialda affects a substance in the body that causes inflammation, tissue damage, and diarrhea.

Lialda is used to treat mild to moderate active [ulcerative colitis](#) in adults and children who weight at least 24kg.

Lialda is also used in adults as a maintenance treatment to prevent the symptoms of ulcerative colitis from recurring.

Warnings

Stop using Lialda and call your doctor at once if you have severe stomach pain, stomach cramping, bloody [diarrhea](#) (may occur with fever, [headache](#), and [skin rash](#)).

Before taking this medicine

You should not use Lialda if you are allergic to mesalamine, [aspirin](#), [sulfasalazine](#), or [salicylates](#) (such as Disalcid, Doans Pills, Dolobid, [Kaopectate](#), [Nuprin Backache](#), [Pepto-Bismol](#), Tricosal, Trilisate, and others).

To make sure Lialda is safe for you, tell your doctor if you have ever had:

- a kidney stone or kidney disease;
- liver disease;
- a blockage in your stomach or intestines (such as pyloric stenosis); or
- a skin condition such as [eczema](#).

Tell your doctor if you are pregnant or plan to become pregnant. It is not known whether mesalamine will harm an unborn baby. However, having untreated or uncontrolled ulcerative colitis during pregnancy may cause complications such as low birth weight or premature birth. The benefit of treating ulcerative colitis may outweigh any risks to the baby.

It may not be safe to breastfeed while using Lialda. Ask your doctor about any risk. If you are breastfeeding, tell your doctor if you notice diarrhea in the nursing baby.

 [Lialda pregnancy and breastfeeding warnings](#) (more detail)

How should I take Lialda?

Take Lialda exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Lialda.

Lialda should be taken with a meal.

Drink plenty of liquids to help prevent [kidney stones](#).

Swallow the Lialda tablet whole and do not crush, chew, or break it. **Tell your doctor if you have trouble swallowing the tablet.**

Call your doctor if your ulcerative colitis symptoms do not improve, or if they get worse.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Lialda.

You may need frequent blood tests to check your blood cells or kidney function.

Store at room temperature away from moisture and heat.

Dosing information

Usual Adult Dose of Lialda for Ulcerative Colitis -- Active:

1200 mg tablets: 2.4 to 4.8 g (two to four tablets) orally once a day with food.

Usual Adult Dose of Lialda for Ulcerative Colitis -- Maintenance:

1200 mg tablets: 2.4 g (two tablets) orally once a day with food.

Usual Pediatric Dose of Lialda for Ulcerative Colitis -- Active:

Patient weight - 24 kg to 35 kg: from week 0 to week 8, two 1.2 g tablets once daily. After week 8, one 1.2 g tablet once daily.

Greater than 35 kg to 50 kg: from week 0 to week 8, three 1.2 g tablets once daily. After week 8, two 1.2 g tablets once daily.

Greater than 50 kg: from week 0 to week 8, four 1.2 g tablets once daily. After week 8, two 1.2 g tablets once daily.

 [Detailed Lialda dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Lialda?

If you have a skin condition such as eczema, mesalamine could make you sunburn more easily. Avoid sunlight or tanning beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Lialda side effects

Get emergency medical help if you have **signs of an allergic reaction to Lialda**: [hives](#); difficulty breathing; swelling of your face, lips, tongue, or throat.

Stop using Lialda and call your doctor at once if you have:

- severe stomach pain, stomach cramping, bloody diarrhea;
- fever, headache, general ill feeling;
- rash, itching, [eye redness](#);
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- **low white blood cell counts** - fever, chills, mouth sores, skin sores, [sore throat](#), cough, feeling light-headed, trouble breathing;
- **signs of a kidney stone** - severe pain in your side and back, frequent need to urinate, foul-smelling urine, dark or cloudy urine;
- **kidney problems** - increased or decreased urination, swelling, weight gain; or
- **liver problems** - loss of appetite, upper stomach pain, tiredness, easy bruising or bleeding, dark urine, clay-colored stools, [jaundice](#) (yellowing of the skin or eyes).

Low white blood cell counts may be more likely in older adults.

Common Lialda side effects may include:

- [burping](#), [constipation](#), [nausea](#), [vomiting](#), stomach pain, diarrhea, gas;
- [dizziness](#);
- [cold symptoms](#) such as stuffy nose, sneezing, sore throat;
- [back pain](#);
- headache;
- rash; or
- abnormal liver function tests.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Lialda side effects](#) (more detail)

Related/similar drugs

Skyrizi

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

Reviews & ratings

7.9 / 10

[56 Reviews](#)

[View more](#)

OmvoH

OmvoH is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

FEATURED

Entyvio

Entyvio (vedolizumab) is used to treat ulcerative colitis (UC) and Crohn's disease, reducing ...

Reviews & ratings

6.2 / 10

[162 Reviews](#)

[View more](#)

Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used alone or with other ...

Reviews & ratings

5.7 / 10

[145 Reviews](#)

[View more](#)

Stelara

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

Reviews & ratings

7.6 / 10

[252 Reviews](#)

[View more](#)

Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

Reviews & ratings

7.0 / 10

[223 Reviews](#)

[View more](#)

Humira

Humira is a tumor necrosis factor blocker that is used to treat many inflammatory conditions such ...

Reviews & ratings

6.4 / 10

[698 Reviews](#)

[View more](#)

What other drugs will affect Lialda?

Mesalamine can harm your kidneys, especially if you also use certain medicines for infections, cancer, [osteoporosis](#), organ transplant rejection, bowel disorders, [high blood pressure](#), or pain or arthritis (including [Advil](#), [Motrin](#), and [Aleve](#)).

Tell your doctor about all your other medicines, especially:

- [azathioprine](#) or 6-mercaptopurine.

This list is not complete. Other drugs may interact with mesalamine, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible drug interactions are listed here.

 [Lialda drug interactions](#) (more detail)

Does Lialda interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Lialda

+

Enter a drug name

Add

Popular FAQ

- What are 7 mesalamine side effects to be aware of?

▼
- How long does it take for mesalamine to work?

▼
- When is the best time to take mesalamine?

▼
- Why is mesalamine so expensive?

▼
- How long can you take mesalamine?

▼
- Is mesalamine an immunosuppressant?

▼
- Can I stop taking mesalamine?

▼
- What is the maximum dosage of mesalamine?

▼

More about Lialda (mesalamine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(105\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [FDA approval history](#)
- [Drug class: 5-aminosalicylates](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

Pentasa, Apriso, Canasa, Rowasa, ... +2 more

Professional resources

- [Lialda prescribing information](#)
- [Mesalamine \(AHFS Monograph\)](#)

Other brands

Pentasa, Apriso, Asacol, Canasa, ... +4 more

Related treatment guides

- [Ulcerative Colitis, Active](#)
- [Ulcerative Colitis, Maintenance](#)
- [Inflammatory Bowel Disease](#)
- [Ulcerative Proctitis](#)
- [Ulcerative Colitis](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Lialda only for the indication prescribed.


Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)


Copyright 1996-2025 Cerner Multum, Inc. Version: 16.01.

DRUG STATUS

Availability
Rx Prescription only

Pregnancy & Lactation
 Risk data available

CSA Schedule*
N/A Not a controlled drug

Approval History
 Drug history at FDA



User Reviews & Ratings

6.8 / 10

[105 Reviews](#)

Images

[Lialda 1.2 g \(S476\)](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT

Copyright © 2000-2025 Drugs.com. All rights reserved.

