

[Home](#)

2. [Natural Products \(Pro\)](#)
3. [Kava](#)

Kava

Scientific Name(s): Piper methysticum Forst.f.

Common Name(s): Ava, Awa, Intoxicating pepper, Kava, Kava kava, Kava pepper, Kava root, Kava-kava, Kavain, Kawa, Kawain, Kew, Rauschpfeffer, Sakau, Tonga, Wurzelstock, Yangona

[Medically reviewed](#) by Drugs.com. Last updated on Jul 15, 2024.

[Overview](#) [History](#) [Dosage](#) [Pregnancy](#) [Interactions](#) [Side effects](#) [FAQ](#)

Clinical Overview

Use

A number of meta-analyses and systematic reviews of kava use in anxiety have found in favor of kava over placebo, but results are not consistent. Kava has also been studied for effects on cognitive function and for potential cancer applications. However, concerns over hepatotoxicity have limited clinical studies.

Dosing

A maximum daily dose of kavalactones 250 mg is suggested to avoid potential hepatotoxicity. Studies in children are lacking, and use is not recommended.

Contraindications

Kava and kava-containing products are not recommended for use in children or in patients with hepatic disease. Kava should be used cautiously in patients with renal or liver disease, blood disorders, Parkinson disease, or depression.

Pregnancy/Lactation

Documented adverse effects. Avoid use.

Interactions

Kava extracts have been shown to interfere with cytochrome P450 (CYP-450) enzymes; however, specific reports on the metabolism of pharmaceuticals are sparse. Case reports exist on interactions with alprazolam, alcohol, barbiturates, and levodopa. Concomitant administration of kava with haloperidol, risperidone, and metoclopramide, among other drugs, may be associated with adverse reactions.

Adverse Reactions

Heavy kava use may cause a scaly skin rash. A variety of adverse reactions, including visual disturbances, urinary

retention, GI discomfort, exacerbation of Parkinson disease, extrapyramidal effects, and rhabdomyolysis, have been reported.

Toxicology

Rare cases of severe liver toxicity have been reported. Fatal intoxication with concomitant IV administration of kavalactones and ethanol has been reported.

Scientific Family

- Piperaceae (black pepper)

Botany

The dried rhizome and roots of *P. methysticum*, a large shrub widely cultivated on many Pacific islands, including Hawaii, Tahiti, and New Guinea, are consumed in various forms as kava. The plant can grow as tall as 3 m, has large, heart-shaped leaves, and is vegetatively propagated exclusively by root cuttings. It is thought to be derived from the wild species *Piper wichmannii* C. DC. Many cultivars of kava are recognized, and the rootstock color ranges from white to dark yellow, depending on the amount of kavalactones present. The comparative chemistry and ethnopharmacology have been studied in detail, and 121 named cultivars from 51 islands have been grouped into 6 chemotypes.[1](#), [2](#), [3](#), [4](#), [5](#)

History

The kava beverage is prepared from the plant's roots, which are chewed or pulverized and then steeped in water or coconut milk. The cloudy mixture is filtered and served at room temperature. Kava has been an important part of Pacific Island ceremonial cultures for many centuries, with elaborate rituals attending its consumption. Its main use has been to induce a relaxed state in ceremony participants by initially causing a numbing and astringent effect in the mouth, followed by anxiolytic and muscle relaxant effects. Eventually, a state of sleep is induced and no hangover effects are experienced. The kava beverage has been used to symbolize respect and hospitality for visiting dignitaries, with traces of kava extract identified on archaeological artifacts from the Fiji islands by mass spectrometry.

Research into the use of kava has been conducted since the late 19th century. In the early 1900s, kava was used as a diuretic and for gonorrhea and nervous disorders. It has been one of the top botanical sellers in the United States and Europe for anxiety and sleep disorders. However, many countries have regulated its sale because of reports of hepatotoxicity. In 2002, the US Food and Drug Administration issued a consumer advisory warning of the potential for liver injury.[5](#), [6](#), [7](#), [8](#)

Chemistry

Fresh kava root contains about 80% water. Dried kava root contains 12% water, 43% starch, and 20% fibers, with the balance being made up of protein, sugars, and minerals. The primary active constituents in the resin include kavalactones, chalcones and other flavanones, and conjugated diene ketones. At least 18 lipid-soluble kavalactones, the primary bioactive constituents, have been isolated from kava root extract, accounting for 3% to 20% of the dry weight. The 6 major kavalactones are kawain, 7,8-dihydrokawain, methysticin, 7,8-dihydromethysticin, yangonin, and demethoxyyangonin, which occur in varying proportions in different cultivars. Toxic alkaloids have been identified in the leaves and stem peelings. Many methods have been developed for the analysis of kavalactones, including thin layer chromatography, gas chromatography, high-performance liquid chromatography, gas chromatography–mass

spectrometry, and nuclear magnetic resonance. Proposals for standardization of constituents exist. [5](#), [9](#), [10](#), [11](#), [12](#), [13](#), [14](#)

Uses and Pharmacology

Chewing kava causes numbness in the mouth because of the local anesthetic action of kavalactones, which is similar to that produced by cocaine and longer lasting than that of benzocaine. In addition, kava produces a mild euphoria characterized by feelings of contentment and fluent, lively speech. Sight, smell, and sound are also reported to be heightened. Higher doses may lead to muscle weakness, especially in the legs, although some observers relate this to sitting for long periods during the kava ceremony rather than to kava itself. Very high doses may induce a deep sleep. [15](#), [16](#)

Anxiety

Animal data

There is no animal data regarding the use of kava for anxiety, aside from studies attempting to elucidate the mechanism of action. Suggested mechanisms include enhanced ligand binding activity at gamma-aminobutyric acid (GABA) receptors, reversible inhibition of monoamine oxidase (MAO)-B, and reduced uptake of norepinephrine and dopamine. Kavalactones, however, do not appear to bind directly to the GABA receptors. [17](#), [18](#), [19](#), [20](#), [21](#), [22](#), [23](#), [24](#), [25](#), [26](#), [27](#), [28](#)

Clinical data

A number of meta-analyses and systematic reviews of kava use for anxiety have been conducted, with findings favoring kava over placebo (5 point reduction in Hamilton Anxiety Scale [HAM-A] score [confidence interval (CI), 1.1 to 8.8]). [28](#), [29](#), [30](#) Efficacy has been shown to be similar to that of oxazepam, opipramol (a tricyclic antidepressant available in Europe), and buspirone. [28](#), [29](#), [30](#), [31](#), [32](#) Studies included in these reviews tested older preparations of kava, while trials evaluating water-derived preparations using accepted dosages are limited. [28](#), [30](#) The Kava Anxiety Depression Spectrum Study evaluated aqueous kava extract 250 mg per day over 3 weeks in 60 adults with generalized anxiety disorder (GAD) and found reductions in HAM-A measurements (reduction of 9.9 points [CI, 7.1 to 12.7]) versus placebo (reduction of 0.8 points [CI, -2.7 to 4.3]). [33](#) Some of the same investigators conducted a 6-week, placebo-controlled trial in GAD that included determination of pharmacogenetic profiles. Response, defined as at least 50% reduction in HAM-A score, occurred in 37% with kava and 23% of placebo patients ($P = 0.046$). Remission (HAM-A up to 7) was seen in 26% and 6% with kava and placebo, respectively ($P = 0.04$). Two of five alleles of GABA transporter were associated with kava response. [89](#) Limited studies suggest efficacy in menopausal-related anxiety. [30](#), [34](#), [35](#), [36](#), [37](#) Studies with negative findings for kava in generalized anxiety disorder have been published, including a single-dose study for acute management of anxiety. [38](#), [39](#), [40](#) Head-to-head trials evaluating efficacy against selective serotonin reuptake inhibitors have not been conducted. [28](#), [31](#)

Cancer

Animal data

Rats fed kava for 14 weeks showed a reduction in colon cancer risk markers. [41](#) Constituents of kava extracts, including the flavokawains, yangonin, and methysticin, have been studied for anticancer effects. Apoptosis and reduced proliferation have been demonstrated in lung, liver, pancreatic, prostate, uterine, and squamous cell lines. Chalcone analogs have also been studied. [42](#), [43](#), [44](#), [45](#), [46](#), [47](#)

Clinical data

Epidemiological data suggest a correlation between kava consumption and a low incidence of cancer.[48](#), [49](#) In vitro data from a study of flavokawain B (FKB), a kava extract, found it arrested the cell cycle and induced apoptosis in human sarcoma cells, and was less toxic to bone marrow and intestinal epithelial cells than doxorubicin.[90](#)

Cognition

Animal data

No animal data exist on the effect of kava on cognition, aside from studies attempting to elucidate the mechanism of action. Suggested mechanisms include enhanced ligand binding activity at GABA receptors, reversible inhibition of MAO-B, and reduced uptake of norepinephrine and dopamine. Kavalactones, however, do not appear to bind directly to the GABA receptors.[17](#), [18](#), [19](#), [20](#), [21](#), [22](#), [23](#), [24](#), [25](#), [26](#), [27](#), [28](#)

Clinical data

Reviews of at least 10 studies on the effects of kava on cognition have been published.[28](#), [50](#), [51](#)

Heterogeneity of dosages/potency and preparations used precludes meta-analysis. Seven studies evaluated the short-term effects of kava on cognition, while 3 evaluated long-term effects. Despite the heterogeneity, kava appears to have little or no overall effect on cognition. At higher dosages, reaction time may be impaired.[28](#), [50](#), [51](#) Kava has been promoted for use in attention deficit hyperactivity disorder and as an alternative to methylphenidate; however, clinical trials are lacking and until safety concerns are resolved, such use is not recommended.[5](#), [51](#)

Other uses

Kawain showed an antithrombotic effect on platelets, dose-dependently blocking platelet aggregation, adenosine 5'-triphosphate release, and synthesis of prostaglandins at high micromolar concentrations.[52](#)

Despite a reputation as an antimicrobial agent in urinary tract infections, kava extracts demonstrated minimal antifungal activity and no antimicrobial or antiviral activity.[53](#) Antitrypanosomal activity has been demonstrated.[54](#)

The Veteran's Administration and Department of Defense (VA/DoD) clinical practice guideline for the management of chronic insomnia disorder and obstructive sleep apnea (2019) recommend against the use of kava for the treatment of chronic insomnia disorder (Strong).[Mysliwiec 2020](#)

Dosing

The Australian Therapeutic Goods Administration requires the maximum strength of tablet or capsule preparations to be kavalactone 125 mg, with a recommended maximum daily dose of kavalactone 250 mg. No adverse reactions have been reported at this dosage, which is less than the traditional consumption.[14](#)

Kava does not appear to be addictive at therapeutic dosages.[55](#)

Studies in children are lacking, and use is not recommended.[51](#), [56](#)

Pregnancy / Lactation

Documented adverse effects. Avoid use.[57](#), [58](#), [59](#)

Interactions

Anti-Parkinson agents (dopamine agonist): Kava may enhance the adverse/toxic effect of anti-Parkinson agents (dopamine agonist). Kava may diminish the therapeutic effect of anti-Parkinson agents (dopamine agonist). Monitor therapy.(66, 94)

CNS depressants: Kava may enhance the CNS depressant effect of CNS depressants. Monitor therapy.(28, 67, 94, 95, 96, 100)

Paroxetine: Kava may enhance the adverse/toxic effect of paroxetine. Monitor therapy.(97)

Kava extracts have been shown to interfere with CYP-450 enzymes; however, reports on specific effects on the metabolism of pharmaceuticals are limited.(60, 61, 62, 63) Case reports exist on interactions with alprazolam, alcohol, barbiturates, and levodopa. Studies in healthy volunteers have demonstrated a lack of interactions with midazolam and digoxin.(15, 63, 64, 65, 66, 67, 68) Interactions with MAO inhibitors, antiplatelets, and hepatotoxic agents may exist. Drugs affecting dopamine, such as haloperidol, risperidone, and metoclopramide, among others, are associated with increased adverse reactions, especially Parkinson-like symptoms, when given concomitantly with kava-containing products.(69, 70)

 [Kava drug interactions](#) (more detail)

Adverse Reactions

Acute adverse reactions of kava consumption may include oral anesthetic effects (especially of the tongue), sedation, euphoria, tremors, and ataxia.15, 16 Visual impairment (near-point accommodation, enlargement of the pupils, and disturbances in oculomotor equilibrium) has also been described.10, 15 One study saw a significantly higher incidence of headache compared with placebo in patients with GAD, but the investigators did not believe headache was related to study treatment.89 There was a significant increase in libido in female patients receiving kava in this same GAD study.91

Heavy consumption of kava has long been known to produce a scaly skin rash similar to pellagra and known as "kava dermatopathy" on the palms of the hands, soles of the feet, and back. Supplementation with niacin did not reverse the condition,15, 71 but cessation of kava use resulted in its reduction or disappearance. Because flavokawain pigments are suspected to be responsible for this toxicity,72 they are commonly removed in the production of commercial extracts despite the lack of scientific proof.73

Urinary retention, exacerbation of Parkinson disease, extrapyramidal effects, myoglobinuria, severe rhabdomyolysis, and GI discomfort, among other effects, have been reported.10, 15, 74, 75, 76

Kava should be used cautiously in patients with renal or liver disease, blood disorders, Parkinson disease, or depression. Regular liver function tests may be warranted.7, 15, 77

Data collected between 2004 and 2013 from 8 US centers in the Drug-induced Liver Injury Network revealed that 15.5% (130) of hepatotoxicity cases were caused by herbals and dietary supplements, whereas 85% (709) of cases were related to prescription medications. Of the 130 cases of liver injury related to supplements, 65% were from non-bodybuilding supplements and occurred most often in Hispanics/Latinos compared with non-Hispanic whites and non-Hispanic blacks. Liver transplant was also more frequent with toxicity from non-bodybuilding supplements (13%) than with conventional medications (3%) ($P<0.001$). Overall, the proportion of severe liver injury cases was significantly higher for supplements than for conventional medications ($P=0.02$). Of the 217 supplement products implicated in liver injury, 175 had identifiable ingredients, of which kava kava was among the 32 (18%) single-ingredient products.93 The European Association for the Study of the Liver (EASL) clinical practice guideline for drug-induced liver injury (2019) recommends

physicians consider herbal and dietary supplements as potential causative agents associated with liver injury (Level 4; Grade C), including kava, which has some of the highest levels of evidence supporting hepatotoxicity.⁹⁹

Related/similar drugs

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Creatine

Creatine is a quickly available source of energy for muscle contractions and is used to enhance ...

Reviews & ratings

[6 Reviews](#)

[View more](#)

Toxicology

Kava has been implicated in several reports of fulminant hepatic failure in Europe and the United States. The FDA as well as the Australian and Canadian authorities have issued warnings to consumers and health care providers on the potential for liver damage from kava products.^{5, 30, 78, 79} Reviews of kava safety issues have been published. Although the incidence of hepatotoxicity in clinical trials has been too low to show causality, patients with any predisposition to hepatic dysfunction should avoid kava use.^{5, 12, 30, 80, 81, 82} Rats fed kava for 14 weeks showed no hepatic lesions on histology.⁴¹

Several theories have been suggested as to why kava may cause hepatotoxicity and to account for its apparent safe use among South Pacific populations for centuries. One theory is that because heavy demand for kava products led to the addition of kava leaves and stems to root products, the alkaloid pipermethystine, which is found in the leaves and stems but not in roots, may be responsible. Cytotoxicity of pipermethystine (50 to 100 mcM) to HepG2 cells has been demonstrated in support of this theory. Another theory, known as the "glutathione theory," suggests that aqueous extracts traditionally used in the South Pacific contain glutathione with the potential to react with kavalactones, providing hepatoprotection, as opposed to products that are chemically derived and appear on the market. Additionally, several studies have indicated that kavalactones are inhibitors of CYP-450 isoenzymes, increasing the risk for drug-herbal interactions, especially with kava formulations containing high concentrations of methysticin and dihydromethysticin. Hepatotoxicity may also result with coadministration of alcohol because kava decreases the conversion of ethanol to acetaldehyde; alcohol may also reduce detoxification of kavalactones. A fatal case of intentional suicide using

kavalactones injected intravenously with ethanol was reported in 36-year-old man with a history of depression.⁹² Another theory postulates that the hepatotoxicity of kava is due to the increased risk of liver damage in patients with deficiencies in CYP2D6, a major metabolic pathway for kavalactones. This phenomenon is seen in approximately 10% of white patients but rarely in Polynesian patients, which may explain the higher incidence of liver toxicity in European patients compared with those in the Pacific Islands. Specifically, kavalactones appear to inhibit cyclooxygenase enzymes COX-1 and COX-2. The mediators derived from the production of COX-2 have been shown to be hepatoprotective, the inhibition of which may contribute to kava hepatotoxicity. Immunological and/or toxicological reactions may also be implicated in hepatotoxicity from kava.^{8, 83, 84, 85, 86, 87, 88}

Concerns over potential toxicity to the kidneys, brain, and hematopoietic system also exist.⁵ Data are insufficient to clarify the safe cosmetic use of kava products.¹³

References

Disclaimer

This information relates to an herbal, vitamin, mineral or other dietary supplement. This product has not been reviewed by the FDA to determine whether it is safe or effective and is not subject to the quality standards and safety information collection standards that are applicable to most prescription drugs. This information should not be used to decide whether or not to take this product. This information does not endorse this product as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this product. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this product. This information is not specific medical advice and does not replace information you receive from your health care provider. You should talk with your health care provider for complete information about the risks and benefits of using this product.

This product may adversely interact with certain health and medical conditions, other prescription and over-the-counter drugs, foods, or other dietary supplements. This product may be unsafe when used before surgery or other medical procedures. It is important to fully inform your doctor about the herbal, vitamins, mineral or any other supplements you are taking before any kind of surgery or medical procedure. With the exception of certain products that are generally recognized as safe in normal quantities, including use of folic acid and prenatal vitamins during pregnancy, this product has not been sufficiently studied to determine whether it is safe to use during pregnancy or nursing or by persons younger than 2 years of age.

Frequently asked questions

- [Does Kava interact with any drugs?](#)
- [Does Kava interact with any drugs?](#)

More about kava

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(14\)](#)
- [Drug class: herbal products](#)

Related treatment guides

- [Herbal Supplementation](#)
- [Anxiety](#)
- [Anxiety and Stress](#)

1. *Piper methysticum* Forst.f. USDA, NRCS. 2012. The PLANTS Database. (<http://plants.usda.gov>. October 12, 2012). National Plant Data Team, Greensboro, NC 27401-4901 USA. Accessed January 3, 2013.
2. Duke JA. *Handbook of Medicinal Herbs*. 2nd ed. Boca Raton, FL: CRC Press; 2002:437-439.
3. Lebot V, Levesque J. Genetic control of kavalactone chemotypes in *Piper methysticum* cultivars. *Phytochem*. 1996;43(2):397-403.
4. Lebot V, Johnston E, Zheng QY, McKern D, McKenna DJ. Morphological, phytochemical, and genetic variation in Hawaiian cultivars of 'awa (kava, *Piper methysticum*, Piperaceae). *Econ Bot*. 1999;53(4):407-418.
5. Fu PP, Xia Q, Guo L, Yu H, Chan PC. Toxicity of kava kava. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev*. 2008;26(1):89-112.18322868
6. Holmes LD. The function of kava in modern Samoan culture. In: Efron DH, Holmstedt B, Kline NS, eds. *Ethnopharmacologic Search for Psychoactive Drugs*. New York, NY: Raven Press; 1979:107.
7. Blumenthal M. *The ABC Clinical Guide to Herbs*. Austin, TX: American Botanical Council; 2003:262-269.
8. Lewin, L. *Über Piper Methysticum (Kawa)*. Berlin: A. Hirschwald; 1886.
9. Denham A, McIntyre M, Whitehouse J. Kava—the unfolding story: report on a work-in-progress. *J Altern Complement Med*. 2002;8(3):237-263.12165183
10. Cloutre DL. Kava kava: examining new reports of toxicity. *Toxicol Lett*. 2004;150(1):85-96.15068826
11. Dharmaratne HR, Nanayakkara NP, Khan IA. Kavalactones from *Piper methysticum*, and their ¹³C NMR spectroscopic analyses. *Phytochemistry*. 2002;59(4):429-433.11830162
12. Rowe A, Zhang LY, Ramzan I. Toxicokinetics of kava. *Adv Pharmacol Sci*. 2011;2011:326724.21541070
13. Robinson V, Bergfeld WF, Belsito DV, et al. Final report on the safety assessment of *Piper methysticum* leaf/root/stem extract and *Piper methysticum* root extract. *Int J Toxicol*. 2009;28(6 suppl):175S-188S.19966149
14. Teschke R, Lebot V. Proposal for a kava quality standardization code. *Food Chem Toxicol*. 2011;49(10):2503-2516.21756963
15. Fetrow CW, Avila JR. *Professional's Handbook of Complementary & Alternative Medicines*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2004:472-477.
16. Freeman LW. *Mosby's Complementary & Alternative Medicine: A Research-Based Approach*. 3rd ed. St. Louis, MO: Mosby Elsevier; 2009:431-434.
17. Davies L, Drew C, Duffield P, Jamieson D. Effects of kava on benzodiazepine and GABA receptor binding. *Eur J Pharmacol*. 1990;183(2):558.
18. Kretzschmar R, Meyer HJ, Teschendorf HJ. Strychnine antagonistic potency of pyrone compounds of the kavaroot (*Piper methysticum* Forst.). *Experientia*. 1970;26(3):283-284.5417498
19. Backhauss C, Krieglstein J. Extract of kava (*Piper methysticum*) and its methysticin constituents protect brain tissue against ischemic damage in rodents. *Eur J Pharmacol*. 1992;215(2-3):265-269.1396990
20. Seitz U, Schüle A, Gleitz J. [3H]-Monoamine uptake inhibition properties of kava pyrones. *Planta Med*. 1997;63(6):548-549.9434608
21. Magura EI, Kopanitsa MV, Gleitz J, Peters T, Krishtal OA. Kava extract ingredients, (+)-methysticin and (±)-kavain inhibit voltage-operated Na(+)-channels in rat CA1 hippocampal neurons [published correction appears in *Neuroscience*. 1998;84(1):323]. *Neuroscience*. 1997;81(2):345-351.9300426
22. Friese J, Gleitz J. Kavain, dihydrokavain, and dihydromethysticin non-competitively inhibit the specific binding of [3H]-batrachotoxinin-A 20- α -benzoate to receptor site 2 of voltage-gated Na⁺ channels. *Planta Med*. 1998;64(5):458-459.9690349
23. Gleitz J, Gottner N, Ameri A, Peters T. Kavain inhibits non-stereospecifically veratridine-activated Na⁺ channels.

Planta Med. 1996;62(6):580-581.17252507

24. Gleitz J, Friese J, Beile A, Ameri A, Peters T. Anticonvulsive action of (\pm)-kavain estimated from its properties on stimulated synaptosomes and Na⁺ channel receptor sites. *Eur J Pharmacol.* 1996;315(1):89-97.8960869
25. Seitz U, Ameri A, Pelzer H, Gleitz J, Peters T. Relaxation of evoked contractile activity of isolated guinea-pig ileum by (\pm)-kavain. *Planta Med.* 1997;63(4):303-306.9270372
26. Boonen G, Pramanik A, Rigler R, Häberlein H. Evidence for specific interactions between kavain and human cortical neurons monitored by fluorescence correlation spectroscopy. *Planta Med.* 2000;66(1):7-10.10705725
27. Holm E, Staedt U, Heep J, et al. The action profile of D,L-kavain. Cerebral sites and sleep-wakefulness-rhythm in animals [in German]. *Arzneimittelforschung.* 1991;41(7):673-683.1772452
28. Sarris J, LaPorte E, Schweitzer I. Kava: a comprehensive review of efficacy, safety, and psychopharmacology. *Aust N Z J Psychiatry.* 2011;45(1):27-35.21073405
29. Pittler MH, Ernst E. Kava extract for treating anxiety. *Cochrane Database Syst Rev.* 2003;(1):CD003383.12535473
30. Ernst E. A re-evaluation of kava (*Piper methysticum*). *Br J Clin Pharmacol.* 2007;64(4):415-417.17555466
31. Witte S, Loew D, Gaus W. Meta-analysis of the efficacy of the acetonic kava-kava extract WS1490 in patients with non-psychotic anxiety disorders. *Phytother Res.* 2005;19(3):183-188.15934028
32. Lehrl S. Clinical efficacy of kava extract WS 1490 in sleep disturbances associated with anxiety disorders. Results of a multicenter, randomized, placebo-controlled, double-blind clinical trial [published correction appears in *J Affect Disord.* 2004;83(2-3):287]. *J Affect Disord.* 2004;78(2):101-110.14706720
33. Sarris J, Kavanagh DJ, Byrne G, Bone KM, Adams J, Deed G. The Kava Anxiety Depression Spectrum Study (KADSS): a randomized, placebo-controlled crossover trial using an aqueous extract of *Piper methysticum*. *Psychopharmacology (Berl).* 2009;205(3):399-407.19430766
34. De Leo V, la Marca A, Morgante G, Lanzetta D, Florio P, Petraglia F. Evaluation of combining kava extract with hormone replacement therapy in the treatment of postmenopausal anxiety. *Maturitas.* 2001;39(2):185-188.11514117
35. Warnecke G. Psychosomatic dysfunctions in the female climacteric. Clinical effectiveness and tolerance of Kava Extract WS 1490 [in German]. *Fortschr Med.* 1991;109(4):119-122.2029982
36. Warnecke G, Pfaender H, Gerster G, Gracza E. Wirksamkeit von kawa-kawa-extrakt beim klimakterischen syndrom. *Z Phytother.* 1990;11:81-86.
37. Cagnacci A, Arangino S, Renzi A, Zanni AL, Malmusi S, Volpe A. Kava-Kava administration reduces anxiety in perimenopausal women. *Maturitas.* 2003;44(2):103-109.12590005
38. Jacobs BP, Bent S, Tice JA, Blackwell T, Cummings SR. An internet-based randomized, placebo-controlled trial of kava and valerian for anxiety and insomnia. *Medicine (Baltimore).* 2005;84(4):197-207.16010204
39. Connor KM, Payne V, Davidson JR. Kava in generalized anxiety disorder: three placebo-controlled trials. *Int Clin Psychopharmacol.* 2006;21(5):249-253.16877894
40. Sarris J, Scholey A, Schweitzer I, et al. The acute effects of kava and oxazepam on anxiety, mood, neurocognition; and genetic correlates: a randomized, placebo-controlled, double-blind study. *Hum Psychopharmacol.* 2012;27(3):262-269.22311378
41. Triolet J, Shaik AA, Gallaher DD, O'Sullivan MG, Xing C. Reduction in colon cancer risk by consumption of kava or kava fractions in carcinogen-treated rats. *Nutr Cancer.* 2012;64(6):838-846.22693990
42. Eskander RN, Randall LM, Sakai T, Guo Y, Hoang B, Zi X. Flavokawain B, a novel, naturally occurring chalcone, exhibits robust apoptotic effects and induces G2/M arrest of a uterine leiomyosarcoma cell line. *J Obstet Gynaecol Res.* 2012;38(8):1086-1094.22540374
43. Johnson TE, Hermanson D, Wang L, et al. Lung tumorigenesis suppressing effects of a commercial kava extract and its selected compounds in A/J mice. *Am J Chin Med.* 2011;39(4):727-742.21721153
44. Sakai T, Eskander RN, Guo Y, et al. Flavokawain B, a kava chalcone, induces apoptosis in synovial sarcoma cell lines. *J Orthop Res.* 2012;30(7):1045-1050.22213202
45. Tang J, Dunlop RA, Rowe A, Rodgers KJ, Ramzan I. Kavalactones Yangonin and Methysticin induce apoptosis in human hepatocytes (HepG2) in vitro. *Phytother Res.* 2011;25(3):417-423.20734326
46. Tanaka A, Hamada N, Fujita Y, et al. A novel kavalactone derivative protects against H2O2-induced PC12 cell death

via Nrf2/ARE activation. *Bioorg Med Chem*. 2010;18(9):3133-3139.20371185

47. Warmka JK, Solberg EL, Zeliadt NA, et al. Inhibition of mitogen activated protein kinases increases the sensitivity of A549 lung cancer cells to the cytotoxicity induced by a kava chalcone analog. *Biochem Biophys Res Commun*. 2012;424(3):488-492.22771807
48. Cassileth B. Kava (*Piper methysticum*). *Oncology (Williston Park)*. 2011;25(4):384-385.21618963
49. Steiner GG. The correlation between cancer incidence and kava consumption. *Hawaii Med J*. 2000;59(11):420-422.11149250
50. LaPorte E, Sarris J, Stough C, Scholey A. Neurocognitive effects of kava (*Piper methysticum*): a systematic review. *Hum Psychopharmacol*. 2011;26(2):102-111.21437989
51. Sarris J, Kean J, Schweitzer I, Lake J. Complementary medicines (herbal and nutritional products) in the treatment of Attention Deficit Hyperactivity Disorder (ADHD): a systematic review of the evidence. *Complement Ther Med*. 2011;19(4):216-227.21827936
52. Gleitz J, Beile A, Wilkens P, Ameri A, Peters T. Anti-thrombotic action of the kava pyrone (+)-kavain prepared from *Piper methysticum* on human platelets. *Planta Med*. 1997;63(1):27-30.9063093
53. Locher CP, Burch MT, Mower HF, et al. Anti-microbial activity and anti-complement activity of extracts obtained from selected Hawaiian medicinal plants. *J Ethnopharmacol*. 1995;49(1):23-32.8786654
54. Otaguro K, Iwatsuki M, Ishiyama A, et al. In vitro antitrypanosomal activity of some phenolic compounds from propolis and lactones from Fijian Kava (*Piper methysticum*). *J Nat Med*. 2012;66(3):558-561.22116743
55. Jorm AF, Christensen H, Griffiths KM, Parslow RA, Rodgers B, Blewitt KA. Effectiveness of complementary and self-help treatments for anxiety disorders. *Med J Aust*. 2004;181(7 suppl):S29-S46.15462640
56. Feucht C, Patel DR. Herbal medicines in pediatric neuropsychiatry. *Pediatr Clin North Am*. 2011;58(1):33-54, x.2128184710.1016/j.pcl.2010.10.006
57. Ernst E. Herbal medicinal products during pregnancy: are they safe? *BJOG*. 2002;109(3):227-235.11950176
58. Pearl PL, Drillings IM, Conry JA. Herbs in epilepsy: evidence for efficacy, toxicity, and interactions. *Semin Pediatr Neurol*. 2011;18(3):203-208.22062945
59. Meyer, JJ, ed. Pharmacology of Kava. *Ethnopharmacologic Search for Psychoactive Drugs*. US Dept Health, Education, and Welfare, Pub No 1645. Washington, DC: Government Printing Office; 1967: 133-40.
60. Mathews JM, Etheridge AS, Black SR. Inhibition of human cytochrome P450 activities by kava extract and kavalactones. *Drug Metab Dispos*. 2002;30(11):1153-1157.12386118
61. Gurley BJ, Gardner SF, Hubbard MA, et al. In vivo effects of goldenseal, kava kava, black cohosh, and valerian on human cytochrome P450 1A2, 2D6, 2E1, and 3A4/5 phenotypes. *Clin Pharmacol Ther*. 2005;77(5):415-426.15900287
62. Gurley BJ, Swain A, Hubbard MA, et al. Supplementation with goldenseal (*Hydrastis canadensis*), but not kava kava (*Piper methysticum*), inhibits human CYP3A activity in vivo. *Clin Pharmacol Ther*. 2008;83(1):61-69.17495878
63. Shi S, Klotz U. Drug interactions with herbal medicines. *Clin Pharmacokinet*. 2012;51(2):77-104.22257149
64. Jamieson DD, Duffield PH. Positive interaction of ethanol and kava resin in mice. *Clin Exp Pharmacol Physiol*. 1990;17(7):509-514.2401104
65. Herberg KW. Effect of kava-special extract WS 1490 combined with ethyl alcohol on safety-relevant performance parameters. *Blutalkohol*. 1993;30(2):96-105.8481222
66. Schelosky L, Raffauf C, Jendroska K, Poewe W. Kava and dopamine antagonism. *J Neurol Neurosurg Psychiatry*. 1995;58(5):639-640.7745421
67. Almeida JC, Grimsley EW. Coma from the health food store: interaction between kava and alprazolam. *Ann Intern Med*. 1996;125(11):940-941.8967683
68. Miller LG. Herbal medicinals: selected clinical considerations focusing on known or potential drug-herb interactions. *Arch Intern Med*. 1998;158(20):2200-2211.9818800
69. Bressler R. Herb-drug interactions: interactions between kava and prescription medications. *Geriatrics*. 2005;60(9):24-25.16153141
70. Anke J, Ramzan I. Pharmacokinetic and pharmacodynamic drug interactions with Kava (*Piper methysticum* Forst.f.). *J Ethnopharmacol*. 2004;93(2-3):153-160.15234747

71. Ruze P. Kava-induced dermopathy: a niacin deficiency? *Lancet*. 1990;335(8703):1442-1445.1972218
72. Shulgin AT. The narcotic pepper—the chemistry and pharmacology of *Piper methysticum* and related species. *Bull Narc*. 1973;25(2):59-74.
73. Schwabe KP. Kava-kava extract, process for the production thereof and use thereof. 1994; US patent 5296224. March 22, 1994.
74. Leung N. Acute urinary retention secondary to kava ingestion. *Emerg Med Australas*. 2004;16(1):94.15239772
75. Ernst E. Safety concerns about kava. *Lancet*. 2002;359(9320):1865.12044412
76. Bodkin R, Schneider S, Rekkerth D, Spillane L, Kamali M. Rhabdomyolysis associated with kava ingestion. *Am J Emerg Med*. 2012;30(4):635.e1-e3.21458945
77. Ernst E. The risk-benefit profile of commonly used herbal therapies: Gingko, St. John's Wort, Ginseng, Echinacea, Saw Palmetto, and Kava [published correction appears in *Ann Intern Med*. 2003;138(1):79]. *Ann Intern Med*. 2002;136(1):42-53.11777363
78. Consumer advisory: kava-containing dietary supplements may be associated with severe liver injury. Food and Drug Administration Center for Food Safety and Applied Nutrition website. <http://www.fda.gov/Food/ResourcesForYou/Consumers/ucm085482.htm>. Published March 25, 2002. Accessed October 12, 2012.
79. From the Centers for Disease Control and Prevention. Hepatic toxicity possibly associated with kava-containing products—United States, Germany, and Switzerland, 1999-2002. *JAMA*. 2003;289(1):36-37.12515265
80. Connor KM, Davidson JR, Churchill LE. Adverse-effect profile of kava. *CNS Spectr*. 2001;6(10):848, 850-853.15334034
81. Teschke R, Wolff A. Kava hepatotoxicity: regulatory data selection and causality assessment. *Dig Liver Dis*. 2009;41(12):891-901.19477698
82. Olsen LR, Grillo MP, Skonberg C. Constituents in kava extracts potentially involved in hepatotoxicity: a review. *Chem Res Toxicol*. 2011;24(7):992-1002.21506562
83. Nerurkar PV, Dragull K, Tang CS. In vitro toxicity of kava alkaloid, pipermethystine, in HepG2 cells compared to kavalactones. *Toxicol Sci*. 2004;79(1):106-111.14737001
84. Anke J, Ramzan I. Kava Hepatotoxicity: Are we any closer to the truth? *Planta Med*. 2004;70(3):193-196.15114493
85. Zhang LY, Rowe A, Ramzan I. Does inflammation play a role in kava hepatotoxicity? *Phytother Res*. 2011;25(4):629-630.20842679
86. Yang X, Salminen WF. Kava extract, an herbal alternative for anxiety relief, potentiates acetaminophen-induced cytotoxicity in rat hepatic cells. *Phytomedicine*. 2011;18(7):592-600.21397479
87. Teschke R, Qiu SX, Xuan TD, Lebot V. Kava and Kava Hepatotoxicity: Requirements for Novel Experimental, Ethnobotanical and Clinical Studies Based on a Review of the Evidence. *Phytother Res*. 2011;25(9):1263-1274.21442674
88. Li Y, Mei H, Wu Q, et al. Methysticin and 7,8-dihydromethysticin are two major kavalactones in kava extract to induce CYP1A1. *Toxicol Sci*. 2011;124(2):388-399.21908763
89. Sarris J, Stough C, Bousman CA, et al. Kava in the treatment of generalized anxiety disorder: a double-blind, randomized, placebo-controlled study. *J Clin Psychopharmacol*. 2013;33(5):643-648.23635869
90. Ji T, Lin C, Krill LS, et al. Flavokawain B, a kava chalcone, inhibits growth of human osteosarcoma cells through G2/M cell cycle arrest and apoptosis. *Mol Cancer*. 2013;12:55.23764122
91. Sarris J, Stough C, Teschke R, et al. Kava for the treatment of generalized anxiety disorder RCT: analysis of adverse reactions, liver function, addiction, and sexual effects. *Phytother Res*. 2013;27(11):1723-1728.23348842
92. Ketola RA, Viinamäki J, Rasanen I, Pelander A, Goebeler S. Fatal kavalactone intoxication by suicidal intravenous injection. *Forensic Sci Int*. 2015;249:e7-e11.25684328
93. Navarro VJ, Barnhart H, Bonkovsky HL, et al. Liver injury from herbals and dietary supplements in the U.S. drug-induced liver injury network. *Hepatology*. 2014;60(4):1399-1408.25043597
94. Toohey TP, Lu BY, Wada C. Toxic effects of psychotropics related to possible p450 enzyme inhibition by kava: report of 2 cases. *Prim Care Companion CNS Disord*. 2013;15(5):PCC.13br01539.24511438

95. Perez J, Holmes JF. Altered mental status and ataxia secondary to acute kava ingestion. *J Emerg Med*. 2005;28(1):49-51.15657005
96. Gurley BJ, Gardner SF, Hubbard MA, et al. In vivo effects of goldenseal, kava kava, black cohosh, and valerian on human cytochrome P450 1A2, 2D6, 2E1, and 3A4/5 phenotypes. *Clin Pharmacol Ther*. 2005;77(5):415-426.15900287
97. Rubin D, McGovern B, Kopelman RI. Back to basics. *Am J Med*. 2006;119(6):482-483.16750959
98. Mysliwiec V, Martin JL, Ulmer CS, et al. The management of chronic insomnia disorder and obstructive sleep apnea: Synopsis of the 2019 U.S. Department of Veterans Affairs and U.S. Department of Defense Clinical Practice Guidelines. *Ann Intern Med*. 2020 Feb 18.
99. European Association for the Study of the Liver. Electronic address: easloffice@easloffice.eu; Clinical Practice Guideline Panel: Chair;; Panel members; EASL Governing Board representative:. EASL Clinical Practice Guidelines: Drug-induced liver injury. *J Hepatol*. 2019;70(6):1222-1261.30926241
100. Gurley BJ, Swain A, Hubbard MA, et al. Supplementation with goldenseal (*Hydrastis canadensis*), but not kava kava (*Piper methysticum*), inhibits human CYP3A activity in vivo. *Clin Pharmacol Ther*. 2008;83(1):61-69.17495878

Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright © 2025 Wolters Kluwer Health

DRUG STATUS

Availability

 Unknown

User Reviews & Ratings

7.9 / 10

[14 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)


[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.