

Home

2. Bentyl

Bentyl 🖘

Generic name: dicyclomine (oral/injection) [dye-SYE-kloe-meen]

Brand names: Bentyl, Dibent, Dicyclocot **Drug class:** Anticholinergics / antispasmodics

Medically reviewed by Sanjai Sinha, MD. Last updated on Nov 13, 2024.

Uses Warnings Before taking Dosage Side effects Interactions

What is Bentyl?

Bentyl is used to treat functional bowel or irritable bowel syndrome.

Bentyl may also be used for purposes not listed in this medication guide.

Warnings

Many drugs can affect Bentyl. Tell your doctor about all your current medicines.

Before taking this medicine

You should not use Bentyl if you are allergic to it, or if you have:

- glaucoma;
- a bladder obstruction or other urination problems;
- a blockage in your digestive tract (stomach or intestines);
- severe ulcerative colitis;
- gastroesophageal reflux disease (GERD);
- · a serious heart condition and active bleeding;
- · myasthenia gravis; or
- if you are breastfeeding a baby.

Not approved for use by anyone younger than 18 years old. **Bentyl should never be given to a child younger than 6** months old.

Tell your doctor if you have ever had:

- heart problems or high blood pressure;
- a stroke;
- ulcerative colitis;
- an ileostomy or colostomy;
- an enlarged prostate; or
- liver or kidney disease.

Older adults may be more sensitive to the effects of Bentyl.

Tell your doctor if you are pregnant.

Do not breastfeed.

Bentyl pregnancy and breastfeeding warnings (more detail)

How should I take Bentyl?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Bentyl oral is taken by mouth.

Measure liquid medicine with the supplied syringe or a dose-measuring device (not a kitchen spoon).

Bentyl injection is given in a muscle if you are unable to take the medicine by mouth.

Call your doctor if your symptoms do not improve after 2 weeks.

Store at room temperature away from moisture and heat.

Detailed Bentyl dosage information

What happens if I miss a dose?

Skip the missed dose and use your next dose at the regular time. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose can cause nausea, vomiting, dilated pupils, weakness or loss of movement in any part of your body, trouble swallowing, fainting, or seizure (convulsions).

What should I avoid while taking Bentyl?

May cause dizziness or blurred vision. Avoid driving or hazardous activity until you know how Bentyl will affect you.

Avoid becoming overheated or dehydrated during exercise and in hot weather. Bentyl can decrease sweating and you

may be more prone to heat stroke. Tell your doctor if you have a fever while taking this medicine.

Avoid using an antacid. Antacids can make it harder for your body to absorb Bentyl oral.

Bentyl side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- · fast or slow heartbeats, pounding heartbeats or fluttering in your chest;
- confusion, agitation, hallucinations, unusual thoughts or behavior;
- problems with memory or speech;
- problems with balance or muscle movement;
- diarrhea, severe constipation, or worsening of bowel symptoms;
- trouble swallowing;
- bruising, swelling, or pain where a Bentyl injection was given; or
- dehydration --dizziness, confusion, feeling very thirsty, less urination or sweating.

Confusion and mood or behavior changes may be more likely in older adults.

Common side effects may include:

- drowsiness, dizziness, weakness, nervousness;
- blurred vision;
- · dry mouth; or
- nausea.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i Bentyl side effects (more detail)

Related/similar drugs

Xifaxan

Xifaxan (rifaximin) is used to treat travelers' diarrhea, irritable bowel syndrome with diarrhea...

Reviews & ratings

6.5 / 10

95 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Levsin

Levsin is used to treat stomach problems such as cramps and irritable bowel syndrome and conditions ...

Reviews & ratings

6.2 / 10

16 Reviews

View more

Librax

Librax is used to treat stomach ulcers and irritable bowel syndrome. Learn about side effects ...

Reviews & ratings

8.6 / 10

72 Reviews

View more

Lotronex

Lotronex is used to treat severe, chronic irritable bowel syndrome (IBS) in women. Learn about side ...

Reviews & ratings

9.6 / 10

24 Reviews

View more

What other drugs will affect Bentyl?

Using Bentyl with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your current medicines. Many drugs can affect Bentyl, especially:

- bronchodilator asthma medication;
- cold or allergy medicine (Benadryl and others);
- glaucoma medication;
- · heart medication;
- medicine to treat depression, anxiety, mood disorders, or mental illness;
- medicine to treat overactive bladder;
- medicine to treat Parkinson's disease; or
- medicine to treat stomach problems, motion sickness, or irritable bowel syndrome.

This list is not complete and many other drugs may affect Bentyl. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Bentyl drug interactions (more detail)

Does Bentyl interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Bentyl	
+	
Enter a drug name	Add

More about Bentyl (dicyclomine)

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (138)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Support group

- Drug class: anticholinergics/antispasmodics
- Breastfeeding
- En español

Professional resources

- · Bentyl prescribing information
- Dicyclomine (AHFS Monograph)

Related treatment guides

• Irritable Bowel Syndrome

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ঝ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

The Drug history at FDA

User Reviews & Ratings

7.3 / 10

138 Reviews

Images

Bentyl 20 mg (BENTYL 20)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.