

[Home](#)

2. Lurasidone

# Lurasidone

**Generic name:** lurasidone [ *loo-RAS-i-done* ]**Brand name:** [Latuda](#)**Dosage form:** oral tablet (120 mg; 20 mg; 40 mg; 60 mg; 80 mg)**Drug class:** [Atypical antipsychotics](#)[Medically reviewed](#) by Drugs.com on Jan 27, 2025. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

## What is lurasidone?

Lurasidone is an antipsychotic medicine that is used to treat [schizophrenia](#) in adults and teenagers who are at least 13 years old.

Lurasidone is also used to treat episodes of depression related to [bipolar disorder](#) ([manic depression](#)) in adults and children who are at least 10 years old.

Lurasidone may also be used for purposes not listed in this medication guide.

## Lurasidone side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

**Report any new or worsening symptoms to your doctor**, such as: mood or behavior changes, anxiety, [panic attacks](#), trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

**High doses or long-term use of lurasidone can cause a serious movement disorder that may not be reversible.**

The longer you use lurasidone, the more likely you are to develop this disorder, especially if you are a woman or an older adult.

**Lurasidone may cause serious side effects. Call your doctor at once if you have:**

- any new or unusual muscle movements you cannot control;
- a light-headed feeling, like you might pass out;
- a seizure (convulsions);
- (in women) irregular menstrual periods, breast or vaginal changes, nipple discharge;

- (in men) breast swelling, impotence;
- trouble swallowing;
- **manic episodes**--racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative;
- **low white blood cell counts**--fever, chills, mouth sores, skin sores, sore throat, cough, trouble breathing;
- **high blood sugar**--increased thirst, increased urination, hunger, dry mouth, fruity breath odor; or
- **severe nervous system reaction**--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out.

**Common side effects of lurasidone may include:**

- drowsiness;
- weight gain;
- tremors, muscle stiffness, slow muscle movement;
- feeling restless or being unable to sit still;
- nausea, vomiting;
- runny nose; or
- sleep problems (insomnia).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Lurasidone side effects](#) (more detail)

## Related/similar drugs

### Lybalvi

Lybalvi is used to treat adults with manic or mixed episodes associated with bipolar I disorder ...

#### Reviews & ratings

8.1 / 10

[16 Reviews](#)

[View more](#)

### Fanapt

Fanapt (iloperidone) is an antipsychotic medication used to treat schizophrenia. Includes Fanapt ...

#### Reviews & ratings

6.4 / 10

[30 Reviews](#)

[View more](#)

#### FEATURED

### Caplyta

Caplyta (lumateperone) may be used to treat schizophrenia or depression associated with bipolar ...

#### Reviews & ratings

6.6 / 10

[184 Reviews](#)

[View more](#)

### Abilify

Abilify is an antipsychotic medicine used to treat the symptoms of schizophrenia and bipolar ...

#### Reviews & ratings

6.1 / 10

[1,187 Reviews](#)

[View more](#)

### Olanzapine

Olanzapine (brand name Zyprexa) is an atypical antipsychotic that may be used to treat adults and ...

#### Reviews & ratings

5.9 / 10

[1,091 Reviews](#)

[View more](#)

### Aripiprazole

Easy-to-read patient leaflet for aripiprazole. Includes indications, proper use, special ...

#### Reviews & ratings

6.1 / 10

[1,540 Reviews](#)

[View more](#)

## Quetiapine

Quetiapine is an atypical antipsychotic used to treat schizophrenia, bipolar disorder, and major ...

### Reviews & ratings

7.2 / 10

[2,249 Reviews](#)

[View more](#)

## Seroquel

Seroquel is used to treat the symptoms of schizophrenia, bipolar disorder and major depressive ...

### Reviews & ratings

7.0 / 10

[694 Reviews](#)

[View more](#)

## Lamictal

Lamictal is an anti-epileptic medication used treat seizures in adults and children over 2 years ...

### Reviews & ratings

8.0 / 10

[792 Reviews](#)

[View more](#)

## Warnings

Lurasidone is not approved for use in older adults with dementia-related psychosis.

Some people have thoughts about suicide while taking lurasidone. Stay alert to changes in your mood or symptoms.

**Report any new or worsening symptoms to your doctor.**

Tell your doctor about all your current medicines and any you start or stop using. **Many drugs can interact, and some drugs should not be used together.**

## Before taking this medicine

You should not use lurasidone if you are allergic to it.

**Many drugs can interact and cause dangerous effects. Some drugs should not be used together with lurasidone.** Your doctor may change your treatment plan if you also use:

- antifungal medicine such as [ketoconazole](#) or [voriconazole](#);
- an antibiotic such as [clarithromycin](#) or rifampin;
- an antiviral such as ritonavir;
- [St. John's wort](#); or
- seizure medicine such as [carbamazepine](#) or phenytoin.

Lurasidone may increase the risk of death in older adults with dementia-related psychosis and is not approved for this use.

Tell your doctor if you have ever had:

- [heart disease](#) or a stroke;
- high or [low blood pressure](#);
- [high cholesterol](#) or triglycerides (a type of fat in the blood);
- diabetes or [high blood sugar](#) (in you or your family);
- a seizure;
- liver or kidney disease;
- low white blood cell (WBC) counts;
- abnormal hormone function tests (thyroid, pituitary gland);
- [breast cancer](#); or
- suicidal thoughts or actions.

Some people have thoughts about suicide while taking lurasidone. Your doctor will need to check your progress at regular visits. Your family or other caregivers should also be alert to changes in your mood or symptoms.

**Taking antipsychotic medicine in the last 3 months of pregnancy may cause breathing problems, feeding problems, or withdrawal symptoms in the newborn.** If you get pregnant, tell your doctor right away. **Do not stop taking lurasidone without your doctor's advice.**

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of lurasidone on the baby.

It may not be safe to breastfeed a baby while you are using lurasidone. Ask your doctor about any risks.

Lurasidone is not approved **for schizophrenia** in anyone younger than 13 years old. Lurasidone is not approved **for depression** in anyone younger than 10 years old.

## How should I take lurasidone?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Lurasidone should be taken with food (at least 350 calories).

You may need frequent blood tests.

It may take several weeks before your symptoms improve. Keep using the medication as directed. Call your doctor if your symptoms do not improve, or if they get worse while using lurasidone.

**You should not stop using lurasidone suddenly.** Stopping suddenly may cause other problems.

It is easier to become dangerously overheated and dehydrated while you are taking lurasidone. **Drink plenty of fluids**, especially in hot weather and during exercise. You may also be more sensitive to temperature extremes (hot or cold).

Store at room temperature away from moisture and heat.

 [Detailed Lurasidone dosage information](#)

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

Get your prescription refilled before you run out of medicine completely.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

## What should I avoid while taking lurasidone?

**Avoid drinking alcohol.** Dangerous side effects could occur.

Avoid driving or operating machinery until you know how lurasidone will affect you. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Grapefruit and [grapefruit juice](#) may interact with lurasidone and lead to unwanted side effects. **Avoid the use of grapefruit products while taking lurasidone.**

## What other drugs will affect lurasidone?

**Using lurasidone with other drugs that slow your breathing can cause dangerous side effects or death.** Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

**Tell your doctor about all your current medicines.** Many drugs can affect lurasidone, especially:

- depression or psychotic episodes;
- sleep problems (insomnia);

- [high blood pressure](#) or a heart rhythm disorder;
- swelling or inflammation;
- seizures; or
- [Parkinson's disease](#).

**This list is not complete and many other drugs may affect lurasidone.** This includes prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

 [Lurasidone drug interactions](#) (more detail)

## Does lurasidone interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).








lurasidone

+

Enter a drug name

Add

## Popular FAQ

- Can you stop taking Latuda immediately? 
- Does Latuda make you sleepy? 
- Is Latuda a mood stabilizer or an antipsychotic? 
- How fast does Latuda work? 
- Does Latuda cause weight gain? 
- Can Latuda be cut in half or split? 
- Is Latuda a controlled substance? 

## More FAQ

- [What drugs cause tardive dyskinesia?](#)

[View more FAQ...](#)

## More about lurasidone

- [Check interactions](#)

- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(951\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: atypical antipsychotics](#)
- [Breastfeeding](#)
- [En español](#)

## Patient resources

- [Lurasidone advanced reading](#)

## Other brands

[Latuda](#)

## Professional resources

- [Lurasidone monograph](#)
- [Lurasidone Tablets \(FDA\)](#)

## Other brands

[Latuda](#)

## Related treatment guides

- [Borderline Personality Disorder](#)
- [Schizoaffective Disorder](#)
- [Bipolar Disorder](#)
- [Schizophrenia](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.


[Medical Disclaimer](#)




DRUG STATUS

- Availability

Rx Prescription only
- Pregnancy & Lactation

 Risk data available
- CSA Schedule\*

N/A Not a controlled drug
- Approval History

 Drug history at FDA



User Reviews & Ratings

6.3 / 10

[951 Reviews](#)

Images

[Lurasidone Hydrochloride 40 mg \(A 40\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

- [About Drugs.com](#)
- [Advertising policy](#)
- [Attribution & citations](#)

Terms & privacy

- [Terms of use](#)
- [Editorial policy](#)
- [Privacy policy](#)

Support

- [Help center](#)
- [Sitemap](#)

Contact us



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.