

Home

2. Levora

Levora 🖘

Generic name: ethinyl estradiol and levonorgestrel [ETH-in-ill-ess-tra-DYE-ol-and-LEE-vo-nor-JESS-trel]

Brand names: Afirmelle, Altavera, Amethyst, Aubra EQ, Aviane, ... show all 24 brands

Drug class: Contraceptives



Medically reviewed by Drugs.com on Mar 31, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Levora?

Levora is used as contraception to prevent pregnancy. There are many brands and forms of Levora available. Not all brands are listed on this leaflet.

Levora may also be used for purposes not listed in this medication guide.

Levora side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have:

- signs of a stroke--sudden numbness or weakness (especially on one side of the body), severe headache, slurred speech, balance problems;
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, swelling or redness in an arm or leg;
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, sweating;
- liver problems--loss of appetite, upper stomach pain, tiredness, fever, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears;
- gallbladder problem--chalky-colored stools, stomach pain after eating, nausea, heartburn, bloating, and severe upper stomach pain that may spread to your back;
- swelling in your hands, ankles, or feet;
- changes in the pattern or severity of migraine headaches;

- a breast lump; or
- symptoms of depression--sleep problems, weakness, tired feeling, mood changes.

Common side effects of Levora may include:

- nausea, vomiting (especially when you first start taking this medicine);
- · breast tenderness;
- · breakthrough bleeding;
- · acne, darkening of facial skin;
- · weight gain; or
- problems with contact lenses.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i Levora side effects (more detail)

Related/similar drugs

Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

Reviews & ratings

6.1 / 10

2,624 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Depo Provera

Depo-Provera is used as contraception to prevent pregnancy. It is also used to reduce pain cause by

Reviews & ratings

4.2 / 10

1,967 Reviews

View more

Nexplanon

Nexplanon is a long-term, reversible birth control implant that lasts for three years. Learn about ...

Reviews & ratings

5.0 / 10

4,864 Reviews

View more

Yaz

Yaz (drospirenone and ethinyl estradiol) is used as a contraceptive to prevent pregnancy. Includes ...

Reviews & ratings

5.9 / 10

997 Reviews

View more

Slynd

Slynd (drospirenone) is a birth control pill used for the prevention of pregnancy. Includes Slynd ...

Reviews & ratings

5.8 / 10

386 Reviews

View more

Warnings

Do not use birth control pills if you are pregnant or if you have recently had a baby.

You should not use birth control pills if you have: uncontrolled high blood pressure, heart disease, coronary artery

disease, circulation problems (especially with diabetes), undiagnosed vaginal bleeding, liver disease or liver cancer, severe migraine headaches, if you also take certain hepatitis C medication, if you will have major surgery, if you smoke and are over 35, or if you have ever had a heart attack, a stroke, a blood clot, jaundice caused by pregnancy or birth control pills, or cancer of the breast, uterus/cervix, or vagina.

Taking birth control pills can increase your risk of blood clots, stroke, or heart attack.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. You should not take Levora if you smoke and are over 35 years old.

Before taking this medicine

Taking this medicine can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, high cholesterol, or if you are overweight. Your risk of stroke or blood clot is highest during your first year of taking birth control pills. Your risk is also high when you restart this medicine after not taking it for 4 weeks or longer.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. Your risk increases the older you are and the more you smoke. You should not take combination birth control pills if you smoke and are over 35 years old.

Do not use if you are pregnant. Stop using this medicine and tell your doctor right away if you become pregnant, or if you miss two menstrual periods in a row. If you have recently had a baby, wait at least 4 weeks before taking birth control pills.

You should not take birth control pills if you have:

- untreated or uncontrolled high blood pressure;
- heart disease (chest pain, coronary artery disease, history of heart attack, stroke, or blood clot);
- an increased risk of having blood clots due to a heart problem or a hereditary blood disorder;
- circulation problems (especially if caused by diabetes);
- a history of hormone-related cancer, or cancer of the breast, uterus/cervix, or vagina;
- unusual vaginal bleeding that has not been checked by a doctor;
- · liver disease or liver cancer;
- severe migraine headaches (with aura, numbness, weakness, or vision changes), especially if you are older than 35;
- a history of jaundice caused by pregnancy or birth control pills; or
- if you take any hepatitis C medication containing ombitasvir/paritaprevir/ritonavir (Technivie).

Tell your doctor if you have ever had:

- heart disease, high blood pressure, or if you are prone to having blood clots;
- high cholesterol or triglycerides, or if you are overweight;
- · depression;
- a seizure or migraine headache;
- diabetes, gallbladder disease;

- liver or kidney disease;
- · irregular menstrual cycles; or
- fibrocystic breast disease, lumps, nodules, or an abnormal mammogram.

Levora can slow breast milk production. You should not breastfeed while using this medicine.

How should I take Levora?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Take your first pill on the first day of your period or on the first Sunday after your period begins. You may need to use back-up birth control, such as condoms or a spermicide, when you first start using Levora. Follow your doctor's instructions.

Take one pill every day, no more than 24 hours apart. When the pills run out, start a new pack the following day. **You** may get pregnant if you do not take one pill daily. Get your prescription refilled before you run out of pills completely.

Some birth control packs contain seven "reminder" pills to keep you on your regular cycle. Your period will usually begin while you are using these reminder pills.

You should chew the **chewable tablet** (pill) and then swallow it with a glass of water, or if you prefer you should swallow the **chewable tablet** (pill) whole. Take it on an empty stomach.

Use a back-up birth control if you are sick with severe vomiting or diarrhea.

You may have breakthrough bleeding, especially during the first 3 months. Tell your doctor if this bleeding continues or is very heavy.

If you need major surgery or will be on long-term bed rest, you may need to stop using this medicine for a short time. Any doctor or surgeon who treats you should know that you are using birth control pills.

While taking birth control pills, you will need to visit your doctor regularly.

Store at room temperature away from moisture, heat, and light.

Detailed Levora dosage information

What happens if I miss a dose?

Follow the patient instructions provided with your medicine. Missing a pill increases your risk of becoming pregnant.

If you miss one active pill, take two pills on the day that you remember. Then take one pill per day for the rest of the pack.

If you miss two active pills in a row in Week 1 or 2, take two pills per day for two days in a row. Then take one pill per day for the rest of the pack. Use back-up birth control for at least 7 days following the missed pills.

If you miss two active pills in a row in Week 3, throw out the rest of the pack and start a new pack the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest

of the pack and start a new pack that day.

If you miss three active pills in a row in Week 1, 2, or 3, throw out the rest of the pack and start a new pack on the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss two or more active pills, you may not have a period during the month. If you miss a period for two months in a row, call your doctor because you might be pregnant.

If you miss a reminder pill, throw it away and keep taking one reminder pill per day until the pack is empty.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose symptoms may include nausea, vomiting, and drowsiness.

What should I avoid while taking Levora?

Do not smoke while taking birth control pills, especially if you are older than 35 years of age.

Birth control pills will not protect you from sexually transmitted diseases--including HIV and AIDS. Using a condom is the only way to protect yourself from these diseases.

What other drugs will affect Levora?

Other drugs may interact with birth control pills, including prescription and over-the-counter medicines, vitamins, and herbal products. Some drugs can make birth control pills less effective, which may result in pregnancy. Tell your doctor about all your current medicines and any medicine you start or stop using.

<u>Levora drug interactions</u> (more detail)

Does Levora interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



More about Levora (ethinyl estradiol / levonorgestrel)

- · Check interactions
- · Compare alternatives
- · Pricing & coupons

- Reviews (231)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: contraceptives
- En español

Patient resources

Other brands

Aviane, Vienva, Jolessa, Altavera, ... +44 more

Professional resources

· Levora prescribing information

Other brands

Aviane, Vienva, Jolessa, Altavera, ... +49 more

Related treatment guides

- Endometriosis
- Abnormal Uterine Bleeding
- Birth Control
- Gonadotropin Inhibition
- Ovarian Cysts
- Polycystic Ovary Syndrome

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 17.03.

Availability

Rx Prescription only

Pregnancy & Lactation

ঝ্য Risk data available

CSA Schedule*

N/A Not a controlled drug

User Reviews & Ratings

5.9 / 10

231 Reviews

Related News

Certain Birth Control Pills, Devices Linked to Higher Heart Risks

Images

Levora inert (WATSON P1)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us







@

Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.