

Home

Treprostinil

Treprostinil

Generic name: treprostinil (oral) [tre-PROS-ti-nil]

Brand names: Orenitram, Orenitram Titration Kit Month 1, Orenitram Titration Kit Month 2, Orenitram Titration Kit Month 3 **Dosage form:** oral tablet, extended release (0.125 mg; 0.125 mg-0.25 mg; 0.125 mg-0.25 mg; 0.25 mg; 1 mg; 2.5 mg; 5 mg)

Drug class: Agents for pulmonary hypertension



Medically reviewed by Drugs.com on Jul 10, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is treprostinil?

Treprostinil is used in adults to treat high blood pressure in the arteries of the lungs also known as pulmonary arterial hypertension (PAH). treprostinil improves your ability to exercise and can help prevent your condition from getting worse.

Treprostinil may also be used for purposes not listed in this medication guide.

Treprostinil side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

Treprostinil may cause serious side effects. Call your doctor at once if you have:

- · chest pain, trouble breathing, tiredness, pale skin, swelling in arms or legs; or
- a light-headed feeling, like you might pass out.

Common side effects of treprostinil may include:

- headache:
- · stomach pain;
- jaw pain, pain in your arms or legs;
- · diarrhea, nausea, vomiting; or
- flushing (sudden warmth, redness, or tingly feeling).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Treprostinil side effects (more detail)

Related/similar drugs

Cialis

Cialis (tadalafil) increase increases blood flow to a certain area of the body and is used to treat ...

Reviews & ratings

8.6 / 10

554 Reviews

View more

Viagra

Viagra (sildenafil) is used to treat erectile dysfunction or impotence in men. Includes Viagra side ...

Reviews & ratings

8.5 / 10

253 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Adcirca

Addirca is used to treat pulmonary arterial hypertension. Learn about side effects, interactions ...

Reviews & ratings

6 Reviews

View more

Opsumit

Opsumit (macitentan) is used to treat

pulliforiary arterial riyperterision. Includes Opsumit side ...

Reviews & ratings

8.2 / 10

10 Reviews

View more

Revatio

Revatio (sildenafil) is used to treat pulmonary arterial hypertension. Includes Revatio side ...

Reviews & ratings

6.9 / 10

10 Reviews

View more

Adempas

Adempas (riociguat) is used to treat pulmonary hypertension. Includes Adempas side effects ...

Reviews & ratings

4 Reviews

View more

Warnings

You should not use treprostinil if you have severe liver disease.

You should not stop taking treprostinil suddenly. Your condition may get worse if you stop taking treprostinil.

Before taking this medicine

You should not use treprostinil if you have severe liver disease.

Tell your doctor if you have or have ever had:

- diverticulosis (an intestinal disorder); or
- liver disease.

It is not known if treprostinil will harm an unborn baby. However, having high lung blood pressure during pregnancy may cause complications or medical problems in both mother and baby. The benefit of treating pulmonary hypertension may

outweigh any risks to the baby.

Ask a doctor if it is safe to breastfeed while using treprostinil.

How should I take treprostinil?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Take with food.

Swallow the tablet whole and do not crush, chew, or break it. Do not use a cracked or broken tablet.

Treprostinil is usually taken 2 to 3 times per day. Follow your doctor's instructions. Your dose may change if you have side effects.

Use treprostinil regularly for the best results.

Your dose needs may change if you switch to a different brand, strength, or form of this medicine. **Avoid medication** errors by using only the medicine your doctor prescribes.

You should not stop taking treprostinil suddenly. Your condition may get worse if you stop taking treprostinil. Follow your doctor's instructions about tapering your dose.

Part of a tablet shell may appear in your stool but this will not make the medicine less effective.

Store at room temperature away from moisture and heat.

Detailed Treprostinil dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

Call your doctor for instructions if you miss 2 or more doses of treprostinil.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking treprostinil?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

What other drugs will affect treprostinil?

Tell your doctor about all your other medicines, especially:

· gemfibrozil; or

• medicine that contains treprostinil--Remodulin, Tyvaso, or Tyvaso DPI.

This list is not complete. Other drugs may affect treprostinil, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Treprostinil drug interactions (more detail)

Does treprostinil interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

treprostinil	
+	
Enter a drug name	Add

More about treprostinil

- · Check interactions
- · Compare alternatives
- · Pricing & coupons
- Reviews (25)
- Latest FDA alerts (1)
- Side effects
- Dosage information
- During pregnancy
- Drug class: agents for pulmonary hypertension
- Breastfeeding

Patient resources

- Treprostinil inhalation drug information
- Treprostinil injection
- Treprostinil (Inhalation) (Advanced Reading)
- Treprostinil (Injection) (Advanced Reading)
- Treprostinil (Oral) (Advanced Reading)

Other brands

Remodulin, Tyvaso, Tyvaso DPI, Orenitram

Professional resources

- Treprostinil monograph
- Treprostinil Injection (FDA)

Other brands

Remodulin, Tyvaso, Tyvaso DPI, Orenitram

Related treatment guides

• Pulmonary Arterial Hypertension

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 3.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ം Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

7.0 / 10

25 Reviews

Images

Orenitram (treprostinil) 1 mg (UT 1)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





