

Home

2. Doxepin

Doxepin (capsules, oral concentrate) ⊲ ¬¬

Generic name: doxepin (capsules, oral concentrate) [DOX-e-pin]

Dosage forms: oral capsule (10 mg; 100 mg; 150 mg; 25 mg; 50 mg; 75 mg), oral concentrate (10 mg/mL)

Drug classes: Miscellaneous anxiolytics, sedatives and hypnotics, Tricyclic antidepressants



Medically reviewed by Drugs.com on Jun 3, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is doxepin?

Doxepin is a tricyclic antidepressant.

Doxepin **capsules** and **oral concentrate** (**liquid**) are used to treat symptoms of depression and/or anxiety associated with alcoholism, manic depression, or other mental illness.

Doxepin tablets (Silenor) are used to treat insomnia in people who have trouble staying asleep.

This medication guide provides only information about doxepin capsules and liquid.

Doxepin may also be used for purposes not listed in this medication guide.

Doxepin side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Doxepin may cause serious side effects. Call your doctor at once if you have:

- blurred vision, tunnel vision, eye pain or swelling, seeing halos around lights;
- a light-headed feeling, like you might pass out;
- tremors, restless muscle movements in your eyes, tongue, jaw, or neck;
- severe constipation;
- · confusion, hallucinations, unusual thoughts, seizure (convulsions); or
- · painful or difficult urination.

Common side effects of doxepin may include:

- drowsiness, dizziness;
- increased blood pressure;
- · vision changes;
- · rash, swelling;
- nausea, vomiting, indigestion;
- little or no urination;
- dry mouth, constipation; or
- · decreased or increased sex drive.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

① Doxepin side effects (more detail)

Related/similar drugs

Lexapro

Lexapro is used to treat anxiety and major depressive disorder. Learn about side effects ...

Reviews & ratings

7.6 / 10

2,376 Reviews

View more

Citalopram

Citalopram is an antidepressant (selective serotonin reuptake inhibitor) and is used to treat ...

Reviews & ratings

7.4 / 10

2,231 Reviews

View more

FEATURED

Xolair

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

Reviews & ratings

7.5 / 10

229 Reviews

View more

Loratadine

Loratadine is a non-drowsy antihistamine used to relieve allergy symptoms like sneezing, runny ...

Reviews & ratings

4.7 / 10

139 Reviews

View more

Cetirizine

Cetirizine treats cold or allergy symptoms such as sneezing, itching, watery eyes, or runny nose ...

Reviews & ratings

5.8 / 10

294 Reviews

View more

Venlafaxine

Venlafaxine is used to treat major depressive disorder, anxiety, and panic disorder. Learn about ...

Reviews & ratings

6.5 / 10

2,786 Reviews

View more

Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

Reviews & ratings

7.9 / 10

View more

Alprazolam

Alprazolam is used to treat anxiety disorders, panic disorders, and anxiety. Learn about side ...

Reviews & ratings

8.6 / 10

1,486 Reviews

View more

Buspirone

Buspirone is used to treat symptoms of anxiety, such as fear, tension, irritability and dizziness ...

Reviews & ratings

5.9 / 10

1,357 Reviews

View more

Escitalopram

Escitalopram is an SSRI antidepressant used to treat major depressive disorder in adults and teens ...

Reviews & ratings

7.5 / 10

3,962 Reviews

View more

Warnings

You should not take doxepin if you have glaucoma or problems with urination.

Do not use doxepin if you have used an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine.

Some young people have thoughts about suicide when first taking an antidepressant. Stay alert to changes in your mood

or symptoms. Report any new or worsening symptoms to your doctor.

Before taking this medicine

You should not use doxepin capsules or liquid if you are allergic to doxepin or similar medicines such as amoxapine or loxapine, or if you have:

- · glaucoma; or
- · urination problems.

Do not use doxepin if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others.

Tell your doctor if you have used an "SSRI" antidepressant in the past 5 weeks, such as citalopram, escitalopram, fluoxetine (Prozac), fluoxamine, paroxetine, sertraline (Zoloft), trazodone, or vilazodone.

Tell your doctor if you have ever had:

• bipolar disorder (manic-depression).

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor should check your progress at regular visits. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Tell your doctor if you are pregnant or breastfeeding.

Doxepin is not approved for use by anyone younger than 12 years old.

How should I take doxepin?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Measure **liquid medicine** carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

Mix the measured dose with 4 ounces of water, milk, orange juice, grapefruit juice, tomato juice, prune juice, or pineapple juice. Do not mix with grape juice or a carbonated soft drink. Stir the mixture and drink all of it right away. Do not save it for later use.

Do not stop using doxepin suddenly, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using doxepin.

It may take up to 3 weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Store at room temperature away from moisture and heat. Keep the bottle tightly closed when not in use.

Detailed Doxepin dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of doxepin can be fatal.**

Overdose symptoms may include irregular heart rhythm, feeling like you might pass out, seizures, or coma.

What should I avoid while taking doxepin?

Do not drink alcohol. Dangerous side effects or death could occur.

Avoid driving or hazardous activity until you know how doxepin will affect you. Your reactions could be impaired.

What other drugs will affect doxepin?

Using doxepin with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Other drugs may affect doxepin, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

1 Doxepin drug interactions (more detail)

Does doxepin interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

What are the strongest sleeping pills?

More FAQ

What are some common side effects of antidepressants?

.

More about doxepin

- Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (433)
- Drug images
- · Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: miscellaneous anxiolytics, sedatives and hypnotics
- Breastfeeding
- En español

Patient resources

- Doxepin (Oral) advanced reading
- Doxepin Tablets
- Doxepin Capsules
- Doxepin Oral Concentrate

Other brands

Sinequan, Silenor

Professional resources

- Doxepin monograph
- Doxepin (FDA)
- Doxepin Capsules (FDA)
- Doxepin Oral Solution Concentrate (FDA)
- Doxepin Tablets (FDA)

Other brands

Sinequan, Silenor

Related treatment guides

- Insomnia
- Anxiety

- Irritable Bowel Syndrome
- Depression

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 14.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ঝ্য Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

6.2 / 10

433 Reviews

Images

Doxepin Hydrochloride 10 mg (AMNEAL 1166)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×