

Home

2. Desipramine

Desipramine 49

Generic name: desipramine [des-IP-ra-meen]

Brand name: Norpramin

Dosage form: oral tablet (10 mg; 100 mg; 150 mg; 25 mg; 50 mg; 75 mg)

Drug class: Tricyclic antidepressants



Medically reviewed by Drugs.com on Jun 12, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is desipramine?

Desipramine is a tricyclic antidepressant that is used to treat symptoms of depression.

Desipramine may also be used for purposes not listed in this medication guide.

Desipramine side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Desipramine may cause serious side effects. Call your doctor at once if you have:

- · easy bruising, unusual bleeding, purple or red spots under your skin;
- a light-headed feeling, like you might pass out;
- confusion, hallucinations, unusual thoughts or behavior;
- blurred vision, tunnel vision, eye pain or swelling, seeing halos around lights;
- new or worsening chest pain, pounding heartbeats or fluttering in your chest;
- sudden numbness or weakness, problems with vision, speech, or balance;
- fever, sore throat;
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum);
- · painful or difficult urination;

- upper stomach pain, loss of appetite, jaundice (yellowing of the skin or eyes);
- · a seizure; or
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Serious side effects may be more likely in older adults and those who are ill or debilitated.

Common side effects of desipramine may include:

- · increased blood pressure;
- tingly feeling, weakness, lack of coordination;
- dry mouth, loss of appetite, nausea, vomiting;
- breast swelling (in men or women); or
- · weight gain or weight loss.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Desipramine side effects (more detail)

Related/similar drugs

Lexapro

Lexapro is used to treat anxiety and major depressive disorder. Learn about side effects ...

Reviews & ratings

7.6 / 10

2,376 Reviews

View more

Zoloft

Zoloft is an antidepressant used to treat major depression, obsessivecompulsive disorder, panic ...

Reviews & ratings

7.3 / 10

1,972 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Seroquel

Seroquel is used to treat the symptoms of schizophrenia, bipolar disorder and major depressive ...

Reviews & ratings

7.0 / 10

694 Reviews

View more

Prozac

Prozac (fluoxetine) is an SSRI antidepressant used to treat depression, OCD, panic disorder ...

Reviews & ratings

7.2 / 10

1,239 Reviews

View more

Cymbalta

Cymbalta (duloxetine) is used to treat major depressive disorder, general anxiety disorder and ...

Reviews & ratings

6.3 / 10

1,899 Reviews

View more

Wellbutrin

Wellbutrin (bupropion) is used to treat major depressive disorder and seasonal affective disorder ...

Reviews & ratings

6.8 / 10

756 Reviews

View more

Celexa

Celexa (citalopram) is an antidepressant in a group of drugs called selective serotonin reuptake ...

Reviews & ratings

7.3 / 10

866 Reviews

View more

Concerta

Concerta is used to treat attention deficit disorder (ADD) and attention deficit hyperactivity ...

Reviews & ratings

6.9 / 10

424 Reviews

View more

Citalopram

Citalopram is an antidepressant (selective serotonin reuptake inhibitor) and is used to treat ...

Reviews & ratings

7.4 / 10

2,231 Reviews

View more

Warnings

You should not use desipramine if you have recently had a heart attack.

Do not use this medicine within 14 days before or after taking an MAO inhibitor, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine.

Some young people have thoughts about suicide when first taking an antidepressant. Stay alert to changes in your mood or symptoms. Report any new or worsening symptoms to your doctor.

Before taking this medicine

You should not use desipramine if you are allergic to it, or:

· if you have recently had a heart attack.

Do not use desipramine if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others. Do not take an MAOI for at least 2 weeks after you stop taking desipramine.

Tell your doctor if you have used an "SSRI" antidepressant in the past 5 weeks, such as citalopram, escitalopram, fluoxetine (Prozac), fluoxamine, paroxetine, sertraline (Zoloft), trazodone, or vilazodone.

Tell your doctor if you have ever had:

- bipolar disorder (manic-depression) or schizophrenia;
- mental illness or psychosis;
- liver disease:
- · heart disease;
- a family history of sudden death related to a heart rhythm disorder;
- a heart attack, stroke, or seizures;
- · a thyroid disorder;
- · glaucoma; or
- problems with urination.

Be sure your doctor knows if you also take stimulant medicine, opioid medicine, herbal products, or medicine for depression, mental illness, Parkinson's disease, migraine headaches, serious infections, or prevention of nausea and vomiting. These medicines may interact with desipramine and cause a serious condition called serotonin syndrome.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor should check your progress at regular visits. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Tell your doctor if you are pregnant or breastfeeding.

Designamine is not approved for use by anyone younger than 18 years old.

How should I take desipramine?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may

occasionally change your dose. Use the medicine exactly as directed.

If you need surgery, tell your surgeon you currently use desipramine. You may need to stop for a short time.

It may take up to 3 weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Do not stop using desipramine suddenly, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine.

Store at room temperature away from moisture and heat. Keep the bottle tightly closed when not in use.

Desipramine dosing information

Usual Adult Dose for Depression:

100 to 200 mg orally per day

-Maximum dose: 300 mg/day

Comments:

- -Doses should be initiated at a lower level and increased according to tolerance and clinical response.
- -In severely ill patients, doses may be further increased to 300 mg per day if needed.
- -Treatment of patients requiring as much as 300 mg should generally be initiated in a hospital setting.

Use: Treatment of depression

Usual Geriatric Dose for Depression:

25 to 100 mg orally per day

-Maximum dose: 150 mg/day

Comments:

- -Doses should be initiated at a lower level and increased according to tolerance and clinical response.
- -Initial treatment may be given once a day or in divided doses.
- -In severely ill patients, doses may be further increased to 150 mg/day if needed.
- -Most patients should not exceed doses of 100 mg/day.

Use: Treatment of depression

Detailed Desipramine dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. An overdose of desipramine can be

Overdose symptoms may include irregular heartbeats, feeling like you might pass out, seizures, or coma.

What should I avoid while taking desipramine?

Do not drink alcohol. Dangerous side effects or death can occur when alcohol is combined with desipramine.

Avoid driving or hazardous activity until you know how desipramine will affect you. Your reactions could be impaired.

Avoid exposure to sunlight or tanning beds. Desipramine can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

What other drugs will affect desipramine?

Using desipramine with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Many drugs can affect desipramine. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all your current medicines and any medicine you start or stop using.

1 Desipramine drug interactions (more detail)

Does desipramine interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

desipramine	
+	
Enter a drug name	Add

More about desipramine

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (38)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: tricyclic antidepressants

- Breastfeeding
- En español

Patient resources

· Desipramine advanced reading

Other brands

Norpramin

Professional resources

- Desipramine monograph
- Desipramine (FDA)

Other brands

Norpramin

Related treatment guides

- Irritable Bowel Syndrome
- Dysautonomia
- ADHD
- Vulvodynia
- Depression

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 12.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

രു Risk data available

CSA Schedule* N/A Not a controlled drug

Approval History Drug history at FDA

User Reviews & Ratings

8.2 / 10

38 Reviews

Images

Desipramine Hydrochloride 150 mg (I 85)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

V