



[Home](#)

2. LoSeasonique

# LoSeasonique

**Generic name:** [ethinyl estradiol and levonorgestrel \(extended-cycle\)](#) [ *ETH-in-ill-ess-tra-DYE-ol-and-lee-voe-nor-JESS-trel* ]

**Drug class:** [Contraceptives](#)

Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on Aug 7, 2023.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#)

## What is LoSeasonique?

LoSeasonique is a combination drug that contains female hormones that prevent ovulation (the release of an egg from an ovary). This medication also causes changes in your cervical mucus and uterine lining, making it harder for sperm to reach the uterus and harder for a fertilized egg to attach to the uterus.

LoSeasonique is used as [contraception](#) to prevent pregnancy.

LoSeasonique may also be used for purposes not listed in this medication guide.

## Warnings

**Do not use LoSeasonique if you are pregnant or if you have recently had a baby.**

You should not take LoSeasonique if you have any of the following conditions: uncontrolled high blood pressure, heart disease, a blood-clotting disorder, circulation problems, diabetic problems with your eyes or kidneys, unusual vaginal bleeding, liver disease or liver cancer, severe migraine headaches, if you smoke and are over 35, or if you have ever had breast or uterine cancer, jaundice caused by birth control pills, a heart attack, a stroke, or a blood clot.

Taking LoSeasonique can increase your risk of blood clots, stroke, or heart attack, especially if you have certain other conditions, or if you are overweight.

**Smoking can greatly increase your risk of blood clots, stroke, or heart attack.** You should not take this medicine if you smoke and are over 35 years old.

Missing a pill increases your risk of becoming pregnant. Carefully follow the "missed dose" instructions if you forget to take your LoSeasonique tablets.

Some drugs can make LoSeasonique less effective in preventing pregnancy, including antibiotics, hepatitis C medications, HIV/AIDS medications, seizure medications, or barbiturate sedatives. Tell your doctor about all other medications you use.

## Before taking this medicine

Taking LoSeasonique can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, high cholesterol, or if you are overweight. Your risk of stroke or blood clot is highest during your first year of taking LoSeasonique. Your risk is also high when you restart this medicine after not taking it for 4 weeks or longer.

**Smoking can greatly increase your risk of blood clots, stroke, or heart attack.** Your risk increases the older you are and the more you smoke. You should not take this medicine if you smoke and are over 35 years old.

**Do not use if you are pregnant.** Stop using this medicine and tell your doctor right away if you become pregnant, or if you miss two menstrual periods in a row. If you have recently had a baby, wait at least 4 weeks before taking LoSeasonique.

You should not take LoSeasonique if you have:

- untreated or uncontrolled high blood pressure;
- heart disease (coronary artery disease, uncontrolled heart valve disorder, history of heart attack, stroke, or blood clot);
- a blood-clotting disorder or circulation problems;
- problems with your eyes, kidneys or circulation caused by diabetes;
- a history of hormone-related cancer such as breast or uterine cancer;
- unusual vaginal bleeding that has not been checked by a doctor;
- liver disease or liver cancer;
- severe migraine headaches (with aura, numbness, weakness, or vision changes), especially if you are older than 35;
- a history of jaundice caused by pregnancy or birth control pills; or
- if you smoke and are over 35 years old.

To make sure LoSeasonique are safe for you, tell your doctor if you have ever had:

- heart disease, high blood pressure, or if you are prone to having blood clots;
- high cholesterol or triglycerides, or if you are overweight;
- depression;
- diabetes, underactive thyroid, gallbladder disease;
- a seizure or migraine headache;
- irregular menstrual cycles; or
- fibrocystic breast disease, lumps, nodules, or an abnormal mammogram.

The hormones in LoSeasonique can pass into breast milk and may harm a nursing baby. This medication may also slow breast milk production. Do not use if you are breast feeding a baby.

 [LoSeasonique pregnancy and breastfeeding warnings](#) (more detail)

## How should I take LoSeasonique?

Take LoSeasonique exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

You will take your first pill on the first day of your period or on the first Sunday after your period begins. You may need to use back-up birth control, such as condoms or a spermicide, when you first start using this medication. Follow your doctor's instructions.

Take one pill every day, no more than 24 hours apart. When the pills run out, start a new pack the following day. You may get pregnant if you do not take one pill daily. Get your prescription refilled before you run out of pills completely.

You will not have a menstrual period every month while you are taking an extended-cycle birth control pill. Instead, your period should occur every 12 weeks.

The 91-day LoSeasonique pack contains three trays with cards that hold 84 "active" pills and seven "reminder" pills. You must use the pills in a certain order to keep you on a regular cycle. Trays 1 and 2 each hold 28 pills. Tray 3 holds 35 pills, including the 7 reminder pills. Your period should begin while you are using these reminder pills.

You may have breakthrough bleeding, especially during the first 3 months. Tell your doctor if this bleeding continues or is very heavy.

Use a back-up birth control if you are sick with severe vomiting or diarrhea.

If you need surgery or medical tests or if you will be on bed rest, you may need to stop using this medication for a short time. Any doctor or surgeon who treats you should know that you are using LoSeasonique.

While taking LoSeasonique, you will need to visit your doctor regularly.

Store LoSeasonique at room temperature away from moisture and heat.

 [Detailed LoSeasonique dosage information](#)

## What happens if I miss a dose?

Missing a LoSeasonique pill increases your risk of becoming pregnant.

If you miss one "active" pill, take two pills on the day that you remember. Then take one pill per day for the rest of the pack.

If you miss two "active" pills in a row, take two pills per day for two days in a row. Then take one pill per day for the rest of the pack. Use back-up birth control for at least 7 days following the missed pills.

If you miss three "active" pills in a row, do not take the missed pills. Continue taking 1 pill per day on schedule according to the pill package and leave the missed pills in the package. You may have some bleeding or spotting if you miss three pills in a row. Use back-up birth control for at least the next 7 days.

If you miss any reminder pills, throw them away and keep taking one pill per day until the pack is empty. You do not need back-up birth control if you miss a reminder pill. If your period does not start while you are taking the reminder pills, call your doctor because you might be pregnant.

# What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose symptoms may include nausea, vomiting, and vaginal bleeding.

## What to avoid

**Do not smoke while taking LoSeasonique**, especially if you are older than 35 years of age.

LoSeasonique will not protect you from sexually transmitted diseases--including HIV and AIDS. Using a condom is the only way to protect yourself from these diseases.

## LoSeasonique side effects

Get emergency medical help if you have **signs of an allergic reaction to LoSeasonique**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Stop using LoSeasonique and call your doctor at once if you have:

- sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance;
- sudden cough, wheezing, rapid breathing, coughing up blood;
- pain, swelling, warmth, or redness in one or both legs;
- chest pain or heavy feeling, pain spreading to the jaw or shoulder, nausea, sweating, general ill feeling;
- a change in the pattern or severity of migraine headaches;
- nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- swelling in your hands, ankles, or feet;
- a breast lump; or
- symptoms of depression (sleep problems, weakness, tired feeling, mood changes).

Common LoSeasonique side effects may include:

- mild nausea (especially when you first start taking this medicine), vomiting, bloating, stomach cramps;
- breast tenderness or swelling, nipple discharge;
- freckles or darkening of facial skin, increased hair growth, loss of scalp hair;
- changes in weight or appetite;
- problems with contact lenses;
- vaginal itching or discharge; or
- changes in your menstrual periods, decreased sex drive.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

## Related/similar drugs

### Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

#### Reviews & ratings

6.1 / 10

[2,624 Reviews](#)

[View more](#)

#### FEATURED

### Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

#### Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

### Depo Provera

Depo-Provera is used as contraception to prevent pregnancy. It is also used to reduce pain cause by ...

#### Reviews & ratings

4.2 / 10

[1,967 Reviews](#)

[View more](#)

### Nexplanon

Nexplanon is a long-term, reversible birth control implant that lasts for three years. Learn about ...

#### Reviews & ratings

5.0 / 10

[4,864 Reviews](#)

[View more](#)

## Yaz

Yaz (drospirenone and ethinyl estradiol) is used as a contraceptive to prevent pregnancy. Includes ...

### Reviews & ratings

5.9 / 10

[997 Reviews](#)

[View more](#)

## Slynd

Slynd (drospirenone) is a birth control pill used for the prevention of pregnancy. Includes Slynd ...

### Reviews & ratings

5.8 / 10

[386 Reviews](#)

[View more](#)

## What other drugs will affect LoSeasonique?

Some drugs can make LoSeasonique less effective, which may result in pregnancy. Other drugs may interact with ethinyl estradiol and levonorgestrel, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

 [LoSeasonique drug interactions](#) (more detail)

## Does LoSeasonique interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

LoSeasonique

+

Enter a drug name

Add

## More about LoSeasonique (ethinyl estradiol / levonorgestrel)

- [Check interactions](#)

- [Compare alternatives](#)
- [Reviews \(118\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [FDA approval history](#)
- [Drug class: contraceptives](#)

## Patient resources

### Other brands

[Aviane](#), [Jolessa](#), [Altavera](#), [Seasonique](#), ... +15 more

## Professional resources

- [LoSeasonique prescribing information](#)

### Other brands

[Aviane](#), [Vienna](#), [Jolessa](#), [Altavera](#), ... +49 more

## Related treatment guides

- [Birth Control](#)
- [Gonadotropin Inhibition](#)
- [Abnormal Uterine Bleeding](#)
- [Endometriosis](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use LoSeasonique only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.


[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.01.

### Availability

**Rx** Prescription only

### Pregnancy & Lactation


 Risk data available

### CSA Schedule\*

**N/A** Not a controlled drug



### Approval History

 Drug history at FDA



## User Reviews & Ratings

5.3 / 10

[118 Reviews](#)

## Related News

[Certain Birth Control Pills, Devices Linked to Higher Heart Risks](#)

## Images

[LoSeasonique ethinyl estradiol 0.02 mg / levonorgestrel 0.1 mg \(b 28\)](#)



[View larger images](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



### About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

### Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

### Support

[Help center](#)

[Sitemap](#)



Contact us



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.