

Home

Oxaprozin

Oxaprozin 🕁

Generic name: oxaprozin [ox-a-PROE-zin]

Brand names: Coxanto, Daypro

Dosage forms: oral capsule (300 mg), oral tablet (600 mg)

Drug class: Nonsteroidal anti-inflammatory drugs



Medically reviewed by Drugs.com on Jan 9, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is oxaprozin?

Oxaprozin is a nonsteroidal anti-inflammatory drug (NSAID) that is used in adults to treat signs and symptoms of osteoarthritis or rheumatoid arthritis. Oxaprozin is also used to treat rheumatoid arthritis in children at least 6 years old.

Oxaprozin may also be used for purposes not listed in this medication guide.

Oxaprozin side effects

Get emergency medical help if you have **signs of an allergic reaction**: (sneezing, runny or stuffy nose; wheezing, trouble breathing, hives, swelling of your face, lips, tongue, or throat) or **a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes.

Get emergency medical help if you have **signs of a heart attack or stroke**: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, or feeling short of breath.

Oxaprozin may cause serious side effects. Stop using oxaprozin and call your doctor at once if you have:

- any skin rash, no matter how mild, or a patchy skin color that worsens in sunlight;
- · flu-like symptoms;
- severe headache, blurred vision, pounding in your neck or ears;
- heart problems--swelling, rapid weight gain, feeling short of breath;
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- **liver problems**--diarrhea, loss of appetite, nausea, vomiting, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);

- kidney problems--swelling, urinating less, feeling tired or short of breath;
- high blood potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement; or
- low red blood cells (anemia)--pale skin, tiredness, feeling light-headed or short of breath, cold hands and feet.

Oxaprozin may worsen asthma attacks in people who have asthma. Talk with your doctor about your own risk.

Common side effects of oxaprozin may include:

- nausea, upset stomach;
- diarrhea, constipation; or
- · rash.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Oxaprozin side effects (more detail)

Related/similar drugs

Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

Reviews & ratings

7.0 / 10

223 Reviews

View more

Cymbalta

Cymbalta (duloxetine) is used to treat major depressive disorder, general anxiety disorder and ...

Reviews & ratings

6.3 / 10

1,899 Reviews

View more

FEATURED

Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used alone or with other ...

Reviews & ratings

5.7 / 10

145 Reviews

View more

Celecoxib

Celecoxib is a nonsteroidal antiinflammatory drug (NSAID) used to treat pain or inflammation ...

Reviews & ratings

7.0 / 10

280 Reviews

View more

Hydroxychloroquine

Hydroxychloroquine is a quinoline drug used to treat or prevent malaria. It's also used to treat ...

Reviews & ratings

6.4 / 10

356 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

Reviews & ratings

7.1 / 10

191 Paviane

View more

Diclofenac

Diclofenac is a nonsteroidal antiinflammatory drug used to treat pain and inflammation associated ...

Reviews & ratings

7.3 / 10

695 Reviews

View more

Naproxen

Naproxen is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

683 Reviews

View more

Duloxetine

Duloxetine is a selective serotonin and norepinephrine reuptake inhibitor antidepressant used to ...

Reviews & ratings

6.2 / 10

2,456 Reviews

View more

Warnings

Oxaprozin can increase your risk of fatal heart attack or stroke. Do not use oxaprozin just before or after heart bypass surgery (coronary artery bypass graft, or CABG). Oxaprozin may also cause stomach or intestinal bleeding, which can be fatal.

Before taking this medicine

Oxaprozin can increase your risk of fatal heart attack or stroke. Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG). Oxaprozin may also cause stomach or intestinal bleeding, which can be fatal. This can occur without warning, especially in older adults.

You should not take oxaprozin if you are allergic to it, or if you have ever had an asthma attack, hives, or a severe allergic reaction (sneezing, runny or stuffy nose, wheezing) after taking aspirin or another NSAID.

Tell your doctor if you have or ever had:

- congestive heart failure, heart disease, high blood pressure, high cholesterol, diabetes, or if you smoke;
- a heart attack, stroke, or blood clot;
- · stomach ulcers or bleeding;
- asthma;
- fluid retention:
- · liver disease; or
- kidney disease (or if you are on dialysis).

It may be harder for you to get pregnant while you are using this medicine.

Do not use oxaprozin during pregnancy unless your doctor tells you to. Using an NSAID in the last 20 weeks can harm the unborn baby or cause pregnancy complications.

Ask a doctor if it is safe to breastfeed while using this medicine.

How should I take oxaprozin?

Follow all directions on your prescription label and read all medication guides or instruction sheets. **Take the lowest dose that is effective in treating your condition.**

Your dose needs may change if you switch to a different brand, strength, or form of oxaprozin. **Avoid medication** errors by using exactly as directed on the label, or as prescribed by your doctor.

Oxaprozin is usually taken once per day. Follow your doctor's instructions.

Doses are based on weight. Your dose may change if you gain or lose weight.

If you use this medicine long-term, you may need frequent medical tests.

This medicine may cause false results on a drug-screening urine test. Tell the laboratory staff that you use oxaprozin.

Store tightly closed at room temperature, away from moisture, heat, and light.

Detailed Oxaprozin dosage information

What happens if I miss a dose?

Call your doctor for instructions if you miss a dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking oxaprozin?

Avoid drinking alcohol and smoking, since it can increase your risk of stomach bleeding.

Avoid taking aspirin or other NSAIDs unless your doctor tells you to.

Ask a doctor or pharmacist before using other medicines for pain, fever, swelling, or cold/flu symptoms. They may contain ingredients similar to oxaprozin (such as aspirin, ibuprofen, ketoprofen, or naproxen).

Oxaprozin could make you sunburn more easily. Avoid sunlight or tanning beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

What other drugs will affect oxaprozin?

Ask your doctor before using oxaprozin if you take an antidepressant. Taking certain antidepressants with an NSAID may cause you to bruise or bleed easily.

Oxaprozin can harm your kidneys, especially if you also use certain medicines for infections, cancer, osteoporosis, organ transplant rejection, high blood pressure, pain or arthritis (including Advil, Motrin, and Aleve).

Tell your doctor about all your current medicines. Many drugs can affect oxaprozin, especially:

- a blood thinner--warfarin, Coumadin, Jantoven;
- medicine used to prevent blood clots--alteplase, clopidogrel, dipyridamole, ticlopidine, and others; or
- steroid medicine (such as prednisone).

This list is not complete and many other drugs may affect oxaprozin. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Oxaprozin drug interactions (more detail)

Does oxaprozin interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

oxaprozin	
+	
Enter a drug name	Add

More about oxaprozin

Check interactions

- · Compare alternatives
- Pricing & coupons
- Reviews (8)
- Drug images
- Latest FDA alerts (4)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: Nonsteroidal anti-inflammatory drugs
- Breastfeeding

Patient resources

· Oxaprozin advanced reading

Other brands

Daypro, Coxanto

Professional resources

- Oxaprozin monograph
- Oxaprozin Capsules (FDA)
- Oxaprozin Tablets (FDA)

Other brands

Daypro, Coxanto

Related treatment guides

- Osteoarthritis
- Rheumatoid Arthritis
- Juvenile Rheumatoid Arthritis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ঝ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

6.4 / 10

8 Reviews

Images

Oxaprozin 600 mg (C 01 70)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright $\ensuremath{\texttt{©}}$ 2000-2025 Drugs.com. All rights reserved.

