

[Home](#)

2. Esgic

Esgic

Generic name: [acetaminophen, butalbital, and caffeine](#) [a-SEET-a-MIN-oh-fen, bue-TAL-bi-tal, and-KAF-eeen]

Brand names: Esgic, [Fioricet](#), [Zebutal](#)

Drug class: [Analgesic combinations](#)



[Medically reviewed](#) by Drugs.com on Jan 13, 2025. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)



The Esgic brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be [generic equivalents available](#).

What is Esgic?

[Acetaminophen](#) is a pain reliever and fever reducer.

Butalbital is in a group of drugs called [barbiturates](#). It relaxes muscle contractions involved in a [tension headache](#).

Caffeine is a central nervous system stimulant. It relaxes muscle contractions in blood vessels to improve blood flow.

Esgic is a combination medicine used to treat tension headaches that are caused by muscle contractions.

Esgic may also be used for purposes not listed in this medication guide.

Esgic side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. **Stop taking Esgic and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.** If you have this type of reaction, you should never again take any medicine that contains acetaminophen.

This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have:

- confusion, a seizure;
- [shortness of breath](#);
- a light-headed feeling, like you might pass out; or

- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Common side effects of Esgic may include:

- drowsiness, dizziness;
- feeling light-headed;
- nausea, vomiting, stomach pain;
- drunk feeling; or
- shortness of breath.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Esgic side effects](#) (more detail)

Related/similar drugs

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-the-counter painkiller and fever ...

Reviews & ratings

6.1 / 10

[13 Reviews](#)

[View more](#)

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

5.8 / 10

[30 Reviews](#)

[View more](#)

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

— — — — —

5.3 / 10

[685 Reviews](#)

[View more](#)

Celebrex

Celebrex is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation. Learn about ...

Reviews & ratings

7.0 / 10

[226 Reviews](#)

[View more](#)

Botox

Botox is used to treat chronic migraines, excessive sweating, bladder conditions, eye muscle ...

Reviews & ratings

5.7 / 10

[473 Reviews](#)

[View more](#)

Celecoxib

Celecoxib is a nonsteroidal anti-inflammatory drug (NSAID) used to treat pain or inflammation ...

Reviews & ratings

7.0 / 10

[280 Reviews](#)

[View more](#)

Diclofenac

Diclofenac is a nonsteroidal anti-inflammatory drug used to treat pain and inflammation associated ...

Reviews & ratings

7.3 / 10

[695 Reviews](#)

[View more](#)

Propranolol

Propranolol is a beta-blocker that is used to treat tremors, chest pain, high blood pressure, heart ...

Reviews & ratings

8.2 / 10

[1,376 Reviews](#)

[View more](#)

Naproxen

Naproxen is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

[683 Reviews](#)

[View more](#)

Amitriptyline

Amitriptyline is a tricyclic antidepressant used to treat depression and, off-label, conditions ...

Reviews & ratings

7.5 / 10

[1,803 Reviews](#)

[View more](#)

Warnings

Do not use this medicine if you have used an MAO inhibitor in the past 14 days. A dangerous [drug interaction](#) could occur. MAO inhibitors include [isocarboxazid](#), linezolid, [methylene blue](#) injection, [phenelzine](#), and [tranylcypromine](#).

Do not take more of this medication than is recommended. **An overdose of acetaminophen can damage your liver or cause death.** Call your doctor at once if you have nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, or jaundice (yellowing of your skin or eyes).

In rare cases, acetaminophen may cause a severe skin reaction. **Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.**

Before taking this medicine

Do not use Esgic if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, [rasagiline](#), [selegiline](#), and tranylcypromine.

You should not use Esgic if you are allergic to it, if you have porphyria, or if you have recently used alcohol, sedatives, tranquilizers, or other opioids.

Tell your doctor if you have ever had:

- [liver disease](#), cirrhosis, a history of [alcoholism](#) or [drug addiction](#), or if you drink more than 3 alcoholic beverages per day;
- kidney disease;
- [stomach ulcer](#) or bleeding;
- a history of skin rash caused by any medication; or
- a history of mental illness or suicidal thoughts.

It is not known whether this medicine will harm an unborn baby. **If you use butalbital while you are pregnant, your baby could become dependent on the drug.** This can cause life-threatening withdrawal symptoms in the baby after it is born. Babies born dependent on habit-forming medicine may need medical treatment for several weeks. Tell your doctor if you are pregnant or plan to become pregnant.

This medicine can pass into breast milk and may harm a nursing baby. Tell your doctor if you are breastfeeding a baby.

Not approved for use by anyone younger than 12 years old.

How should I take Esgic?

Follow all directions on your prescription label. Do not take more of this medication than recommended. **An overdose can damage your liver or cause death.** Tell your doctor if the medicine seems to stop working as well in relieving your pain.

Butalbital may be habit-forming. **Never share this medicine with another person, especially someone with a history of drug abuse or addiction.** Keep the medication in a place where others cannot get to it. Selling or giving away this medicine is against the law.

Store at room temperature away from moisture and heat.

Keep track of the amount of medicine used from each new bottle. Butalbital is a drug of abuse and you should be aware if anyone is using your medicine improperly or without a prescription.

 [Detailed Esgic dosage information](#)

What happens if I miss a dose?

Since Esgic is used when needed, you may not be on a dosing schedule. If you are on a schedule, use the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of Esgic can be fatal.**

The first signs of an [acetaminophen overdose](#) include loss of appetite, nausea, vomiting, stomach pain, sweating, and confusion or weakness. Later symptoms may include pain in your upper stomach, dark urine, and yellowing of your skin or the whites of your eyes.

Overdose symptoms may also include insomnia, restlessness, tremor, diarrhea, increased shallow breathing, uneven heartbeats, seizure (convulsions), or fainting.

What should I avoid while taking acetaminophen, butalbital, and caffeine?

This medication can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert.

Avoid drinking alcohol. It may increase your risk of liver damage while taking acetaminophen.

Ask a doctor or pharmacist before using any other cold, allergy, pain, or sleep medication. Acetaminophen (sometimes abbreviated as APAP) is contained in many combination medicines. **Taking certain products together can cause you to get too much acetaminophen which can lead to a fatal overdose.** Check the label to see if a medicine contains acetaminophen or APAP.

While you are taking this medication, avoid taking diet pills, caffeine pills, or other stimulants (such as ADHD medications) without your doctor's advice.

What other drugs will affect Esgic?

Taking Esgic with other drugs that make you sleepy or slow your breathing can cause dangerous or life-threatening side effects. Ask your doctor before taking Esgic with a sleeping pill, opioid pain medicine, muscle relaxer, or medicine for anxiety, [depression](#), or seizures.

Other drugs may affect Esgic, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Tell your doctor about all other medicines you use.

 [Esgic drug interactions](#) (more detail)

Does Esgic interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

+

Add

More about Esgic (acetaminophen / butalbital / caffeine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(11\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: analgesic combinations](#)
- [En español](#)

Patient resources

- [Esgic advanced reading](#)

Other brands

[Fioricet](#), [Esgic-Plus](#), [Medigesic](#), [Geone](#), ... +9 more

Professional resources

- [Esgic prescribing information](#)
- [Esgic Tablets \(FDA\)](#)

Other brands

[Fioricet](#), [Esgic-Plus](#), [Phrenilin Forte](#), [Vanatol LQ](#), ... +3 more

Related treatment guides

- [Headache](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

Availability
Discontinued

Pregnancy & Lactation
Risk data available

CSA Schedule*
N/A Not a controlled drug



User Reviews & Ratings

9.6 / 10

[11 Reviews](#)

Images

[Esgic 325 mg / 50 mg / 40 mg \(535 11 Logo\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.