

Home

2. Bacitracin/neomycin/polymyxin b

Bacitracin, neomycin, and polymyxin B (topical)

Generic name: bacitracin, neomycin, and polymyxin B (topical) [BAS-i-TRAY-sin, NEE-oh-MYE-sin, POL-ee-MIX-in-B]

Brand names: Medi-Quik, Neosporin, Triple Antibiotic, Septa, Mycitracin Triple Antibiotic, ... show all 8 brands

Dosage form: topical ointment (400 units-3.5 mg-5000 units/g)

Drug class: Topical antibiotics



Medically reviewed by Drugs.com on Sep 13, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking What to avoid Interactions

What is bacitracin, neomycin, and polymyxin B?

Bacitracin, neomycin, and polymyxin B are antibiotics that kill bacteria on your skin.

Bacitracin, neomycin, and polymyxin B topical (for the skin) is a combination medicine used as a first aid antibiotic to prevent infections in minor cuts, scrapes, or burns on your skin.

Bacitracin, neomycin, and polymyxin B topical may also be used for other purposes not listed in this medication guide.

bacitracin, neomycin, and polymyxin B side effects

Get emergency medical help if you have any of these **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

A rare but serious side effect of neomycin is hearing loss, which has occurred in people using other forms of neomycin. It is unlikely that you would absorb enough of this medicine through your skin to cause this effect. Call your doctor at once if you notice any changes in your hearing.

Call your doctor at once if you have severe redness or irritation, swelling, pus, oozing, or other signs of infection.

Common side effects of bacitracin, neomycin, and polymyxin B may include:

- mild itching or rash; or
- minor skin irritation after using the medicine.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Bacitracin / neomycin / polymyxin b topical side effects (more detail)

Related/similar drugs

Bactrim

Bactrim (sulfamethoxazole and trimethoprim) is an antibiotic used to treat ear infections, urinary ...

Reviews & ratings

5.4 / 10

616 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

SMZ-TMP DS

SMZ-TMP DS (sulfamethoxazole and trimethoprim) is an antibiotic used to treat ear infections ...

Reviews & ratings

4.6 / 10

54 Reviews

View more

Septra

Septra is an antibiotic containing sulfamethoxazole and trimethoprim used to treat ear infections ...

Reviews & ratings

5.4 / 10

19 Reviews

View more

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use this medication if you are allergic to bacitracin, neomycin, or polymyxin B.

Ask a doctor or pharmacist if it is safe for you to take this medicine if you have other medical conditions, especially:

- · chronic ear infections; or
- a ruptured (torn) eardrum.

It is not known whether bacitracin, neomycin, and polymyxin B topical will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medication.

It is not known whether bacitracin, neomycin, and polymyxin B topical passes into breast milk or if it could harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

Bacitracin, neomycin, and polymyxin B topical should not be used on a child younger than 2 years old without medical advice.

How should I use bacitracin, neomycin, and polymyxin B?

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended.

Before you apply this medication, wash the skin area with soap and water and then dry it thoroughly.

Do not apply the ointment over large skin areas. Do not use on a deep cut, an animal bite, or a serious burn. Contact your doctor for instructions on how to treat these more severe skin injuries.

This medication can be applied up to 3 times each day, or as directed on the medicine label. Cover the wound with a bandage if desired. Clean the wound and use a new bandage each time you apply the medicine.

Use the ointment for as many days as recommended on the label or by your doctor, even if your symptoms start getting better. Your symptoms may improve before the infection is completely cleared.

Call your doctor if your symptoms do not improve after 7 days of treatment, or if they get worse while using bacitracin, neomycin, and polymyxin B topical.

Store at room temperature away from moisture and heat. Throw away any ointment not used before the expiration date on the medicine label.

What happens if I miss a dose?

Apply the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** use extra medicine to make up the missed dose.

What happens if I overdose?

An overdose of bacitracin, neomycin, and polymyxin B topical is not expected to be dangerous. Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

What should I avoid while using bacitracin, neomycin, and polymyxin B?

Do not take by mouth. Bacitracin, neomycin, and polymyxin B topical is for use only on your skin. Avoid getting this medicine in your eyes, nose, or mouth. If this does happen, rinse with water.

Avoid applying other creams, lotions, ointments, or other medicated skin products to the same areas you treat with bacitracin, neomycin, and polymyxin B.

What other drugs will affect bacitracin, neomycin, and polymyxin B?

It is not likely that other drugs you take orally or inject will have an effect on topically applied bacitracin, neomycin, and polymyxin B. But many drugs can interact with each other. Tell each of your health care providers about all medicines you use, including prescription and over-the-counter medicines, vitamins, and herbal products.

Bacitracin / neomycin / polymyxin b topical drug interactions (more detail)

Does bacitracin/neomycin/polymyxin b topical interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

bacitracin/neomycin/polymyxin b topical
+
Enter a drug name
Add

More about bacitracin / neomycin / polymyxin b topical

- Check interactions
- · Compare alternatives
- Reviews (6)
- · Side effects
- · Drug class: topical antibiotics
- En español

Patient resources

• Neomycin, polymyxin b, and bacitracin (Topical) advanced reading

Other brands

Neosporin, Triple Antibiotic, Curad Triple Antibiotic, Medi-Quik

Professional resources

Other brands

Triple Antibiotic

Related treatment guides

- · Bacterial Skin Infection
- Paronychia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 2.01.

Antibiotics 101

List of Common Antibiotics & Types

Antibiotics and Alcohol

Probiotics and Antibiotics

Antibiotics for Strep Throat

Antibiotics for a Tooth Infection

Antibiotics for UTI's

DRUG STATUS

Availability

OTC Over the counter

CSA Schedule*

N/A Not a controlled drug

User Reviews & Ratings

6 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





