

Home

2. Belbuca

Belbuca 🖘

Generic name: buprenorphine (oral/buccal) [BUE-pre-NOR-feen]

Drug class: Opioids (narcotic analgesics)

Medically reviewed by Philip Thornton, DipPharm. Last updated on Jul 22, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is Belbuca?

Belbuca buccal films contain buprenorphine, an opioid medication. An opioid is sometimes called a narcotic.

Belbuca buccal films are for around-the-clock treatment of moderate to severe chronic pain that is not controlled by other medicines.

Belbuca is not for use on an as-needed basis for occasional pain.

The Belbuca buccal film is placed between the gum and cheek where it adheres to the moist buccal mucosa to deliver the medicine as the film dissolves.

Warnings

Belbuca can slow or stop your breathing, and may be habit-forming. MISUSE OF OPIOID MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH, especially in a child or other person using Belbuca without a prescription.

Using Belbuca during pregnancy may cause life-threatening withdrawal symptoms in the newborn.

Fatal side effects can occur if you use Belbuca with alcohol, or with other drugs that cause drowsiness or slow your breathing.

Before using this medicine

You should not use Belbuca if you are allergic to buprenorphine, or if you have:

• severe asthma or trouble breathing, or a blockage in your stomach or intestines.

To make sure Belbuca is safe for you, tell your doctor if you have ever had:

- tooth problems, including a history of cavities;
- breathing problems, sleep apnea;

- kidney or liver disease (especially hepatitis B or C);
- enlarged prostate, urination problems;
- problems with your gallbladder, pancreas, or thyroid;
- a head injury, brain tumor, or seizures;
- drug addiction, methadone use, alcoholism, mental illness;
- mouth sores caused by cancer;
- an electrolyte imbalance (such as low levels of potassium or magnesium in your blood); or
- long QT syndrome, or if you take heart rhythm medication.

If you use Belbuca while you are pregnant, your baby could become dependent on the drug. This can cause lifethreatening withdrawal symptoms in the baby after it is born. Babies born dependent on habit-forming medicine may need medical treatment for several weeks. Tell your doctor if you are pregnant or plan to become pregnant.

Buprenorphine can pass into breast milk and may harm a nursing baby. You should not breastfeed while you are using Belbuca.

Belbuca is not approved for use by anyone younger than 18 years old.

Belbuca pregnancy and breastfeeding warnings (more detail)

How should I use Belbuca?

Use Belbuca exactly as prescribed by your doctor. Follow all directions on your prescription label. Buprenorphine can slow or stop your breathing. Never use Belbuca in larger amounts, or for longer than prescribed. Tell your doctor if Belbuca seems to stop working as well in relieving your pain.

Buprenorphine may be habit-forming, even at regular doses. Never share Belbuca with another person, especially someone with a history of drug abuse or addiction. MISUSE OF NARCOTIC MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH, especially in a child or other person using Belbuca without a prescription. Selling or giving away Belbuca is against the law.

Read all patient information, medication guides, and instruction sheets provided to you. Ask your doctor or pharmacist if you have any questions.

Place the Belbuca film against the inside of your cheek and hold it in place for 5 seconds. Once in place, the film will dissolve completely in about 30 minutes. Rinse your mouth with water after the medicine dissolves. Wait one hour after the film dissolves to brush your teeth to prevent damage to the teeth and gums. **Do not chew the Belbuca film or swallow it whole.**

Do not eat or drink anything until the Belbuca film has completely dissolved in your mouth.

You should receive regular dental checkups while using Belbuca.

While using Belbuca, you may need frequent blood tests to check your liver function.

If you need surgery, tell the surgeon ahead of time that you are using Belbuca.

Do not stop using Belbuca suddenly, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely

stop using buprenorphine.

Store at room temperature away from moisture and heat. Keep Belbuca in a place where a child cannot get to it. Keep track of your medicine. Buprenorphine is a drug of abuse and you should be aware if anyone is using your medicine improperly or without a prescription.

Do not keep leftover opioid medication. **Just one dose can cause death in someone using Belbuca accidentally or improperly.** Ask your pharmacist where to locate a drug take-back disposal program. If there is no take-back program, remove any unused films from the foil pack and flush the films down the toilet. Throw the empty foil pack into the trash.

Detailed Belbuca dosage information

What happens if I miss a dose?

Since Belbuca is used for pain, you are not likely to miss a dose. Skip any missed dose if it is almost time for your next scheduled dose. Do not use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. A buprenorphine overdose can be fatal, especially in a child or other person using Belbuca without a prescription.

Overdose symptoms may include extreme drowsiness or weakness, cold or clammy skin, severe muscle weakness, pinpoint pupils, weak pulse, very slow breathing, or coma.

What should I avoid while using Belbuca?

Do not drink alcohol. Dangerous side effects or death could occur.

Avoid applying the buccal film to an area where you have a mouth sore.

Buprenorphine may impair your thinking or reactions. Avoid driving or operating machinery until you know how Belbuca will affect you. Dizziness or severe drowsiness can cause falls, accidents, or severe injuries.

Belbuca side effects

Get emergency medical help if you have **signs of an allergic reaction to Belbuca**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Like other narcotic medicines, buprenorphine can slow your breathing. Death may occur if breathing becomes too weak.

A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up.

Call your doctor at once if you have:

- noisy breathing, sighing, shallow breathing, breathing that stops during sleep;
- slow heartbeat or weak pulse;
- a light-headed feeling, like you might pass out;

- · chest pain, fast heart rate, trouble breathing;
- severe constipation;
- opioid withdrawal symptoms shivering, goose bumps, increased sweating, feeling hot or cold, runny nose, watery eyes, diarrhea, muscle pain;
- low cortisol levels nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness; or
- liver problems nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Serious side effects may be more likely in older adults and those who are overweight, malnourished, or debilitated.

Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent.

Common Belbuca side effects may include:

- constipation, nausea, vomiting;
- headache, dizziness, drowsiness;
- increased sweating;
- sleep problems (insomnia); or
- pain anywhere in your body.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Belbuca side effects (more detail)

Related/similar drugs

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-thecounter painkiller and fever ...

Reviews & ratings

6.1 / 10

13 Reviews

View more

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions

such as headaches ...

Reviews & ratings

5.8 / 10

30 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Cymbalta

Cymbalta (duloxetine) is used to treat major depressive disorder, general anxiety disorder and ...

Reviews & ratings

6.3 / 10

1,899 Reviews

View more

Oxycodone

Oxycodone is an opioid analgesic used to treat moderate to severe pain; it has a high potential for ...

Reviews & ratings

6.9 / 10

1,142 Reviews

View more

Amitriptyline

Amitriptyline is a tricyclic antidepressant used to treat depression and, off-label, conditions ...

Reviews & ratings

View more

Naproxen

Naproxen is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

683 Reviews

View more

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

1,404 Reviews

View more

Tramadol

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

Reviews & ratings

7.0 / 10

1,876 Reviews

View more

Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

Reviews & ratings

6.0 / 10

618 Reviews

What other drugs will affect Belbuca?

You may have breathing problems or withdrawal symptoms if you start or stop taking certain other medicines.

Tell your doctor if you also use an antibiotic, antifungal medication, heart or blood pressure medication, seizure medication, or medicine to treat HIV or hepatitis C.

Opioid medication can interact with many other drugs and cause dangerous side effects or death. Be sure your doctor knows if you also use:

- cold or allergy medicines, bronchodilator asthma/COPD medication, or a diuretic ("water pill");
- · medicines for motion sickness, irritable bowel syndrome, or overactive bladder;
- other opioid medications opioid pain medicine or prescription cough medicine;
- a sedative like Valium diazepam, alprazolam, lorazepam, Xanax, Klonopin, Versed, and others;
- drugs that make you sleepy or slow your breathing a sleeping pill, muscle relaxer, medicine to treat mood disorders or mental illness; or
- drugs that affect serotonin levels in your body a stimulant, or medicine for depression, Parkinson's disease, migraine headaches, serious infections, or nausea and vomiting.

This list is not complete. Other drugs may interact with buprenorphine, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

Belbuca drug interactions (more detail)

How long does buprenorphine stay in your system?

Does Belbuca interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Belbuca	
+	
Enter a drug name	Add

Popular FAQ

What are the different brands of buprenorphine?	~
s buprenorphine an opiate / narcotic?	~
How long does opioid withdrawal last?	~

What are the different types of buprenorphine/naloxone?

More FAQ

• Is Belbuca a controlled substance?

View more FAQ...

More about Belbuca (buprenorphine)

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (250)
- Drug images
- Latest FDA alerts (4)
- Side effects
- Dosage information
- During pregnancy
- Support group
- FDA approval history
- Drug class: Opioids (narcotic analgesics)
- Breastfeeding
- En español

Patient resources

Other brands

Subutex, Sublocade, Butrans, Brixadi, ... +2 more

Professional resources

- Belbuca prescribing information
- Buprenorphine (AHFS Monograph)

Other brands

Sublocade, Butrans, Brixadi, Buprenex, Probuphine

V

Related treatment guides

- Chronic Pain
- Pain

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Belbuca only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 7.01.

DRUG STATUS

Availability Rx Prescription only Pregnancy & Lactation Risk data available CSA Schedule* 3 Moderate abuse potential Approval History Drug bistory et EDA

Drug history at FDA

WADA Class

്റ്റ് Anti-Doping Classification

User Reviews & Ratings

4.5 / 10

250 Reviews

Related News

More ER Docs Prescribing Meds To Fight Opioid Addiction

Indivior Announces FDA Approval of Label Changes for Sublocade (buprenorphine extended-release) Injection

Images

Belbuca 150 mcg buccal film (E1)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.