

Home

Losartan

Losartan 🕁

Generic name: losartan [loe-SAR-tan]

Brand names: Cozaar, Arbli

Dosage form: oral tablet (100 mg; 25 mg; 50 mg), oral suspension (10 mg/mL)

Drug class: Angiotensin receptor blockers

Medically reviewed by Melisa Puckey, BPharm. Last updated on Apr 6, 2025.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is losartan?

Losartan is used alone or in combination with other medications to treat high blood pressure in adults and children at least 6 years old.

Losartan is also used to decrease the risk of stroke in people who have high blood pressure and a heart condition called left ventricular hypertrophy (enlargement of the walls of the left side of the heart).

Losartan may not decrease the risk of stroke in African Americans who have these conditions.

Losartan is also used to treat kidney disease in people who have type 2 diabetes (condition in which the body does not use insulin normally and therefore cannot control the amount of sugar in the blood) and high blood pressure.

Losartan belongs to a class of medications called angiotensin II receptor antagonists. It works by blocking the action of certain natural substances that tighten the blood vessels, allowing the blood to flow more smoothly and the heart to pump more efficiently.

Warnings

Do not use losartan if you are pregnant, and tell your doctor right away if you become pregnant. It can cause injury or death to the unborn baby during your second or third trimester.

If you have diabetes, do not use losartan together with any medication that contains aliskiren (a blood pressure medicine).

You may also need to avoid taking losartan with aliskiren if you have kidney disease.

In rare cases, this medicine can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine.

Before taking this medicine

You should not use losartan if you are allergic to it.

If you have diabetes, do not take losartan with any medication that contains aliskiren (a blood pressure medicine).

To make sure this medicine is safe for you, tell your doctor if you have ever had:

- heart disease or congestive heart failure;
- an electrolyte imbalance (such as high levels of potassium in your blood);
- if you are on a low-salt diet;
- · liver disease; or
- kidney disease.

You may also need to avoid taking losartan with aliskiren if you have kidney disease.

Stop using this medicine and tell your doctor right away if you become pregnant. Losartan can cause injury or death to the unborn baby if you use the medicine during your second or third trimester.

Tell your doctor if you are breastfeeding.

i Losartan pregnancy and breastfeeding warnings (more detail)

How should I take losartan?

Take losartan exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

- · Losartan is usually taken once per day.
- You may take losartan with or without food.

Call your doctor if you are sick with vomiting or diarrhea, or if you are sweating more than usual. You can easily become dehydrated while taking losartan. This can lead to very low blood pressure, a serious electrolyte imbalance, or kidney failure.

- Your blood pressure will need to be checked often, and you may need frequent blood tests.
- If you have high blood pressure, keep using this medicine even if you feel well. High blood pressure often has no symptoms.

Store tightly closed at room temperature, away from moisture, heat, and light.

1 Losartan patient tips (more detail)

Dosing information

Usual Adult Dose for Diabetic Nephropathy:

Initial dose: 50 mg orally once a day Maximum dose: 100 mg orally once a day

Comments:

- -If hypertensive patients with left ventricular hypertrophy or type 2 diabetic nephropathy patients require titration from initial dose, increase to 100 mg orally once a day.
- -Antihypertension is substantially present within 1 week but in some studies the maximal effect occurred in 3 to 6 weeks.

Uses:

- -Treatment of hypertension.
- -To reduce the risk of stroke in patients with hypertension and left ventricular hypertrophy.
- -Treatment of diabetic nephropathy with an elevated serum creatinine and proteinuria (urinary albumin to creatinine ratio 300 mg/g or greater) in patients with type 2 diabetes and a history of hypertension.

Usual Adult Dose for Hypertension:

Initial dose: 50 mg orally once a day

Maximum dose: 100 mg orally once a day

Comments:

- -If hypertensive patients with left ventricular hypertrophy or type 2 diabetic nephropathy patients require titration from initial dose, increase to 100 mg orally once a day.
- -Antihypertension is substantially present within 1 week but in some studies the maximal effect occurred in 3 to 6 weeks.

Uses:

- -Treatment of hypertension.
- -To reduce the risk of stroke in patients with hypertension and left ventricular hypertrophy.
- -Treatment of diabetic nephropathy with an elevated serum creatinine and proteinuria (urinary albumin to creatinine ratio 300 mg/g or greater) in patients with type 2 diabetes and a history of hypertension.

Usual Pediatric Dose for Hypertension:

6 Years or Older:

Initial dose: 0.7 mg/kg orally once a day (up to 50 mg total)

Comments:

- -May be given as a tablet or suspension. If using the suspension, use an oral dosing syringe or an oral dosing cup to measure the prescribed dose. Do not use a teaspoon or tablespoon.
- -Doses above 1.4 mg/kg (or 100 mg) daily have not been studied in pediatric patients.

<u>Detailed Losartan dosage information</u>

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking losartan?

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

Do not use potassium supplements or salt substitutes, unless your doctor has told you to.

Losartan side effects

Get emergency medical help if you have **signs of an allergic reaction to losartan:** hives, difficulty breathing, swelling of your face, lips, tongue, or throat.

Losartan may cause serious side effects. Call your doctor at once if you have:

- · a light-headed feeling, like you might pass out;
- high blood potassium nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement; or
- kidney problems swelling, urinating less, feeling tired or short of breath.

Common losartan side effects may include:

- · dizziness, tiredness;
- low blood pressure;
- low blood sugar;
- · diarrhea;
- · back pain; or
- · cold symptoms such as stuffy nose, sneezing, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Losartan side effects (more detail)

Related/similar drugs

Cozaar

Cozaar (losartan) is used to treat high blood pressure (hypertension). Includes Cozaar side ...

Reviews & ratings

6.4 / 10

62 Reviews

View more

Altace

Altace (ramipril) is used to treat high blood pressure, heart failure, and to improve survival ...

Reviews & ratings

6.9 / 10

12 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Captopril

Captopril is used to treat hypertension, heart failure and to improve survival after a heart ...

Reviews & ratings

7 Reviews

View more

Lisinopril

Lisinopril is an ACE inhibitor. It is used to treat high blood pressure, congestive heart failure ...

Reviews & ratings

5.0 / 10

626 Reviews

View more

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

56 Reviews

View more

Ramipril

Ramipril treats high blood pressure, heart failure, and improves survival after a heart attack ...

Reviews & ratings

5.7 / 10

133 Reviews

View more

Enalapril

Enalapril is used to treat high blood pressure, congestive heart failure and to improve survival ...

Reviews & ratings

5.6 / 10

32 Reviews

View more

What other drugs will affect losartan?

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which may increase side effects or make the medicines less effective.

Tell your doctor about all your other medicines, especially:

- a diuretic or "water pill" that may increase blood potassium such as spironolactone, triamterene, amiloride;
- NSAIDs (nonsteroidal anti-inflammatory drugs) aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others; or
- · heart or blood pressure medication.

This list is not complete. Other drugs may affect losartan, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Losartan drug interactions (more detail)

Does losartan interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

losartan	
+	
Enter a drug name	Add

Popular FAQ

Can you eat bananas when taking losartan?

More FAQ

- Does Losartan block the receptor used by the Coronavirus?
- Losartan vs Valsartan What's the difference between them?
- Does losartan cause rapid heart rate, irregular heartbeat or low blood pressure?
- · Are losartan and losartan potassium the same or different drugs?

View more FAQ...

References

- 1. Arbli (losartan) Prescribing Information
- 2. Cozaar (losartan) Prescribing Information

More about losartan

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (578)
- Drug images
- · Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: angiotensin receptor blockers
- Breastfeeding

• En español

Patient resources

Other brands

Cozaar

Professional resources

- Losartan Potassium monograph
- Losartan (FDA)

Other brands

Cozaar, Arbli

Related treatment guides

- High Blood Pressure
- · Diabetic Kidney Disease
- Alport Syndrome

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medicine only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 19.01.

DRUG STATUS

Availability Rx Prescription only Pregnancy & Lactation Risk data available CSA Schedule* N/A Not a controlled drug Approval History Drug history at FDA

578 Reviews

Related News

FDA Approves Arbli (losartan potassium) Oral Suspension as the First Ready-to-Use Oral Liquid Losartan in the U.S.

Images

Losartan Potassium 50 mg (I 6)







View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







