

#### Home

- 2. Lidex
- Consumer

# Lidex ⊲»

Generic name: Fluocinonide Cream, Gel, Ointment, and Solution [ floo-oh-SIN-oh-nide ]

Brand names: Lidex, Lidex-E Drug class: Topical steroids

Medically reviewed by Drugs.com. Last updated on Jan 11, 2024.

Uses Before taking Warnings Dosage Side effects Overdose FAQ



The Lidex brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be generic equivalents available.

## **Uses of Lidex:**

- · It is used to treat skin irritation.
- It is used to treat psoriasis.
- It is used to treat skin rashes.

# What do I need to tell my doctor BEFORE I take Lidex?

• If you are allergic to Lidex (fluocinonide cream, gel, ointment, and solution); any part of Lidex (fluocinonide cream, gel, ointment, and solution); or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.

This medicine may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take Lidex (fluocinonide cream, gel, ointment, and solution) with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

# What are some things I need to know or do while I take Lidex?

- Tell all of your health care providers that you take Lidex (fluocinonide cream, gel, ointment, and solution). This includes your doctors, nurses, pharmacists, and dentists.
- Talk with your doctor before you use other drugs or products on your skin.

- Do not put on cuts, scrapes, or damaged skin.
- Use care when putting on a large part of the skin or where there are open wounds. Talk with the doctor.
- Do not use longer than you have been told by the doctor.
- Do not use tight-fitting diapers or plastic pants if treated part is in the diaper area. This may cause more drug to get into the body.
- Use with care in children. Talk with the doctor.
- This medicine may affect growth in children and teens in some cases. They may need regular growth checks. Talk
  with the doctor.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

# How is this medicine (Lidex) best taken?

Use Lidex (fluocinonide cream, gel, ointment, and solution) as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Use as you have been told, even if your signs get better.
- Do not take Lidex (fluocinonide cream, gel, ointment, and solution) by mouth. Use on your skin only. Keep out of your mouth, nose, and eyes (may burn).
- Wash your hands before and after use. Do not wash your hands after use if putting this on your hand.
- Clean affected part before use. Make sure to dry well.
- Put a thin layer on the affected skin and rub in gently.
- Do not use coverings (bandages, dressings) unless told to do so by the doctor.

#### What do I do if I miss a dose?

- Put on a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not put on 2 doses at the same time or extra doses.
- Detailed Lidex dosage information

# What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever;
   wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit.

- Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss.
- Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache, or slow healing.
- Skin changes (pimples, stretch marks, slow healing, hair growth).
- · Skin irritation.
- Thinning of the skin.
- i <u>Lidex side effects</u> (more detail)

# Related/similar drugs

## **Dupixent**

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

#### **Reviews & ratings**

7.2 / 10

499 Reviews

#### View more

#### **Stelara**

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

## Reviews & ratings

7.6 / 10

252 Reviews

### View more

#### FEATURED

### **Skyrizi**

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

#### **Reviews & ratings**

7.9 / 10

56 Reviews

#### View more

### Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

### **Reviews & ratings**

7.0 / 10

223 Reviews

View more

# **Hydrocortisone topical**

Hydrocortisone cream is used to treat eczema, dermatitis, psoriasis, hemorrhoids, insect bites or ...

### Reviews & ratings

7.0 / 10

63 Reviews

View more

### **Triamcinolone**

Triamcinolone is used to treat allergies, skin conditions, ulcerative colitis, and arthritis. Learn ...

#### **Reviews & ratings**

6.7 / 10

226 Reviews

View more

### **Dexamethasone**

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

### **Reviews & ratings**

5.9 / 10

162 Reviews

View more

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

#### **Reviews & ratings**

7.1 / 10

484 Reviews

View more

#### **Methotrexate**

Methotrexate is used to treat certain types of cancer of the breast, skin, head and neck, or lung ...

#### **Reviews & ratings**

6.5 / 10

335 Reviews

View more

### What are some other side effects of Lidex?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Burning or stinging.
- · Dry skin.
- · Itching.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

# If OVERDOSE is suspected:

If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## How do I store and/or throw out Lidex?

- Store at room temperature.
- · Protect from heat.

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so.
   Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

# **Consumer Information Use and Disclaimer**

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions
  about Lidex (fluocinonide cream, gel, ointment, and solution), please talk with your doctor, nurse, pharmacist, or
  other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

# Frequently asked questions

- How long should you use fluocinonide for?
- What causes Plaque Psoriasis?
- What is a substitute for fluocinonide cream?
- Is fluocinonide used for hair loss?
- Is fluocinonide an antifungal cream?
- · What is fluocinonide cream good for?
- Is fluocinonide a steroid?
- Is fluocinonide an antibiotic?
- Is fluocinonide safe to use on your face?

# More about Lidex (fluocinonide topical)

- · Check interactions
- · Compare alternatives
- Reviews (10)
- · Side effects
- Dosage information
- During pregnancy
- · Drug class: topical steroids
- Breastfeeding

### **Patient resources**

· Lidex advanced reading

#### Other brands

Vanos, Lidex-E

### **Professional resources**

- · Lidex prescribing information
- Lidex Gel (FDA)
- Lidex Ointment (FDA)
- Lidex Topical Solution (FDA)

### Other brands

Vanos

# Related treatment guides

- Atopic Dermatitis
- Dermatitis
- Eczema
- Psoriasis

# **Further information**

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

### **DRUG STATUS**

**Availability** 



**Pregnancy & Lactation** 

& Risk data available

**CSA Schedule\*** 

N/A Not a controlled drug

Approval History

10+ years FDA approved 1971

### **User Reviews & Ratings**

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com
Advertising policy
Attribution & citations

#### **Terms & privacy**

Terms of use

**Editorial policy** 

Privacy policy

### **Support**

Help center

**Sitemap** 

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







×