

[Home](#)

2. Paxil

# Paxil

**Generic name:** [paroxetine](#) [ *pa-ROX-a-teen* ]**Drug class:** [Selective serotonin reuptake inhibitors](#)Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on Aug 15, 2023.[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

## What is Paxil?

Paxil is an antidepressant that belongs to group of drugs called [selective serotonin reuptake inhibitors \(SSRIs\)](#).

Paroxetine affects chemicals in the brain that may be unbalanced in people with depression, anxiety, or other disorders.

Paxil is used to treat [depression](#), including major depressive disorder.

Paxil is also used to treat [panic disorder](#), obsessive-compulsive disorder (OCD), [anxiety disorders](#), [post-traumatic stress disorder \(PTSD\)](#), and premenstrual dysphoric disorder (PMDD).

## Warnings

You should not use Paxil if you are also taking [pimozide](#) or [thioridazine](#).

Do not use Paxil within 14 days before or 14 days after you have used an MAO inhibitor, such as [isocarboxazid](#), [linezolid](#), [methylene blue](#) injection, [phenelzine](#), [rasagiline](#), [selegiline](#), or [tranylcypromine](#).

Some young people have thoughts about suicide when first taking an antidepressant. Stay alert to changes in your mood or symptoms. Report any new or worsening symptoms to your doctor

**Seek medical attention right away if you have symptoms such as:** [agitation](#), hallucinations, muscle stiffness, twitching, loss of coordination, [dizziness](#), warmth or tingly feeling, [nausea](#), [vomiting](#), [diarrhea](#), fever, sweating, tremors, racing heartbeats, or a seizure (convulsions).

**Do not stop using Paxil without first asking your doctor.**

## Before taking this medicine

You should not use Paxil if you are allergic to paroxetine, or if you are also taking pimozide or thioridazine.

Do not use an MAO inhibitor within 14 days before or 14 days after you take paroxetine. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

After you stop taking Paxil you must wait at least 14 days before you start taking an MAO inhibitor.

To make sure Paxil is safe for you, tell your doctor if you have:

- heart disease, [high blood pressure](#), or a stroke;
- liver or kidney disease;
- a bleeding or blood clotting disorder;
- [seizures](#) or [epilepsy](#);
- [bipolar disorder](#) (manic depression), [drug addiction](#), or suicidal thoughts;
- sexual problems;
- narrow-angle [glaucoma](#); or
- low levels of sodium in your blood.

Be sure your doctor knows if you also take stimulant medicine, opioid medicine, [herbal products](#), or medicine for depression, mental illness, [Parkinson's disease](#), [migraine](#) headaches, serious infections, or prevention of [nausea and vomiting](#). These medicines may interact with paroxetine and cause a serious condition called serotonin syndrome.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor should check your progress at regular visits. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Taking an SSRI antidepressant such as Paxil during pregnancy may cause serious lung problems or other complications in the baby. However, you may have a relapse of depression if you stop taking your antidepressant. Tell your doctor right away if you become pregnant. **Do not start or stop taking this medicine without your doctor's advice.**

You should not breastfeed while using this medicine.

Paxil is not approved for use by anyone younger than 18 years old.

 [Paxil pregnancy and breastfeeding warnings](#) (more detail)

## How should I take Paxil?

Take Paxil exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Swallow the Paxil CR **extended-release tablet** whole and do not crush, chew, or break the tablet.

Shake the Paxil **oral suspension** (liquid) before you measure a dose. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

It may take up to 4 weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

**Tell your doctor if you have any changes in sexual function**, such as loss of interest in sex, trouble having an orgasm, or (in men) problems with erections or ejaculation. Some sexual problems can be treated.

Do not stop using Paxil suddenly, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine. Follow your doctor's instructions about tapering your dose.

Store at room temperature away from moisture, heat, and light.

 [Detailed Paxil dosage information](#)

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of paroxetine can be fatal.**

## What to avoid

Avoid driving or hazardous activity until you know how paroxetine will affect you. Your reactions could be impaired.

Ask your doctor before taking a nonsteroidal anti-inflammatory drug (NSAID) such as [aspirin](#), [ibuprofen \(Advil, Motrin\)](#), [naproxen \(Aleve\)](#), [celecoxib \(Celebrex\)](#), [diclofenac](#), [indomethacin](#), [meloxicam](#), and others. Using an NSAID with Paxil may cause you to bruise or bleed easily.

Drinking alcohol with this medicine can cause side effects.

## Paxil side effects

Get emergency medical help if you have **signs of an allergic reaction to Paxil** ([hives](#), difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, [sore throat](#), burning eyes, skin pain, red or purple [skin rash](#) with blistering and peeling).

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, [panic attacks](#), trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Call your doctor at once if you have:

- racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual;
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights;
- unusual bone pain or tenderness, swelling or bruising;
- changes in weight or appetite;
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), coughing up blood;
- **severe nervous system reaction** - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, fainting; or
- **low levels of sodium in the body** - [headache](#), confusion, slurred speech, severe weakness, loss of coordination, feeling unsteady.

**Seek medical attention right away if you have symptoms of serotonin syndrome, such as:** agitation, hallucinations, fever, sweating, shivering, [fast heart rate](#), muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Common Paxil side effects may include:

- vision changes;
- weakness, drowsiness, dizziness, tiredness;
- sweating, anxiety, shaking;
- sleep problems ([insomnia](#));
- loss of appetite, nausea, vomiting, diarrhea, [constipation](#);
- dry mouth, yawning;
- infection;
- headache; or
- decreased sex drive, [impotence](#), abnormal ejaculation, or difficulty having an orgasm.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Paxil side effects](#) (more detail)

## Related/similar drugs

### Lexapro

Lexapro is used to treat anxiety and major depressive disorder. Learn about side effects ...

#### Reviews & ratings

7.6 / 10

[2,376 Reviews](#)

[View more](#)

### Zoloft

Zoloft is an antidepressant used to treat major depression, obsessive-compulsive disorder, panic ...

#### Reviews & ratings

7.3 / 10

[1,972 Reviews](#)

[View more](#)

FEATURED

## Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

### Reviews & ratings

5.3 / 10

[685 Reviews](#)

[View more](#)

## Citalopram

Citalopram is an antidepressant (selective serotonin reuptake inhibitor) and is used to treat ...

### Reviews & ratings

7.4 / 10

[2,231 Reviews](#)

[View more](#)

## Venlafaxine

Venlafaxine is used to treat major depressive disorder, anxiety, and panic disorder. Learn about ...

### Reviews & ratings

6.5 / 10

[2,786 Reviews](#)

[View more](#)

## Clonazepam

Clonazepam is used to treat seizures and panic disorder. Learn about side effects, interactions and ...

### Reviews & ratings

8.2 / 10

[2,176 Reviews](#)

[View more](#)

## Prozac

Prozac (fluoxetine) is an SSRI antidepressant used to treat

antidepressant used to treat  
depression, OCD, panic disorder ...

### Reviews & ratings

7.2 / 10

[1,239 Reviews](#)

[View more](#)

## Buspirone

Buspirone is used to treat symptoms  
of anxiety, such as fear, tension,  
irritability and dizziness ...

### Reviews & ratings

5.9 / 10

[1,357 Reviews](#)

[View more](#)

## Xanax

Xanax (alprazolam) is a  
benzodiazepine medication used to  
treat anxiety and panic disorders ...

### Reviews & ratings

8.6 / 10

[915 Reviews](#)

[View more](#)

## Hydroxyzine

Hydroxyzine is an antihistamine used  
to treat itching, hives, and anxiety. It  
also acts as a ...

### Reviews & ratings

5.9 / 10

[1,404 Reviews](#)

[View more](#)

## What other drugs will affect Paxil?

Using Paxil with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- [cimetidine \(Tagamet\)](#), [digoxin](#), [St. John's wort](#), [tamoxifen](#), [theophylline](#), [tryptophan](#) (sometimes called L-tryptophan), [warfarin \(Coumadin, Jantoven\)](#);
- a diuretic or "water pill";
- heart rhythm medicine;
- [HIV](#) or AIDS medications;
- **certain medicines to treat [narcolepsy](#) or [ADHD](#)** - [amphetamine](#), [atomoxetine](#), [dextroamphetamine](#), [Adderall](#), [Dexedrine](#), [Evekeo](#), [Vyvanse](#), and others;
- **narcotic pain medicine** - [fentanyl](#), [tramadol](#);
- **medicine to treat anxiety, mood disorders, thought disorders, or mental illness** - such as [buspirone](#), [lithium](#), other [antidepressants](#), or [antipsychotics](#);
- **migraine headache medicine** - [sumatriptan](#), [rizatriptan](#), [zolmitriptan](#), and others; or
- **seizure medicine** - [phenobarbital](#), [phenytoin](#).

This list is not complete. Other drugs may interact with paroxetine, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible interactions are listed in this medication guide.

 [Paxil drug interactions](#) (more detail)

## Does Paxil interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

+

Add

## Popular FAQ

Does Benadryl help with or cause anxiety?



Why give Taxol (Paxel) before carboplatin?



## More FAQ

- [SSRIs vs SNRIs - What's the difference between them?](#)
- [What are some common side effects of antidepressants?](#)

[View more FAQ...](#)

## More about Paxil (paroxetine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(960\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Support group](#)
- [Drug class: selective serotonin reuptake inhibitors](#)
- [Breastfeeding](#)
- [En español](#)

## Patient resources

### Other brands

[Brisdelle](#), [Pexeva](#)

## Professional resources

- [Paxil prescribing information](#)
- [PARoxetine \(AHFS Monograph\)](#)

### Other brands

[Brisdelle](#), [Pexeva](#)

## Other formulations

- [Paxil CR](#)

## Related treatment guides

- [Anxiety and Stress](#)
- [Dysautonomia](#)
- [Anxiety](#)
- [Generalized Anxiety Disorder](#)



# Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Paxil only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 30.01.

## DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
10+ years	FDA approved 1992

## User Reviews & Ratings

7.1 / 10

[960 Reviews](#)

## Images

[Paxil 20 mg \(PAXIL 2 0\)](#)



[View larger images](#)

# Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)  
[Advertising policy](#)  
[Attribution & citations](#)

## Terms & privacy

[Terms of use](#)  
[Editorial policy](#)  
[Privacy policy](#)

## Support

[Help center](#)  
[Sitemap](#)  
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.