



[Home](#)

2. [Nortrel 7/7/7](#)

3. [Dosage](#)

## Nortrel 7/7/7 Dosage

**Generic name:** [NORETHINDRONE 0.5mg, ETHINYL ESTRADIOL 0.035mg; NORETHINDRONE 0.75mg, ETHINYL ESTRADIOL 0.035mg; NORETHINDRONE 1mg, ETHINYL ESTRADIOL 0.035mg;](#)

**Dosage form:** tablets

**Drug class:** [Contraceptives](#)

[Medically reviewed](#) by Drugs.com. Last updated on Dec 23, 2024.

To achieve maximum contraceptive effectiveness, Nortrel 7/7/7 tablets must be taken exactly as directed and at intervals not exceeding 24 hours. Nortrel 7/7/7 tablets are available with the blister pack tablet dispenser which is preset for a Sunday Start. Day 1 Start is also available.

### Sunday Start

When taking Nortrel 7/7/7, the first “active” tablet should be taken on the first Sunday after menstruation begins. If the period begins on Sunday, the first “active” tablet should be taken that day. Take one active tablet daily for 21 days followed by one white “reminder” tablet daily for 7 days. After 28 tablets have been taken, a new course is started the next day (Sunday). For the first cycle of a Sunday Start regimen, another method of contraception such as a condom or spermicide should be used until after the first 7 consecutive days of administration.

If the patient misses one (1) “active” tablet in Weeks 1, 2, or 3, the tablet should be taken as soon as she remembers. If the patient misses two (2) “active” tablets in Week 1 or Week 2, the patient should take two (2) tablets the day she remembers and two (2) tablets the next day; and then continue taking one (1) tablet a day until she finishes the pack. The patient should be instructed to use a back-up method of birth control such as a condom or spermicide if she has sex in the seven (7) days after missing pills. If the patient misses two (2) “active” tablets in the third week or misses three (3) or more “active” tablets in a row, the patient should continue taking one tablet every day until Sunday. On Sunday the patient should throw out the rest of the pack and start a new pack that same day. The patient should be instructed to use a back-up method of birth control if she has sex in the seven (7) days after missing pills.

Complete instructions to facilitate patient counseling on proper pill usage may be found in the Detailed Patient Labeling (“How to Take the Pill” section).

### Day 1 Start

The dosage of Nortrel 7/7/7, for the initial cycle of therapy, is one “active” tablet administered daily from the 1st through the 21st day of the menstrual cycle, counting the first day of menstrual flow as “Day 1” followed by one white “reminder” tablet daily for 7 days. Tablets are taken without interruption for 28 days. After 28 tablets have been taken, a new course is started the next day.

If the patient misses one (1) “active” tablet in Weeks 1, 2, or 3, the tablet should be taken as soon as she remembers. If the patient misses two (2) “active” tablets in Week 1 or Week 2, the patient should take two (2) tablets the day she remembers and two (2) tablets the next day; and then continue taking one (1) tablet a day until she finishes the pack. The

patient should be instructed to use a back-up method of birth control such as a condom or spermicide if she has sex in the seven (7) days after missing pills. If the patient misses two (2) “active” tablets in the third week or misses three (3) or more “active” tablets in a row, the patient should throw out the rest of the pack and start a new pack that same day. The patient should be instructed to use a back-up method of birth control if she has sex in the seven (7) days after missing pills.

Complete instructions to facilitate patient counseling on proper pill usage may be found in the Detailed Patient Labeling (“How to Take the Pill” section).

The use of Nortrel 7/7/7 for contraception may be initiated 4 weeks postpartum in women who elect not to breastfeed. When the tablets are administered during the postpartum period, the increased risk of thromboembolic disease associated with the postpartum period must be considered. (See [CONTRAINDICATIONS](#) and [WARNINGS](#) concerning thromboembolic disease. See also [PRECAUTIONS](#), [Nursing Mothers](#).) The possibility of ovulation and conception prior to initiation of medication should be considered.

(See Discussion of Dose Related Risk of Vascular Disease from Oral Contraceptives.)

Breakthrough bleeding, spotting, and amenorrhea are frequent reasons for patients discontinuing oral contraceptives. In breakthrough bleeding, as in all cases of irregular bleeding from the vagina, nonfunctional causes should be borne in mind. In undiagnosed persistent or recurrent abnormal bleeding from the vagina, adequate diagnostic measures are indicated to rule out pregnancy or malignancy. If pathology has been excluded, time or a change to another formulation may solve the problem. Changing to an oral contraceptive with a higher estrogen content, while potentially useful in minimizing menstrual irregularity, should be done only if necessary since this may increase the risk of thromboembolic disease.

Use of oral contraceptives in the event of a missed menstrual period:

1. If the patient has not adhered to the prescribed schedule, the possibility of pregnancy should be considered at the time of the first missed period and oral contraceptive use should be discontinued if pregnancy is confirmed.
2. If the patient has adhered to the prescribed regimen and misses two consecutive periods, pregnancy should be ruled out.

## More about Nortrel 7 / 7 / 7 (ethinyl estradiol / norethindrone)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(24\)](#)
- [Drug images](#)
- [Side effects](#)
- [During pregnancy](#)
- [Drug class: contraceptives](#)
- [En español](#)

## Patient resources

- [Nortrel 7/7/7 birth control drug information](#)

Other brands

[Lo Loestrin Fe](#), [Junel Fe 1/20](#), [Blisovi Fe 1/20](#), [Loestrin 21 1/20](#), ... +77 more

Professional resources

- [Nortrel 7/7/7 prescribing information](#)

Other brands

[Lo Loestrin Fe](#), [Blisovi Fe 1/20](#), [Loestrin 21 1/20](#), [Hailey Fe 1/20](#), ... +81 more

Related treatment guides

- [Birth Control](#)

Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	Drug history at FDA

User Reviews & Ratings

3.5 / 10

[24 Reviews](#)

Related Drugs

[Mirena](#), [Provera](#), [Depo Provera](#), [Nexplanon](#), [Sprintec](#), [Yaz](#), [Slynd](#), [drospirenone](#), [levonorgestrel](#), [medroxyprogesterone](#)

Related News

[Certain Birth Control Pills, Devices Linked to Higher Heart Risks](#)

## Images

Nortrel 7 7 7 inert (b 944)



[View larger images](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



### About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

### Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

### Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.