

Home

2. Piperacillin/tazobactam

Piperacillin and tazobactam 🖏

Generic name: piperacillin and tazobactam [PI-per-a-SIL-in-and-TAZ-oh-BAK-tam]

Brand name: Zosyn

Dosage forms: intravenous powder for injection (12 g-1.5 g; 2 g-0.25 g; 3 g-0.375 g; 36 g-4.5 g; 4 g-0.5 g), ... show all 2 dosage forms

Drug class: Penicillins / beta-lactamase inhibitors



Medically reviewed by Drugs.com on Jul 12, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is piperacillin and tazobactam?

Piperacillin and tazobactam is a combination penicillin antibiotic that is used to treat many different infections caused by bacteria, such as stomach infections, skin infections, pneumonia, and severe uterine infections.

piperacillin and tazobactam is sometimes given together with other antibiotics.

Piperacillin and tazobactam may also be used for purposes not listed in this medication guide.

Piperacillin and tazobactam side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes.

Piperacillin and tazobactam may cause serious side effects. Call your doctor at once if you have:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose);
- · confusion;
- muscle twitching or stiffness, trouble walking;
- a seizure;
- low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing; or
- **low potassium level**--leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling.

Common side effects of piperacillin and tazobactam may include:

- diarrhea, constipation;
- nausea;
- · headache; or
- sleep problems (insomnia).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Piperacillin / tazobactam side effects (more detail)

Related/similar drugs

Augmentin

Augmentin is a prescription antibiotic combining amoxicillin and clavulanate to treat bacterial ...

Reviews & ratings

5.4 / 10

447 Reviews

View more

Nitrofurantoin

Nitrofurantoin is an antibiotic used to treat urinary tract infections. Learn about side effects ...

Reviews & ratings

4.5 / 10

1,878 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Amoxicillin/clavulanate

Amoxicillin and clavulanate potassium is a combination antibiotic used to treat bacterial ...

Reviews & ratings

5.3 / 10

910 Reviews

View more

Bactrim

Bactrim (sulfamethoxazole and trimethoprim) is an antibiotic used to treat ear infections, urinary ...

Reviews & ratings

5.4 / 10

616 Reviews

View more

Levofloxacin

Levofloxacin is a fluoroquinolone antibiotic used to treat serious bacterial infections and prevent ...

Reviews & ratings

5.1 / 10

741 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Ciprofloxacin

Ciprofloxacin is an antibiotic belong to

a group or grugs caneu fluoroquinolones. Learn about side ...

Reviews & ratings

5.0 / 10

935 Reviews

View more

Warnings

You should not use piperacillin and tazobactam if you have ever had an allergic reaction to certain antibiotics (especially a penicillin or cephalosporin antibiotic).

Before taking this medicine

You should not use this medicine if you are allergic to:

- piperacillin or any other penicillin antibiotic (amoxicillin, ampicillin, Augmentin, dicloxacillin, oxacillin, penicillin, ticarcillin, or others);
- · tazobactam; or
- a cephalosporin antibiotic such as cefdinir (Omnicef), cephalexin (Keflex), or others.

Tell your doctor if you have ever had:

- kidney disease (or if you are on dialysis);
- a bleeding or blood clotting disorder;
- an electrolyte imbalance such as low levels of potassium in your blood;
- · cystic fibrosis;
- · any type of allergy; or
- · if you are on a low-salt diet.

Tell your doctor if you are pregnant or breastfeeding.

How should I use piperacillin and tazobactam?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Piperacillin and tazobactam is given as an infusion into a vein. A healthcare provider will give your first dose and may teach you how to properly use the medication by yourself.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you don't understand all instructions.

Prepare an injection only when you are ready to give it. Do not use if the medicine has changed colors or has

particles in it. Call your pharmacist for new medicine.

Use piperacillin and tazobactam for the full prescribed length of time, even if your symptoms quickly improve. Skipping doses can increase your risk of infection that is resistant to medication. Piperacillin and tazobactam will not treat a viral infection such as the flu or a common cold.

You may need frequent blood tests.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using piperacillin and tazobactam.

Carefully follow your healthcare provider's instructions about storing your medicine. You may need to keep the medicine in a refrigerator or at room temperature. How you store the medicine and for how long will depend on how the medicine was mixed at the pharmacy.

Use a needle and syringe only once and then place them in a puncture-proof "sharps" container. Follow state or local laws about how to dispose of this container. Keep it out of the reach of children and pets.

Detailed Piperacillin / tazobactam dosage information

What happens if I miss a dose?

Call your doctor for instructions if you miss a dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while using piperacillin and tazobactam?

Antibiotic medicines can cause diarrhea, which may be a sign of a new infection. If you have diarrhea that is watery or bloody, call your doctor before using anti-diarrhea medicine.

What other drugs will affect piperacillin and tazobactam?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Other drugs may affect piperacillin and tazobactam, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

1 Piperacillin / tazobactam drug interactions (more detail)

Does piperacillin/tazobactam interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

piperacillin/tazobactam

Enter a drug name Add

More about piperacillin / tazobactam

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (8)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: penicillins/beta-lactamase inhibitors
- En español

Patient resources

Piperacillin and tazobactam advanced reading

Other brands

Zosyn

Professional resources

- Piperacillin/Tazobactam monograph
- Piperacillin and Tazobactam (FDA)

Other brands

Zosyn

Related treatment guides

- Aspiration Pneumonia
- Bacteremia
- Appendicitis
- Appendectomy

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

Antibiotics 101

List of Common Antibiotics & Types

Antibiotics and Alcohol

Probiotics and Antibiotics

Antibiotics for Strep Throat

Antibiotics for a Tooth Infection

Antibiotics for UTI's

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

প্ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

6.9 / 10

8 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









 \times

Copyright © 2000-2025 Drugs.com. All rights reserved.