



[Home](#)

2. [Uribel](#)

Uribel

Generic name: [hyoscyamine](#), [methenamine](#), [methylene blue](#), [phenyl salicylate](#), [Sodium biphosphate monobasic](#) [*HYE-oh-SYE-a-meen*, *meth-EN-a-meen*, *METH-il-een-BLUE*, *FEEN-il-sa-LIS-il-ate*, *SOE-dee-um-bye-FOS-fate*]

Brand names: [Phosphasal](#), [urelle](#), [Uribel](#), [Urimar-T](#), [Urin D/S](#), ... [show all 14 brands](#)

Drug class: [Urinary antispasmodics](#)



[Medically reviewed](#) by Drugs.com on Oct 16, 2023. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is Uribel?

Hyoscyamine produces many effects in the body, including relief from muscle spasms.

Methenamine and [methylene blue](#) work as mild antiseptics that fight bacteria in the urine and bladder.

Phenyl salicylate is a mild pain reliever.

Sodium biphosphate monobasic is a form of phosphorus, which is a naturally occurring substance that is important in every cell in the body.

Uribel is a combination medicine used to treat bladder irritation (pain, spasms, inflammation) caused by [urinary tract infection](#). Uribel is also used to prevent bladder discomfort during a medical procedure.

Uribel may also be used for purposes not listed in this medication guide.

Side Effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have:

- severe dizziness, blurred vision, [fast heart rate](#);
- painful or difficult urination; or
- [shortness of breath](#).

Older adults may be more likely to have side effects from this medication.

Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm.

Common side effects of Uribel may include:

- dizziness, drowsiness;
- dry mouth; or
- nausea, vomiting.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Uribel side effects](#) (more detail)

Related/similar drugs

Augmentin

Augmentin is a prescription antibiotic combining amoxicillin and clavulanate to treat bacterial ...

Reviews & ratings

5.4 / 10

[447 Reviews](#)

[View more](#)

Bactrim

Bactrim (sulfamethoxazole and trimethoprim) is an antibiotic used to treat ear infections, urinary ...

Reviews & ratings

5.4 / 10

[616 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Macrobid

Macrobid (nitrofurantoin) is an antibiotic used to treat urinary tract infections. Includes side ...

Reviews & ratings

4.5 / 10

[827 Reviews](#)

[View more](#)

Nitrofurantoin

Nitrofurantoin is an antibiotic used to treat urinary tract infections. Learn about side effects ...

Reviews & ratings

4.5 / 10

[1,878 Reviews](#)

[View more](#)

Amoxicillin/clavulanate

Amoxicillin and clavulanate potassium is a combination antibiotic used to treat bacterial ...

Reviews & ratings

5.3 / 10

[910 Reviews](#)

[View more](#)

Levofloxacin

Levofloxacin is a fluoroquinolone antibiotic used to treat serious bacterial infections and prevent ...

Reviews & ratings

5.1 / 10

[741 Reviews](#)

[View more](#)

Ciprofloxacin

Ciprofloxacin is an antibiotic belong to a group of drugs called

a group of drugs called
fluoroquinolones. Learn about side ...

Reviews & ratings

5.0 / 10

[935 Reviews](#)

[View more](#)

Warnings

Many drugs can interact with methylene blue. **During your treatment with this medicine, do not start or stop using any other medications without your doctor's advice.**

If you take an antidepressant or psychiatric medication, call your doctor right away if you have signs of a serious drug interaction, including: confusion, memory problems, feeling hyperactive (mentally or physically), loss of coordination, [muscle twitching](#), shivering, sweating, diarrhea, and/or fever.

Before taking this medicine

You should not use this medicine if you are allergic to hyoscyamine, methenamine, methylene blue, phenyl salicylate, or Sodium biphosphate monobasic.

During your treatment with this medicine, do not start or stop using any other medications unless your doctor tells you to.

To make sure this medicine is safe for you, tell your doctor if you have:

- [heart disease](#);
- a heart rhythm disorder;
- congestive [heart failure](#);
- [coronary heart disease](#);
- a heart valve disorder;
- glaucoma;
- an [enlarged prostate](#) or bladder obstruction;
- [myasthenia gravis](#);
- an ulcer or obstruction in your stomach;
- an allergy to belladonna (Donnatal and others); or
- an allergy to salicylates (such as aspirin, [choline magnesium trisalicylate](#), [magnesium salicylate](#)).

It is not known whether this medication will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medicine.

This medicine can pass into breast milk and may harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

Do not give this medicine to a child without medical advice.

How should I take Uribel?

Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Drink plenty of liquids while you are taking this medicine.

Hyoscyamine can slow your digestion, and it may take longer for your body to absorb any medicines you take by mouth.

Store at room temperature away from moisture, heat, and light. Keep the bottle tightly closed when not in use.

 [Detailed Uribel dosage information](#)

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose.

Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Uribel?

Avoid taking an antacid or anti-diarrhea medicine within 1 hour before or after you take hyoscyamine, methenamine, methylene blue, phenyl salicylate, and Sodium biphosphate monobasic. Antacids or anti-diarrhea medicine can make it harder for your body to absorb hyoscyamine.

If you also take ketoconazole (Nizoral), wait at least 2 hours after taking it before you take hyoscyamine, methenamine, methylene blue, phenyl salicylate, and Sodium biphosphate monobasic.

What other drugs will affect Uribel?

Many drugs can interact with methylene blue. Tell your doctor about all other medications you use. You may need to stop using certain medicines (in some cases for up to 5 weeks before you start treatment with Uribel). **However, do not stop taking any of your medications without your doctor's advice.** This includes:

- any type of antidepressant;
- diet pills, stimulants, cold or allergy medicines, ADHD medication;
- narcotic pain medicine;
- medication to treat [Parkinson's disease](#) or restless leg syndrome; or
- a "triptan" [migraine headache](#) medication.

Taking any of the above medications while you are treated with medicine that contains methylene blue can cause **high**

levels of serotonin in the body. Call your doctor right away if you have symptoms of this condition: agitation, hallucinations, fever, fast heart rate, overactive reflexes, nausea, vomiting, diarrhea, loss of coordination, or fainting.

Many drugs can interact with this medicine. Not all possible interactions are listed here. Also, hyoscyamine can make it harder for your body to absorb other medications you take by mouth. **Tell your doctor about all your medications and any you start or stop using**, especially:

- a diuretic or "water pill";
- medicines that change the acidity of your urine--[sodium bicarbonate](#), [potassium citrate](#), sodium citrate with citric acid or potassium;
- bladder or urinary medicines--darifenacin, [fesoterodine](#), oxybutynin, tolterodine, solifenacin; or
- a sulfa drug.

This list is not complete and many other drugs can interact with Uribel. This includes prescription and over-the-counter medicines, vitamins, and [herbal products](#). Give a list of all your medicines to any healthcare provider who treats you.

 [Uribel drug interactions](#) (more detail)

Does Uribel interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Uribel

+

Enter a drug name

Add

More about Uribel (hyoscyamine / methenamine / methylene blue / phenyl salicylate / Sodium biphosphate monobasic)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(40\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [Drug class: urinary antispasmodics](#)
- [En español](#)

Patient resources

- [Uribel advanced reading](#)

Other brands

[Uro-MP](#), [Phosphasal](#), [Uretron DS](#), [Urimar-T](#), ... +15 more

Professional resources

- [Uribel prescribing information](#)

Other brands

[Phosphasal](#), [Uretron DS](#), [Urimar-T](#), [Urelle](#), ... +7 more

Related treatment guides

- [Urinary Tract Infection](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 3.02.

Antibiotics 101

[List of Common Antibiotics & Types](#)

[Antibiotics and Alcohol](#)

[Probiotics and Antibiotics](#)

[Antibiotics for Strep Throat](#)

[Antibiotics for a Tooth Infection](#)

[Antibiotics for UTI's](#)

DRUG STATUS

Availability	
Rx	Prescription only
CSA Schedule*	
N/A	Not a controlled drug



User Reviews & Ratings

6.3 / 10

[40 Reviews](#)

Images

Uribel hyoscyamine 0.12 mg / methenamine 118 mg/methylene blue 10 mg / phenyl salicylate 36 mg / sodium phosphate 40.8 mg (S 111)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.