

#### Home

ProAir RespiClick

# ProAir RespiClick

Generic name: albuterol inhalation [ al-BUE-ter-ol ]

Brand names: ProAir Digihaler, ProAir RespiClick, Ventolin HFA

Drug class: Adrenergic bronchodilators



Medically reviewed by Drugs.com on Aug 12, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

# What is ProAir RespiClick?

Albuterol **inhalers** are used in adults and children at least 4 years old to to treat or prevent narrowing and swelling inside the lungs (bronchospasm) that may cause breathing problems. Albuterol **inhalers** are also used to prevent breathing problems while exercising.

ProAir RespiClick solution is used to treat acute bronchospasm attacks in children at least 2 years old with asthma.

ProAir RespiClick may also be used for purposes not listed in this medication guide.

# ProAir RespiClick side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

**ProAir RespiClick may increase the risk of asthma-related death or hospitalization.** Use only the prescribed dose and follow all instructions for safe use.

ProAir RespiClick may cause serious side effects. Call your doctor at once if you have:

- wheezing, choking, or other breathing problems after using ProAir RespiClick;
- · chest pain, fast heart rate, pounding heartbeats or fluttering in your chest;
- severe headache, pounding in your neck or ears;
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor; or
- **low blood potassium**--leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling.

#### Common side effects of ProAir RespiClick may include:

· chest pain, fast or pounding heartbeats;

- pain and burning when you urinate;
- upset stomach, vomiting;
- · dizziness;
- tremors or shaking, nervousness;
- headache, back pain, body aches; or
- cough, sore throat, sinus pain, runny or stuffy nose.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

ProAir RespiClick side effects (more detail)

# Related/similar drugs

## **Breztri Aerosphere**

Breztri (budesonide/glycopyrrolate/formoterol fumarate) is a combination inhaler that may be used ...

#### **Reviews & ratings**

6.9 / 10

119 Reviews

#### View more

#### **Xolair**

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

## **Reviews & ratings**

7.5 / 10

229 Reviews

#### View more

#### FEATURED

## **Dupixent**

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

#### **Reviews & ratings**

7.2 / 10

. . . \_ .

#### View more

# Ciprofloxacin

Ciprofloxacin is an antibiotic belong to a group of drugs called fluoroquinolones. Learn about side ...

#### **Reviews & ratings**

5.0 / 10

935 Reviews

View more

# Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

#### **Reviews & ratings**

7.1 / 10

484 Reviews

View more

## **Azithromycin**

Azithromycin is an antibiotic used to treat many different types of infections caused by bacteria ...

## **Reviews & ratings**

6.8 / 10

1,306 Reviews

View more

## **Montelukast**

Montelukast is a daily oral medication used to prevent asthma attacks, exercise-induced ...

## **Reviews & ratings**

5.6 / 10

571 Reviews

View more

## **Doxycycline**

Doxycycline is a tetracycline antibiotic used to treat many different bacterial infections such as ...

#### **Reviews & ratings**

6.4 / 10

1,635 Reviews

View more

#### **Amoxicillin**

Amoxicillin is a penicillin antibiotic that fights bacteria. It is used to treat many types of ...

## **Reviews & ratings**

6.4 / 10

366 Reviews

View more

## **Augmentin**

Augmentin is a prescription antibiotic combining amoxicillin and clavulanate to treat bacterial ...

#### **Reviews & ratings**

5.4 / 10

447 Reviews

View more

# **Warnings**

Use only as directed. Tell your doctor if you use other medicines or have other medical conditions or allergies.

# Before taking this medicine

You should not use ProAir RespiClick if you are allergic to ProAir RespiClick.

Do not use albuterol **inhalation powder** (**ProAir RespiClick or ProAir Digihaler**) if you are allergic to lactose or milk proteins.

Tell your doctor if you have ever had:

- heart problems, high blood pressure;
- · a thyroid disorder;
- · seizures;
- if you have used an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, or tranylcypromine;
- · diabetes; or
- low blood levels of potassium.

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of albuterol inhalation on the baby.

Tell your doctor if you are pregnant or plan to become pregnant. It is not known whether ProAir RespiClick will harm an unborn baby. However, having uncontrolled asthma during pregnancy may increase the risk of premature birth, low birth weight, or eclampsia (dangerously high blood pressure that can lead to medical problems in both mother and baby). The benefit of preventing bronchospasm may outweigh any risks to the baby.

Do not allow a young child to use ProAir RespiClick without help from an adult.

# How should I use ProAir RespiClick?

Follow the directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed. **Do not use the medicine more often than prescribed.** 

Your dose needs may change if you switch to a different brand, strength, or form of this medicine. **Avoid medication** errors by using exactly as directed on the label, or as prescribed by your doctor.

The effects of albuterol inhalation last about 4 to 6 hours. To prevent exercise-induced bronchospasm, use this medicine 15 to 30 minutes before you exercise. Talk with your doctor if you need to time your dosing around any planned activities.

Read and carefully follow any instructions for preparing and using the medicine for the first time and for following doses. Ask your doctor or pharmacist if you do not understand these instructions.

Your dose needs may change due to surgery, illness, stress, or a recent asthma attack. Do not change your dose or stop using asthma medication without your doctor's advice. **Tell your doctor if your medicine seems to stop working.** 

Call your doctor if your symptoms do not improve, or if they get worse.

Keep the cover on your inhaler closed when not in use. Store away from open flame or high heat. Do not puncture or burn an empty canister.

Do not take apart an inhaler device, wash, or put any part of your inhaler in water. Follow all storage and cleaning instructions provided with your inhaler. Your pharmacist can provide more information about how to store this medicine.

Get your prescription refilled before you run out of medicine completely. Always use the new inhaler device provided.

Store unused albuterol solution vials in the foil pouch at room temperature away from moisture, heat and light. Call

your pharmacist if the medicine looks cloudy, has changed colors or has particles in it.

**ProAir RespiClick solution is stable for only a certain number of days or weeks** after being removed from the foil pouch. Throw away any medicine not used within that time. Talk to your pharmacist if you have any questions.

Detailed ProAir RespiClick dosage information

# What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

# What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of albuterol can be fatal.** 

Overdose symptoms may include dry mouth, tremors, chest pain, fast heartbeats, nausea, general ill feeling, seizure, feeling light-headed or fainting.

# What should I avoid while using ProAir RespiClick?

Avoid using other fast-acting rescue medications or asthma medications with ProAir RespiClick unless you doctor tells you to.

Avoid getting ProAir RespiClick in your eyes.

# What other drugs will affect ProAir RespiClick?

Tell your doctor about all your other medicines, especially:

- any other inhaled medicines or bronchodilators such as ipratropium o tiotropium;
- digoxin;
- · epinephrine;
- a diuretic ("water pill")
- an antidepressant--amitriptyline, desipramine, imipramine, doxepin, nortriptyline, and others;
- a beta blocker--atenolol, carvedilol, labetalol, metoprolol, nadolol, nebivolol, propranolol, sotalol, and others; or
- an MAO inhibitor--isocarboxazid, linezolid, methylene blue injection, phenelzine, tranylcypromine, and others.

This list is not complete. Other drugs may affect ProAir RespiClick, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 ProAir RespiClick drug interactions (more detail)

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

ProAir RespiClick	
+	
Enter a drug name	Add

# **Popular FAQ**

ProAir Digihaler vs Respiclick, what's the difference?	~
Can asthma inhalers cause a sore throat?	~
Can you use an expired albuterol inhaler?	<b>\</b>

## **More FAQ**

- How do you use the ProAir Digihaler?
- Does either Ventolin or albuterol contain steroids?
- Does coffee help with asthma?
- What is albuterol sulfate and can I take it if I'm allergic to sulfa?

# More about ProAir RespiClick (albuterol)

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (108)
- Latest FDA alerts (5)
- Side effects
- Dosage information
- During pregnancy
- FDA approval history
- Drug class: adrenergic bronchodilators
- Breastfeeding

## **Patient resources**

· Proair Respiclick advanced reading

#### Other brands

Ventolin HFA, Ventolin, ProAir HFA, Proventil, ... +5 more

## **Professional resources**

- ProAir Respiclick prescribing information
- Albuterol Sulfate, Levalbuterol Hydrochloride, Levalbuterol Tartrate (Systemic, Oral Inhalation) (AHFS Monograph)

## Other brands

Ventolin, ProAir HFA, Proventil HFA, ProAir Digihaler

# Related treatment guides

- Bronchiectasis
- Bronchospasm Prophylaxis
- · Asthma, acute
- Asthma, Maintenance
- · COPD, Acute
- · COPD, Maintenance

# **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.02.

#### **DRUG STATUS**

# Availability Rx Prescription only Pregnancy & Lactation Risk data available CSA Schedule\* N/A Not a controlled drug Approval History Drug history at FDA

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com

Advertising policy

Attribution & citations

#### **Terms & privacy**

Terms of use

**Editorial policy** 

Privacy policy

#### **Support**

Help center

**Sitemap** 

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.