

[Home](#)

2. Aspirin/carisoprodol/codeine

Aspirin, carisoprodol, and codeine

Generic name: aspirin, carisoprodol, and codeine [*AS-pir-in, kar-EYE-soe-PROE-dol, and-KOE-deen*]**Brand name:** [Soma Compound with Codeine](#)**Dosage form:** oral tablet (325 mg-200 mg-16 mg)**Drug class:** [Skeletal muscle relaxant combinations](#)[Medically reviewed](#) by Drugs.com on Mar 31, 2025. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is aspirin, carisoprodol, and codeine

Aspirin, [carisoprodol](#), and codeine is a combination medicine used together with rest and physical therapy to treat pain, muscle spasm, and other symptoms related to injuries and other painful muscular conditions.

aspirin, carisoprodol, and codeine contains codeine, an opioid (narcotic) medicine, and may be habit-forming.

This medication may also be used for purposes not listed in this medication guide.

What are the possible side effects of aspirin, carisoprodol, and codeine

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up.

Aspirin may cause stomach or intestinal bleeding, which can be fatal. This can occur without warning while you are taking this medicine.

Aspirin, carisoprodol, and codeine may cause serious side effects. Call your doctor at once if you have:

- noisy breathing, sighing, shallow breathing;
- a [slow heart rate](#) or weak pulse;
- a light-headed feeling, like you might pass out;
- confusion, unusual thoughts or behavior;
- easy bruising or bleeding (nosebleeds, bleeding gums);
- severe [constipation](#);

- **symptoms of stomach bleeding**--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds; or
- **low cortisol levels**--nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, [fast heart rate](#), muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Serious side effects may be more likely in older adults and those who are overweight, malnourished, or debilitated.

Common side effects of aspirin, carisoprodol, and codeine may include:

- dizziness, drowsiness, tiredness;
- nausea, vomiting, stomach pain, constipation; or
- headache.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Aspirin / carisoprodol / codeine side effects](#) (more detail)

Related/similar drugs

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

5.8 / 10

[30 Reviews](#)

[View more](#)

Naproxen

Naproxen is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

[683 Reviews](#)

[View more](#)

FEATURED

Ozempic

[Learn about Ozempic \(semaglutide\)](#)

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

[1,404 Reviews](#)

[View more](#)

Tramadol

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

Reviews & ratings

7.0 / 10

[1,876 Reviews](#)

[View more](#)

Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

Reviews & ratings

6.0 / 10

[618 Reviews](#)

[View more](#)

Meloxicam

Meloxicam is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.3 / 10

[600 Reviews](#)

[View more](#)

Aspirin

Aspirin is used to treat mild to moderate pain and to reduce fever or inflammation. Learn about ...

Reviews & ratings

7.8 / 10

[51 Reviews](#)

[View more](#)

Ibuprofen

Ibuprofen (Advil, Motrin) treats minor aches and pains caused by the common cold, headaches ...

Reviews & ratings

7.3 / 10

[240 Reviews](#)

[View more](#)

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-the-counter painkiller and fever ...

Reviews & ratings

6.1 / 10

[13 Reviews](#)

[View more](#)

Warnings

MISUSE OF OPIOID MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medication in a place where others cannot get to it.

Do not give this medicine to anyone younger than 12 years old, or anyone under 18 who recently had surgery to remove

the tonsils or adenoids.

Taking this medicine during pregnancy may cause life-threatening withdrawal symptoms in the newborn.

Fatal side effects can occur if you use opioid medicine with alcohol, or with other drugs that cause drowsiness or slow your breathing.

Before taking this medicine

You should not use aspirin, carisoprodol, and codeine if you are allergic to aspirin, carisoprodol, or codeine, or if you have:

- severe asthma, asthma with runny nose and nasal polyps, or other breathing problems;
- a stomach or [bowel obstruction](#) (including [paralytic ileus](#));
- an allergy to meprobamate or NSAIDs ([nonsteroidal anti-inflammatory drugs](#));
- porphyria (a genetic enzyme disorder that causes symptoms affecting the skin or nervous system);
- hemophilia; or
- if you have used an MAO inhibitor in the past 14 days (such as [isocarboxazid](#), linezolid, [methylene blue](#) injection, phenelzine, rasagiline, selegiline, or [tranylcypromine](#)).

Aspirin, carisoprodol, and codeine is **not approved** for use by anyone younger than 12 years old.

Do not give this medicine to anyone younger than 18 years old who recently had surgery to remove the tonsils or adenoids.

Tell your doctor if you have ever had:

- sleep apnea or other breathing disorders;
- a [stomach ulcer](#) or bowel obstruction;
- a [vitamin K deficiency](#);
- seizures, head injury, or brain tumor;
- liver or kidney disease;
- alcoholism or a [drug addiction](#);
- problems with your pancreas, thyroid, or gallbladder.
- urination problems; or
- if you take medicine to treat or prevent blood clots (such as warfarin, enoxaparin, clopidogrel, rivaroxaban, Coumadin, Effient, or Xarelto).

Do not give this medicine to a teenager with a fever, flu symptoms, or chickenpox. **Aspirin can cause Reye's syndrome, a serious and sometimes fatal condition in adolescents.**

If you use opioid medicine while you are pregnant, your baby could become dependent on the drug. This can cause life-threatening withdrawal symptoms in the baby after it is born. Babies born dependent on opioids may need medical treatment for several weeks.

Taking aspirin during late pregnancy may cause bleeding in the mother or the baby during delivery.

Do not breast-feed. Codeine can pass into breast milk and may cause drowsiness, breathing problems, or death in a nursing baby.

How should I take aspirin, carisoprodol, and codeine

Follow the directions on your prescription label and read all medication guides. **Use the lowest dose that is effective in treating your condition.** Tell your doctor if you feel an increased urge to take more of this medicine.

Never share this medicine with another person, especially someone with a history of drug abuse or addiction. MISUSE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medication in a place where others cannot get to it. Selling or giving away opioid medicine is against the law.

If you need surgery or medical tests, tell the doctor ahead of time that you are using this medicine.

Aspirin, carisoprodol, and codeine is for short term use only. Call your doctor if your symptoms do not improve after 2 to 3 weeks.

Do not stop using this medicine suddenly after long-term use, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using the medicine.

Store at room temperature away from moisture and heat. Keep track of your medicine. You should be aware if anyone is using it improperly or without a prescription.

Do not keep leftover opioid medication. **Just one dose can cause death in someone using this medicine accidentally or improperly.** Ask your pharmacist where to locate a drug take-back disposal program. If there is no take-back program, flush the unused medicine down the toilet.

Aspirin, carisoprodol, and codeine dosing information

Usual Adult Dose for Muscle Spasm:

1 to 2 tablets 4 times a day

Maximum daily dose: 8 tablets/day (carisoprodol: 1600 mg; aspirin: 2600 mg; codeine: 128 mg)

Duration of therapy: Up to 2 or 3 weeks

Comments:

-Initiate therapy, taking into account severity of pain, response, prior analgesic treatment experience, and risk factors for addiction, abuse, and misuse

-Use should be limited to short periods (up to 3 weeks) since acute, painful musculoskeletal conditions are generally of short duration and evidence of effectiveness beyond this time period has not been established.

Use: For the relief of discomfort associated with acute, painful musculoskeletal conditions.

Usual Adult Dose for Pain:

1 to 2 tablets 4 times a day

Maximum daily dose: 8 tablets/day (carisoprodol: 1600 mg; aspirin: 2600 mg; codeine: 128 mg)

Duration of therapy: Up to 2 or 3 weeks

Comments:

-Initiate therapy, taking into account severity of pain, response, prior analgesic treatment experience, and risk factors for addiction, abuse, and misuse

-Use should be limited to short periods (up to 3 weeks) since acute, painful musculoskeletal conditions are generally of short duration and evidence of effectiveness beyond this time period has not been established.

Use: For the relief of discomfort associated with acute, painful musculoskeletal conditions.

 [Detailed Aspirin / carisoprodol / codeine dosage information](#)

What happens if I miss a dose?

Since aspirin, carisoprodol, and codeine is used for pain, you are not likely to miss a dose. Skip any missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of this medicine can be fatal, especially in a child or another person using the medicine without a prescription.**

Overdose symptoms may include extreme drowsiness, confusion, muscle weakness or stiffness, slow heartbeats, dilated or pinpoint pupils, vision problems, ringing in your ears, muffled hearing, hallucinations, feeling hot, vomiting, fainting, blue lips, cold and clammy skin, rapid or very slow breathing, seizure, or coma.

What should I avoid while taking aspirin, carisoprodol, and codeine

Avoid driving or operating machinery until you know how aspirin, carisoprodol, and codeine will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Do not drink alcohol. Dangerous side effects or death could occur.

Ask a doctor or pharmacist before using other medicines for pain, fever, swelling, or cold/flu symptoms. They may contain ingredients similar to aspirin (such as ibuprofen, ketoprofen, or naproxen).

What other drugs will affect aspirin, carisoprodol, and codeine

You may have breathing problems or withdrawal symptoms if you start or stop taking certain other medicines.

Tell your doctor if you also use an antibiotic, antifungal medication, heart or blood pressure medication, seizure medication, or medicine to treat HIV or hepatitis C.

Opioid medication can interact with many other drugs and cause dangerous side effects or death. Be sure your doctor knows if you also use:

- cold or allergy medicines, bronchodilator asthma/COPD medication, or a diuretic ("water pill");
- medicines for [motion sickness](#), [irritable bowel syndrome](#), or [overactive bladder](#);
- **other narcotic medications**--opioid pain medicine or prescription cough medicine;

- **a sedative like Valium**--diazepam, alprazolam, lorazepam, Xanax, Klonopin, Versed, and others;
- **drugs that make you sleepy or slow your breathing**--a sleeping pill, muscle relaxer, medicine to treat mood disorders or mental illness; or
- **drugs that affect serotonin levels in your body**--a stimulant, or medicine for depression, [Parkinson's disease](#), migraine headaches, serious infections, or nausea and vomiting.

This list is not complete. Other drugs may affect aspirin, carisoprodol, and codeine, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Not all possible interactions are listed here.**

i [Aspirin / carisoprodol / codeine drug interactions](#) (more detail)

Does aspirin/carisoprodol/codeine interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

aspirin/carisoprodol/codeine

+

Enter a drug name

Add

More about aspirin / carisoprodol / codeine

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(6\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: skeletal muscle relaxant combinations](#)
- [En español](#)

Patient resources

- [Carisoprodol, aspirin, and codeine advanced reading](#)

Professional resources

- [Carisoprodol, Aspirin, Codeine prescribing information](#)

Other brands

[Soma Compound with Codeine](#)

Related treatment guides

- [Muscle Spasm](#)
- [Pain](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.01.

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
3	Moderate abuse potential
Approval History	
	Drug history at FDA

User Reviews & Ratings

[6 Reviews](#)

Images

[Aspirin, Carisoprodol and Codeine 325 mg / 200 mg / 16 mg \(A 138\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.